Parallel Computation

1 Computational Operations

1.1 Set of Canonical Functions

Define $\mathbb{C}$; the set of canonical functions

$$\mathbb{C} = \{I, \neg, \vee, \wedge, ==, >, <, \cup, \cap, ||, ...\}$$

1.2 Definition of Assignment

Define assignment $\leftarrow$

$$a \leftarrow b \ = \leftarrow [a, b] :=$$
$$a = \varnothing \cup b$$

1.3 Definition of Canonical Instruction

Define canonical instruction c as the assignment of a canonical program $l[X_n]$ to element a

$$c := a \leftarrow l[X_n]$$

1.4 Definition of Computational Instruction

Define computational instruction s as a set of canonical instructions with output a_{2M-1}

$$X_n = \{x_1, x_2, ..., x_n\};$$

$$s[X_n] := \{c_1, c_3, c_5, ..., c_{2M-1}\} \rightarrow a_1 \mid$$
$$c_{2i-1} = a_i \leftarrow l_i[\hat{X}_i] \quad l_i \in \mathbb{C} \quad \forall i \leqslant M$$

$$= \{\{a_1 \leftarrow l[\hat{X}_1]\}, \{a_2 \leftarrow l[\hat{X}_2]\}, ..., \{a_M \leftarrow l[\hat{X}_M]\}\} \rightarrow a_M \mid$$
$$l_i \in \mathbb{C} \quad \forall i$$

2 Computational Instructions

2.1 Set Union $\cup$

2.1.1 Canonical Union $\cup$

Restate the definion of Union $\cup$ for elements $a; b$

$$a \cup b = \cup[a, b] \quad a \in \Omega \;\; b \in \Omega :=$$

$a = b \neq \varnothing$

$$a \cup a = b \cup b \rightarrow \{a\} = \{b\}$$

$\varnothing \neq a \neq b \neq \varnothing$

$$a \cup b \rightarrow \{a, b\}$$

$\varnothing = a \neq b$

$$\varnothing \cup b \rightarrow \{b\}$$

$a \neq b = \varnothing$

$$a \cup \varnothing \rightarrow \{a\}$$

$a = b = \varnothing$

$$\varnothing \cup \varnothing \rightarrow \varnothing$$

2.1.2 Domain of Canonical Union $\cup$

$$\{a \mid a \in \Omega\}$$
$$\{b \mid b \in \Omega\}$$

2.1.3 Set Theory Definition of Set Union $\cup$

Define the overloaded symbol $\cup$; Set Union

$$A = \{a_1, a_2, ..., a_N\} : a_i \in \Omega \quad \forall a_i \in A$$
$$B = \{b_1, b_2, ..., b_M\} : b_j \in \Omega \quad \forall b_j \in A$$

$$A \cup B := \tilde{A} = \{\tilde{a}_1, \tilde{a}_2, ..., \tilde{a}_K\} :$$
$$\tilde{a}_i \in A \vee \tilde{a}_i \in B \quad \forall \tilde{a}_i \in \tilde{A}$$

2.1.4 Domain Set Union

$$\{A \mid A \subseteq \Omega\}$$
$$\{B \mid B \subseteq \Omega\}$$

2.1.5 Overloaded $\cup$

Union is defined distinctly according to the domain of inputs

2.1.6 Computational Defintion of Set Union $\cup$

2.2 Set Minus $\setminus$

2.2.1 Formal Definition

2.2.2 Computational Defintion of Set Minus $\setminus$

2.2.3 Domain

2.3 Set is equal to $==$

2.3.1 Formal Definition

2.3.2 Computational Defintion of Set is equal to $==$

2.3.3 Domain

2.4 Plus $+$

2.4.1 Formal Definition

$$N + M :=$$
$$|\{a_1, a_2, ..., a_N\} \cup \{b_1, b_2, ..., b_M\}|$$
$$= |a_1 \cup a_2 \cup ... \cup a_N \cup b_1 \cup b_2 ... \cup b_M|$$

Let $\hat{a}_i = a_i \quad \forall i \leqslant N; \quad \hat{a}_{N+j} = b_j \quad \forall j \leqslant M;$

$$= |\{\hat{a}_1\} \cup \{\hat{a}_2\} \cup ... \cup \{\hat{a}_N\} \cup \{\hat{a}_{N+1}\} \cup \{\hat{a}_{N+2}\} \cup ... \cup \{\hat{a}_{N+M}\}|$$

2.4.2 Computational Instruction Definition of Plus $+$

$$s[X_n] := \{c_1, c_3, c_5, ..., c_{2M-1}\} \rightarrow a_1 \mid$$
$$c_{2i-1} = a_i \leftarrow l_i[\hat{X}_i] \quad l_i \in \mathbb{C} \quad \forall i \leqslant M$$

$$+[N, M] :=$$
$$c_1 = a_1 \leftarrow I_1[1]$$
$$c_3 = a_2 \leftarrow I_2[1]$$
$$c_5 = a_1 \leftarrow a_1 \cup a_2$$
$$c_7 = a_3 \leftarrow I_3[1]$$
$$c_9 = a_1 \leftarrow a_1 \cup a_3$$
$$...$$
$$c_{4(M+N-2)+3} = a_{M+N} \leftarrow I_{M+N}[1]$$
$$c_{4(M+N-2)+5} = a_1 \leftarrow a_1 \cup a_{M+N}$$
$$c_{4(M+N-1)+3} = a_1 \leftarrow |a_1|$$

2.4.3 Domain

$$\{N \mid N \in \mathbb{R}\}$$
$$\{M \mid M \in \mathbb{R}\}$$

2.5 Minus -

2.5.1 Formal Definition

$$N - M := |\{a_1 \cup a_2 \cup ... \cup a_N\} \backslash \{a_1 \cup a_2 \cup ... \cup a_M\}|$$
$$= |a_1 \cup a_2 \cup ... \cup a_N \setminus a_1 \setminus a_2 ... \setminus a_M|$$
$$= |a_{M+1} \cup a_{M+2} \cup ... \cup a_N|$$

Let $\hat{a}_i = a_{M+i} \quad M < i \leqslant N$

$$= |\hat{a}_1 \cup \hat{a}_2 \cup ... \cup \hat{a}_{N-M}|$$

2.5.2 Computational Instruction Definition of Minus -

$$s[X_n] := \{c_1, c_3, c_5, ..., c_{2M-1}\} \rightarrow a_1 \mid$$
$$c_{2i-1} = a_i \leftarrow l_i[\hat{X}_i] \quad l_i \in \mathbb{C} \quad \forall i \leqslant M$$

$$-[N, M] :=$$

$$c_1 = a_1 \leftarrow I_1[1]$$
$$c_3 = a_2 \leftarrow I_2[1]$$
$$c_5 = a_1 \leftarrow a_1 \cup a_2$$
$$c_7 = a_3 \leftarrow I_3[1]$$
$$c_9 = a_1 \leftarrow a_1 \cup a_3$$
$$...$$
$$c_{4(N-2)+3} = a_N \leftarrow I_N[1]$$
$$c_{4(N-2)+5} = a_1 \leftarrow a_1 \cup a_N$$
$$c_{4(N-1)+3} = a_{N+1} \leftarrow I_1[1]$$
$$c_{4(N-1)+5} = a_1 \leftarrow a_1 \setminus a_{N+1}$$
$$c_{4N+3} = a_{N+2} \leftarrow I_2[1]$$
$$c_{4N+5} = a_1 \leftarrow a_1 \setminus a_{N+2}$$
$$...$$
$$c_{4(M+N-2)+3} = a_M \leftarrow I_M[1]$$
$$c_{4(M+N-2)+5} = a_1 \leftarrow a_1 \setminus a_M$$
$$c_{4(M+N-1)+3} = a_1 \leftarrow |a_1|$$

2.5.3 Domain

$$\{a \mid a \in \mathbb{R}\}$$
$$\{b \mid b \in \mathbb{R} \ \wedge b \leqslant a\}$$

3 Compound Instruction Theorem

Any Instruction that can be defined as a set of computational instructions "Compound Instruction" is also a computational instruction

3.1 Definition of Compound Instruction

3.2 $! =$

3.3 $\geqslant$

3.4 $\leqslant$

3.5 $\sum$

3.6 Set Intersection $\cap$

3.6.1 Formal Definition

3.6.2 Computational Defintion of $\cap$

3.6.3 Domain

$$\{A \mid A \in \Omega\}$$
$$\{B \mid B \in \Omega\}$$

3.7 *

3.8 Exponentiation a^b

3.9 /

3.9.1 Definition

3.9.2 Prove / is a Computational Instruction

3.10 N^{th} root $\sqrt[n]{a}$

3.10.1 Definition

3.10.2 Prove $\sqrt[n]{a}$ is a Computational Instruction

4 Programs

4.1 Instructions

Define $\mathcal{L}$; an ordered set of computational instructions s_i

$$\mathcal{L} := \{s_1, s_2, ..., s_N\}$$

4.2 Memory

Define Memory $\mathcal{M}$; an ordered set of elements s_i

$$\mathcal{M} := \{b_1, b_2, ..., b_M\}$$

4.3 Program

Define Program; an ordered set of operations and memory

$$P := \{s_1, s_2, ..., s_N, b_1, b_2, ..., b_M\}$$

4.3.1 State Variable Notation

It is convention to use | to separate instructions and memory

$$P := \{s_1, s_2, ..., s_N, b_1, b_2, ..., b_M\}$$
$$= \{s_1, s_2, ..., s_N | b_1, b_2, ..., b_M\}$$

4.4 Boolean Programs

Define a boolean program; a program with inputs x_i and boolean output y_o

$$X_n = \{x_1, ..., x_n\}$$
$$P_{boolean} = P[X_n] := \{s_1, s_2, ..., s_N \mid b_1, b_2, ..., b_M, y_o\} : y_o \in \{\mathbb{T}, \mathbb{F}\} \ \ \forall X_n$$

4.4.1 Function Notation

$$P_{boolean} = P[X_n] \rightarrow y_o : y_o \in \{\mathbb{T}, \mathbb{F}\} \ \ \forall X_n$$

4.5 Void Programs

Define a void program; a program with inputs x_i and no output

$$X_n = \{x_1, ..., x_n\}$$
$$P = P[X] := \{s_1, s_2, ..., s_N \mid b_1, b_2, ..., b_M\}$$

4.6 Numerical Programs

Define a numerical program; a program with inputs x_i, input set C, and real, rational output y_o

$$X = \{x_1, ..., x_n, C\}$$
$$P = P[X] := \{s_1, s_2, ..., s_N \mid b_1, b_2, ..., b_M, y_o\} =$$
$$P[X] \rightarrow y_o \in \mathbb{Q} \ \ y_o \geqslant 0$$

4.7 System Programs

Define a system program; a program with inputs x_i, input set C, and real, output set Y_o

$$X = \{x_1, ..., x_n, C\}$$
$$P = P[X] := \{s_1, s_2, ..., s_N \mid b_1, b_2, ..., b_M, Y_o\} =$$
$$P[X] \rightarrow Y_o = \{y_1, y_2, ..., y_K\}$$

4.8 Mathematical Programs

Define a mathematical program; a program with inputs x_i, input set C and numerical output y_o

$$X = \{x_1, ..., x_n, C\}$$
$$P = P[X] := \{s_1, s_2, ..., s_N \mid b_1, b_2, ..., b_M, y_o\} =$$
$$P[X] \rightarrow y_o \in \mathbb{Q}$$

4.9 Instruction Memory Notation

$$X_n = \{x_1, ..., x_n\};$$
$$P[X_n] \rightarrow y_o = \{s_1, s_2, ..., s_{O_T[n]}, b_1, b_2, ..., b_{O_S[n]}, y_o\} \rightarrow y_o$$
$$\mathcal{L} := \{s_1, s_2, ..., s_{O_T[n]}\}$$
$$\mathcal{M} := \{b_1, b_2, ..., b_{O_S[n]}\}$$
$$P[X_n] = \{\mathcal{L}, \mathcal{M}, y_o\}$$

5 Problem Definition

Also denoted as a "Question"

$$X_i = \{x_1, ..., x_n\}$$
$$Q := f[X_i] = Y_o \subseteq \Omega \quad \forall X_i$$

5.1 Set of Questions

Define $\mathbb{Q}$; the set of questions

$$\mathbb{Q} := \{Q_1, Q_2, ...\} :$$
$$Q_i = f[X_j] = Y_o \subseteq \Omega \ \ \forall X_j, i$$

5.2 Decision Questions / Decision Problems

5.2.1 Definition

Define decision problem; a function with inputs x_i and boolean output "answer" a_o

$$X_i = \{x_1, ..., x_n\}$$
$$D := f[X_i] = a_o \in \{\mathbb{T}, \mathbb{F}\} \quad \forall X_i$$

5.3 Numerical Questions / Numerical Problems

5.3.1 Definition

Define numerical problem; a function with inputs x_i and numerical output y_o

$$X_i = \{x_1, ..., x_n\}$$
$$Q := f[X_i] = y_o \in \mathbb{R} \quad \forall X_i$$

5.4 System Questions / System Problems

5.4.1 Definition

Define system problem; a function with inputs x_i and outputs y_j

$$X_i = \{x_1, ..., x_n\}$$
$$Q := f[X_i] = Y_o = \{y_1, ..., y_m\} \quad \forall X_i$$

6 Solutions

6.1 Definition

Program P is a solution s^+ to decision problem D if

1. P outputs answer a_o for all inputs $X_i \;\; \forall i$
and
2. $s^+[X_i]$ is a subset of $s^+[\hat{X}_i]$

$$X_i = \{x_1, ..., x_n, C\}; \;\; \hat{X}_i = \{x_1, ..., x_n, x_{n+1}, C\}$$
$$D := f[X_i] \rightarrow a_o \in \{\mathbb{T}, \mathbb{F}\} \;\; \forall X_i$$
$$s^+ = s^+[X_i] := P :$$
$$(P[X_i] \rightarrow y_o == a_o \;\; \forall X_i) \;\; \cap \;\; (P[\hat{X}_i] \supseteq P[X_i] \;\; \forall X_i, \hat{X}_i)$$
$$P[X_i] = \{s_1, s_2, ..., s_N | b_1, b_2, ..., b_M, y_o\}$$
$$s^+ = P[X_i] = \{s_1, s_2, ..., s_{O_T[n]}, b_1, b_2, ..., b_{O_S[n]}, y_o\} \;\; \forall X_i$$

6.1.1 Property of No-op ;

No-op ; can be added to any solution S_i without modifying the output y_o or memory b_i

$$; := \emptyset$$
$$s^+ = \{s_1, s_2, ..., s_{O_T[n]}, b_1, b_2, ..., b_{O_S[n]}, y_o\}$$
$$\hat{s}^+ = \{s_1, s_2, ..., \; ; \; , ..., s_{O_T[n]+1}, \hat{b}_1, \hat{b}_2, ..., \hat{b}_{O_S[n]}, \hat{y}_o\}$$
$$\hat{y}_o = y_o \;\; \forall k$$

6.2 Definition of S^+

Define S^+; the set of solutions to decision problem D

$$X_i = \{x_1, ..., x_n, C\}; \;\; \hat{X}_i = \{x_1, ..., x_n, x_{n+1}, C\}$$
$$D := f[X_i] \rightarrow a_o \in \{\mathbb{T}, \mathbb{F}\} \;\; \forall X_i$$
$$s_j^+ = s_j^+[X_i] := P :$$
$$(P[X_i] \rightarrow y_o == a_o \;\; \forall X_i) \;\; \cap \;\; (P[\hat{X}_i] \supseteq P[X_i] \;\; \forall X_i, \hat{X}_i)$$
$$S^+ := \{s_j^+, ...\} \;\; \forall j$$

6.3 Definition of Solvable

Define solvable

$$X_i = \{x_1, ..., x_n, C\}; \;\; \hat{X}_i = \{x_1, ..., x_n, x_{n+1}, C\}$$
$$D := f[X_i] \rightarrow a_o \in \{\mathbb{T}, \mathbb{F}\} \quad \forall X_i$$
$$solvable = solvable[D] = b_o \in \{\mathbb{T}, \mathbb{F}\} :=$$
$$\exists P : (P[X_i] \rightarrow y_o == a_o \quad \forall X_i) \;\; \cap \;\; (P[\hat{X}_i] \supseteq P[X_i] \quad \forall X_i, \hat{X}_i)$$

7 The set of all Decision Problems $\mathbb{D}$

7.1 Definition

Define the set of decision problems $\mathbb{D}$

$$X_i = \{x_1, ..., x_n, C\}$$
$$D_j := f_j[X_i] \rightarrow a_o \in \{\mathbb{T}, \mathbb{F}\} \quad \forall X_i$$
$$\mathbb{D} := \{D_j, ...\} \quad \forall j$$

8 Complexity

8.1 Time Complexity of a Decision Problem $O_T[n]$

Define Time Complexity $O_T[n]$ of solution s^+ to Decision Problem D as the total number of logical operations

$$X_i = \{x_1, ..., x_n, C\}; \;\; \hat{X}_i = \{x_1, ..., x_{n+1}, C\}$$

$$D := f[X_i] \rightarrow a_o \in \{\mathbb{T}, \mathbb{F}\} \quad \forall X_i$$

$$s^+[X_i] := P :$$
$$(P[X_i] \rightarrow y_o == a_o \quad \forall X_i) \;\; \cap \;\; (P[\hat{X}_i] \supseteq P[X_i] \quad \forall X_i, \hat{X}_i)$$

$$s^+ = \{s_1, s_2, ..., s_N | b_1, b_2, ..., b_M, y_o\} = \{s_1, s_2, ..., s_{O_T[n]}, b_1, b_2, ..., b_{O_S[n]}, y_o\}$$
$$= \{\mathcal{L}, \mathcal{M}, y_o\}$$

$$O_T[n] := |\mathcal{L}| = N$$

8.2 Space Complexity $O_S[n]$

Define Space Complexity $O_S[n]$ of solution s^+ to Decision Problem D as the total number of memory elements

$$X_i = \{x_1, ..., x_n, C\}; \;\; \hat{X}_i = \{x_1, ..., x_{n+1}, C\}$$

$$D := f[X_i] \rightarrow a_o \in \{\mathbb{T}, \mathbb{F}\} \quad \forall X_i$$

$$s^+[X_i] := P :$$
$$(P[X_i] \rightarrow y_o == a_o \quad \forall X_i) \;\; \cap \;\; (P[\hat{X}_i] \supseteq P[X_i] \quad \forall X_i, \hat{X}_i)$$

$$s^+ = \{s_1, s_2, ..., s_N | b_1, b_2, ..., b_M, y_o\} = \{s_1, s_2, ..., s_{O_T[n]}, b_1, b_2, ..., b_{O_S[n]}, y_o\}$$
$$= \{\mathcal{L}, \mathcal{M}, y_o\}$$

$$O_S[n] := |\mathcal{M}| + |y_o|^* = M + 1$$

*It is convention to reserve one memory element for output y_o.
Void programs do not require the y_o memory element for output

9 Definition of Complexity

Define Complexity $O[n]$ as a vector of dimension V

$$\mathbf{O}[n] := \;\; < O_T[n], O_S[n], O_3[n], O_4[n] ..., O_V[n] >$$

10 Simple Computational Complexity

The remainder of this document assumes simple computational complexity of dimension 2

10.1 Definition

Define simple computational complexity of dimension 2

$$\mathbf{O}[n] := \ < O_T[n], O_S[n] >$$

10.2 Total Complexity

Define Total Complexity of solution s^+

$$O[n] := |s^+[X_n]| = |\{\mathcal{L}, \mathcal{M}, y_o\}|$$

$$= |\mathcal{L}| + |\mathcal{M}| + |y_o| = N + M + 1$$

10.3 Time Complexity

Restate definition of Time Complexity $O_T[n]$ of solution s^+

$$s^+ = \{\mathcal{L}, \mathcal{M}, y_o\}$$

$$O_T[n] := |\mathcal{L}| = N$$

10.4 Space Complexity

Restate definition of Time Complexity $O_S[n]$ of solution s^+

$$s^+ = \{\mathcal{L}, \mathcal{M}, y_o\}$$

$$O_S[n] := |\mathcal{M}| + |y_o| = M + 1$$

10.5 Total Complexity as a Function of Time and Space Complexity

$$O[n] := |s^+[X_n]| = |\{\mathcal{L}, \mathcal{M}, y_o\}|$$

$$= |\mathcal{L}| + |\mathcal{M}| + |y_o|$$

$$= O_T[n] + O_S[n]$$

10.6 $O_S[n] > 0^*$

Assuming Program is not void

10.6.1 Proof

Assume $O_S[n] = 0$

$$O_S[n] := |\mathcal{M}| + |y_o|$$
$$O_S[n] = 0 \Rightarrow \mathcal{M} = y_o = \varnothing$$
$$y_o = \varnothing; \;\; y_o \in \{\mathbb{T}, \mathbb{F}\} \text{ by definition of } s^+$$

$\therefore O_S[n] = 0$ contradicts the definition of solution s^+ of a decision problem

$$O_S[n] \geqslant 0 \text{ by definition of magnitude}$$
$$\therefore O_S[n] > 0$$

10.7 $O_T[n] > 0^*$

Assuming Program is not void

10.7.1 Proof

Assume $O_T[n] = 0$

$$O_T[n] := |\mathcal{L}|$$
$$O_T[n] = 0 \Rightarrow y_o \notin \{\mathbb{T}, \mathbb{F}\}$$
$$y_o \notin \{\mathbb{T}, \mathbb{F}\}; \;\; y_o \in \{\mathbb{T}, \mathbb{F}\} \text{ by definition of } s^+$$

$\therefore O_T[n] = 0$ contradicts the definition of solution s^+ of a decision problem

$$O_T[n] \geqslant 0 \text{ by definition of magnitude}$$
$$\therefore O_T[n] > 0$$

10.8 $O[n] > 0^*$

Assuming Program is not void

10.8.1 Proof

$$O[n] := O_T[n] + O_S[n]$$

$$O_T[n] > 0; \;\; O_S[n] > 0$$

$$\therefore O[n] > 0$$

10.9 $O[n] > O_T[n]*$

Assuming Program is not void

10.9.1 Proof

$$O[n] := O_T[n] + O_S[n]$$

$$O_S[n] > 0$$

$$\therefore O[n] > O_T[n]$$

10.10 $O[n] > O_S[n]^*$

Assuming Program is not void

10.10.1 Proof

$$O[n] := O_T[n] + O_S[n]$$

$$O_T[n] > 0$$

$$\therefore O[n] > O_S[n]$$

10.11 $O[n+1] \geqslant O[n]$

10.11.1 Proof

$$X_i = \{x_1, ..., x_n, C\}; \;\; \hat{X}_i = \{x_1, ..., x_{n+1}, C\}$$

$$O[n] = \mid s^+[X_i] \mid$$

$$O[n+1] = \hat{O}[n] = \mid s^+[\hat{X}_i] \mid$$

For general solutions s^+

$$s^+[\hat{X}_i] \supseteq s^+[X_i]$$
$$\Rightarrow |s^+[\hat{X}_i]| \geqslant |s^+[X_i]|$$
$$\therefore \hat{O}[n] = O[n+1] \geqslant O[n]$$

11 Complexity of Canonical Instructions

$$c := a \leftarrow l[X_n]$$

12 Complexity of Computational Operations

12.1 $+$

Express the bounds of complexity for Computational Operation $+$

13 Inductive Functions

13.1 Inductive Function f_{n+1}

$$O[n] = O_T[n] + O_S[n]$$
$$O[n+1] = O_T[n+1] + O_S[n+1]$$

$$f_{n+1}[n] := O[n+1] - O[n]$$

13.2 Inductive Space and Time Formulas

$$f^T_{n+1}[n] := O_T[n+1] - O_T[n]$$
$$O_T[n+1] = O_T[n] + f^T_{n+1}[n]$$
$$f^S_{n+1}[n] := O_S[n+1] - O_S[n]$$
$$O_S[n+1] = O_S[n] + f^S_{n+1}[n]$$

13.3 Inductive Function Expressions

Relate $f_{n+1}[n]$ to equivalence functions

$$O[n] = O_T[n] + O_S[n]$$
$$O[n+1] = O_T[n+1] + O_S[n+1] = O[n] + f_{n+1}[n]$$
$$O_T[n] = O[n] - O_S[n]$$
$$O_S[n] = O[n] - O_T[n]$$

$$f_{n+1}[n] = O[n+1] - O[n]$$
$$f_{n+1}[n] = O_T[n+1] + O_S[n+1] - O[n]$$
$$f_{n+1}[n] = O_T[n+1] - O_T[n] + O_S[n+1] - O_S[n]$$
$$f_{n+1}[n] = O[n+1] - O_T[n] - O_S[n]$$
$$f_{n+1}[n] = f^T_{n+1}[n] + f^S_{n+1}[n]$$

13.4 Zero Order Space Inductive Function

$$Let\ \ O_S[n] \sim n^0$$
$$f_{n+1}[n] = O_T[n+1] - O_T[n] + O_S[n+1] - O_S[n] = O_T[n+1] - O_T[n]$$

14 Polynomial Complexity

14.1 Definition

Decision problem D with solution s^+ has polynomial total complexity $O[n]$ if

$$\exists K, C, \lambda_1 ... \lambda_K \quad :$$
$$O[n] = (\lambda_K n)^K + (\lambda_{K-1} n)^{K-1} ... + \lambda_1 n + C \quad \forall n$$

14.2 Polynomial Problems

Define $\mathbb{P}$, the set of Decision Problems that can be solved with Polynomial Complexity

$$\mathbb{P} := \{D_1, D_2, ...\} :$$

$$\exists K, C, \lambda_1 ... \lambda_K :$$
$$O[n] = (\lambda_K n)^K + (\lambda_{K-1} n)^{K-1} ... + \lambda_1 n + C \quad \forall n, D_i \in \mathbb{P}$$

14.3 Polynomial Order of Complexity

Solution s^+ with total complexity $O[n]$ is said to be of order n^K

$$O[n] \sim n^K$$
$$O[n] = (\lambda_K n)^K + (\lambda_{K-1} n)^{K-1} ... + \lambda_1 n + C \quad \forall n$$

14.4 Property of Polynomial Complexity 1

Solutions with polynomial complexity have convergent complexity

$$lim_{n \to \infty} \frac{O[n+1]}{O[n]} = 1$$

14.4.1 Proof

$$O[n] = (\lambda_K n)^K + (\lambda_{K-1} n)^{K-1} ... + \lambda_1 n + C$$
$$O[n+1] = (\lambda_K (n+1))^K + (\lambda_{K-1} (n+1))^{K-1} ... + \lambda_1 (n+1) + C$$
$$= (\lambda_K n)^K + (\tilde{\lambda}_{K-1} n)^{K-1} ... + \tilde{\lambda_1} n + \tilde{C}$$

$$lim_{n \to \infty} \frac{O[n+1]}{O[n]}$$

$$= lim_{n \to \infty} \frac{(\lambda_K n)^K + (\tilde{\lambda}_{K-1} n)^{K-1} ... + \tilde{\lambda_1} n + \tilde{C}}{(\lambda_K n)^K + (\lambda_{K-1} n)^{K-1} ... + \lambda_1 n + C}$$

$$= lim_{n\to\infty} \frac{(\lambda_K n)^K}{(\lambda_K n)^K + (\lambda_{K-1} n)^{K-1} ... + \lambda_1 n + C} + \frac{(\tilde{\lambda}_{K-1} n)^{K-1}}{(\lambda_K n)^K + (\lambda_{K-1} n)^{K-1} ... + \lambda_1 n + C} + ... + \frac{\tilde{\lambda}_1 n}{(\lambda_K n)^K + (\lambda_{K-1} n)^{K-1} ... + \lambda_1 n + C} + \frac{\tilde{C}}{(\lambda_K n)^K + (\lambda_{K-1} n)^{K-1} ... + \lambda_1 n + C}$$

$$= 1 = lim_{n\to\infty} \frac{O[n+1]}{O[n]}$$

14.5 Property of Polynomial Complexity 2

$$\exists K, \hat{C}, \hat{\lambda}_1, ..., \hat{\lambda}_{K-1} :$$
$$O[n+1] - O[n] = f_{n+1}[n] = (\hat{\lambda}_{K-1} n)^{K-1} ... + \hat{\lambda}_1 n + \hat{C} \quad \forall n$$

14.5.1 Proof

$$O[n] = (\lambda_K n)^K + (\lambda_{K-1} n)^{K-1} ... + \lambda_1 n + C$$
$$O[n+1] = (\lambda_K (n+1))^K + (\lambda_{K-1} (n+1))^{K-1} ... + \lambda_1 (n+1) + C$$
$$= (\lambda_K n)^K + (\tilde{\lambda}_{K-1} n)^{K-1} ... + \tilde{\lambda_1} n + \tilde{C}$$
$$O[n+1] - O[n] = ((\tilde{\lambda}_{K-1} - \lambda_{K-1}) n)^{K-1} ... + (\tilde{\lambda_1} - \lambda_1) n + (\tilde{C} - C)$$
$$O[n+1] - O[n] = (\hat{\lambda}_{K-1} n)^{K-1} ... + \hat{\lambda}_1 n + \hat{C}$$

14.6 Property of Polynomial Complexity 3

$$limit_{n\to\infty} \frac{f_{n+1}[n]}{O[n]} = 0$$

14.6.1 Proof

$$limit_{n\to\infty} \frac{O[n+1]}{O[n]} = 1$$
$$limit_{n\to\infty} \frac{O[n] + f_{n+1}[n]}{O[n]} = 1$$
$$limit_{n\to\infty} \frac{O[n]}{O[n]} + \frac{f_{n+1}[n]]}{O[n]} = 1$$
$$limit_{n\to\infty} 1 + \frac{f_{n+1}[n]]}{O[n]} = 1$$
$$limit_{n\to\infty} \frac{f_{n+1}[n]]}{O[n]} = 0$$

14.7 Total Polynomial Complexity Implies Time bounded Polynomial Complexity

$$D \in \mathbb{P} \Longrightarrow O_T[n] < (\lambda_K n)^K + (\lambda_{K-1} n)^{K-1} ... + \lambda_1 n + C$$

14.7.1 Proof

$$O[n] = (\lambda_K n)^K + (\lambda_{K-1} n)^{K-1} ... + \lambda_1 n + C \;\; \forall n$$

$$O[n] := O_T[n] + O_S[n]; \;\; O_S[n] > 0$$

$$\therefore O_T[n] < (\lambda_K n)^K + (\lambda_{K-1} n)^{K-1} ... + \lambda_1 n + C \;\; \forall n$$

14.8 Total Polynomial Complexity Implies Space bounded Polynomial Complexity

$$D \in \mathbb{P} \Longrightarrow O_S[n] < (\lambda_K n)^K + (\lambda_{K-1} n)^{K-1} ... + \lambda_1 n + C$$

14.8.1 Proof

$$O[n] = (\lambda_K n)^K + (\lambda_{K-1} n)^{K-1} ... + \lambda_1 n + C \;\; \forall n$$

$$O[n] := O_T[n] + O_S[n]; \;\; O_T[n] > 0$$

$$\therefore O_S[n] < (\lambda_K n)^K + (\lambda_{K-1} n)^{K-1} ... + \lambda_1 n + C \;\; \forall n$$

14.9 Polynomial Complexity in Space and Time Implies Polynomial Total Complexity

$$(O_S[n] == (\lambda_K n)^K + (\lambda_{K-1} n)^{K-1} ... + \lambda_1 n + \lambda_0)$$

$$\wedge$$

$$(O_T[n] == (\hat{\lambda}_M n)^M + (\hat{\lambda}_{M-1} n)^{M-1} ... + \hat{\lambda}_1 n + \hat{\lambda}_0)$$

$$\Rightarrow D \in \mathbb{P}$$

14.9.1 Proof

$$O_S[n] = \lambda_K n^K + \lambda_{K-1} n^{K-1} + ... + \lambda_1 n + \lambda_0$$

$$O_T[n] = \hat{\lambda}_M n^M + \hat{\lambda}_{M-1} n^{M-1} + ... + \hat{\lambda}_1 n + \hat{\lambda}_0$$

$$O[n] = O_S[n] + O_T[n]$$

$$^{*}O[n] = (\hat{\lambda}_0 + \lambda_0) + n(\lambda_1 + \hat{\lambda_1}) + ... + n^K(\lambda_K + \hat{\lambda_K}) + \hat{\lambda}_{K+1} n^{K+1} + ... + \hat{\lambda}_M n^M$$

$$\therefore O[n] \text{ has polynomial total complexity by definition}$$

* Assume K < M, similar proof for K=M, K>M

15 Non-Polynomial Complexity

15.1 Definition

Decision problem $\tilde{D}$ with solution s^+ has non-polynomial total complexity $O[n]$ if

$$\nexists K, \lambda_0, \lambda_1, ..., \lambda_K \ :$$
$$O[n] = (\lambda_K n)^K + (\lambda_{K-1} n)^{K-1} ... + \lambda_1 n + C \quad \forall n$$

15.2 Non-Polynomial Problems

Define $\mathcal{N}$, the set of Decision Problems that cannot be solved with Polynomial Complexity

$$\mathcal{N} := \{\tilde{D}_1, \tilde{D}_2, ...\} :$$
$$\nexists K, C, \lambda_1 ... \lambda_K \ :$$
$$O[n] = (\lambda_K n)^K + (\lambda_{K-1} n)^{K-1} ... + \lambda_1 n + C \quad \forall n, \ s^+ \in S_i^+, \ \tilde{D}_i \in \mathcal{N}$$

15.3 $\mathbb{P}$ and $\mathcal{N}$ are disjoint

$$\mathbb{P} \cap \mathcal{N} = \varnothing$$

15.3.1 Proof

Let D $\in \mathcal{N}$

$$\nexists K, C, \lambda_1 ... \lambda_K \ :$$
$$O[n] = (\lambda_K n)^K + (\lambda_{K-1} n)^{K-1} ... + \lambda_1 n + C \quad \forall n$$

Assume D $\in \mathbb{P}$

$$\exists K, C, \lambda_1 ... \lambda_K \ :$$
$$O[n] = (\lambda_K n)^K + (\lambda_{K-1} n)^{K-1} ... + \lambda_1 n + C \quad \forall n$$

Contradicts the definition of $\mathcal{N}$

$$\therefore D \in \mathcal{N} \Rightarrow D \notin \mathbb{P}$$

Let D $\in \mathbb{P}$

$$\exists K, C, \lambda_1 ... \lambda_K \ :$$
$$O[n] = (\lambda_K n)^K + (\lambda_{K-1} n)^{K-1} ... + \lambda_1 n + C \quad \forall n$$

Assume D $\in \mathcal{N}$

$$\nexists K, C, \lambda_1 ... \lambda_K \ :$$
$$O[n] = (\lambda_K n)^K + (\lambda_{K-1} n)^{K-1} ... + \lambda_1 n + C \quad \forall n$$

Contradicts the definition of $\mathbb{P}$

$$\therefore D \in \mathbb{P} \Rightarrow D \notin \mathcal{N}$$

$$D \in \mathcal{N} \Rightarrow D \notin \mathbb{P}; D \in \mathbb{P} \Rightarrow D \notin \mathcal{N}$$
$$\therefore \mathbb{P} \cap \mathcal{N} = \varnothing$$

16 Discrete Derivative; Z Transform

16.1 Discrete Derivative

Define derivative for discrete function f[n]

$$\Delta_n^1 f[n] := f[n+1] - f[n]$$

We will use the above definition for the remainder of this document

16.2 Zero Order Derivative

$$\Delta_n^0 f[n] = f[n]$$

16.3 Kth Discrete Derivative

Define the Kth derivative of discrete function $f[n]$

$$\Delta_n^K f[n] := \Delta_n^{K-1} f[n+1] - \Delta_n^{K-1} f[n]$$

16.4 Kth Discrete Derivative as an Alternating Sum

$$\Delta_n^K f[n] := \Delta_n^{K-1} f[n+1] - \Delta_n^{K-1} f[n]$$

$$= (\Delta_n^{K-2} f[n+2] - \Delta_n^{K-2} f[n+1]) - (\Delta_n^{K-2} f[n+1] - \Delta_n^{K-2} f[n])$$

$$= (\Delta_n^{K-2} f[n+2] - 2\Delta_n^{K-2} f[n+1] - \Delta_n^{K-2} f[n])$$

$$= \sum_{i=0}^{K} (-1)^j \ ({}_K C_j) \ \Delta_n^0 f[n+j]$$

$$= \sum_{i=0}^{K} (-1)^j \ ({}_K C_j) \ f[n+j]$$

16.5 Z Transform

Define the Z Transform for discrete function f[n]

$$\mathcal{Z}(f[n]) := \sum_{n=0}^{\infty} f[n] z^{-n}$$

16.6 Z Transform of 0 Order Derivative

$$\Delta_n^0 f[n] := f[n]$$

$$\mathcal{Z}(\Delta_n^0 f[n]) = \mathcal{Z}(f[n])$$

16.7 Z Transform of 1st Derivative

$$\Delta_n^1 f[n] := f[n+1] - f[n]$$

$$\mathcal{Z}(\Delta_n^1 f[n]) = \mathcal{Z}(f[n+1] - f[n])$$

$$= \sum_{n=0}^{\infty} (f[n+1] - f[n]) z^{-n}$$

$$= \sum_{n=0}^{\infty} (f[n+1] z^{-n} - f[n] z^{-n})$$

$$= \sum_{n=0}^{\infty} f[n+1] z^{-n} - \sum_{n=0}^{\infty} f[n] z^{-n}$$

$$= \sum_{m=0}^{\infty} f[m+1] z^{-m} - \sum_{n=0}^{\infty} f[n] z^{-n}$$

Let

$$\hat{m} = m + 1; \ \ m = \hat{m} - 1$$

$$= \sum_{m=0}^{\infty} f[\hat{m}] z^{-(\hat{m}-1)} - \mathcal{Z}(f[n])$$

$$= z^1 \sum_{\hat{m}=1}^{\infty} f[\hat{m}] z^{-\hat{m}} - \mathcal{Z}(f[n])$$

$$= z^1 \sum_{\hat{m}=1}^{\infty} f[\hat{m}] z^{-\hat{m}} + f[0] - f[0] - \mathcal{Z}(f[n])$$

$$= z^1 \sum_{\hat{m}=0}^{\infty} f[\hat{m}] z^{-\hat{m}} - f[0] - \mathcal{Z}(f[n])$$

$$= z^1 \mathcal{Z}(f[n]) - f[0] - \mathcal{Z}(f[n])$$

$$\mathcal{Z}(\Delta_n^1 f[n]) = \mathcal{Z}(f[n])(z^1 - 1) - f[0]$$

16.8 Z Transform of Kth Derivative

$$\mathcal{Z}(f[n]) := \sum_{n=0}^{\infty} f[n] z^{-n}$$

$$\mathcal{Z}(\Delta_n^K f[n]) = \sum_{n=0}^{\infty} \Delta_n^K f[n] z^{-n}$$

$$= \sum_{n=0}^{\infty} \sum_{i=0}^{K} (-1)^j \ ({}_K C_j) \ f[n+j] z^{-n}$$

$$= \sum_{n=0}^{\infty} (f[n+K] - ({}_K C_1) f[n+K-1] + ({}_K C_2) f[n+K-2] - ... \pm f[n]) z^{-n}$$

$$= \sum_{n=0}^{\infty} f[n+K] z^{-n} - ({}_K C_1) f[n+K-1] z^{-n} + ({}_K C_2) f[n+K-2] z^{-n} - \\ ... \pm f[n] z^{-n}$$

$$= z^K \mathcal{Z}(f[n]) + \sum_{i=0}^{K-1} f[i] - ({}_K C_1) z^{K-1} \mathcal{Z}(f[n]) - \sum_{j=0}^{K-2} f[j] + \\ ({}_K C_2) z^{K-2} \mathcal{Z}(f[n]) + \sum_{k=0}^{K-3} f[k] - ... \pm \mathcal{Z}(f[n])$$

When K is odd

$$\mathcal{Z}(\Delta_n^K f[n]) = (z-1)^K \mathcal{Z}(f[n]) + \sum_{i=0}^{\frac{n+1}{2}} f[2i] \quad K > 0$$

When K is even

$$\mathcal{Z}(\Delta_n^K f[n]) = (z-1)^K \mathcal{Z}(f[n]) + \sum_{j=0}^{\frac{n}{2}} f[2j+1] \quad K > 0$$

17 Divergent Complexity

17.1 Definition of Converges to

??Can f[n] = 0??

$$f[n] \ converges \ to \ C = convergent[f[n], C] = a_o; \ a_o \in \{\mathbb{T}, \mathbb{F}\} =$$

$$|C - f[n+1]| < |C - f[n]| \ \ \forall n$$

$$\wedge$$

$$\nexists K : |C - f[\hat{n}]| > K \quad \forall n; K > 0$$

17.1.1 Notation

C is commonly denoted by a limit

$$C = lim_{n \to \infty} f[n]$$

17.2 Definition of General Convergence

$$f[n] \text{ is } convergent = convergent[f[n]] = a_o;\ a_o \in \{\mathbb{T}, \mathbb{F}\} =$$

$$\exists C :$$

$$convergent[f[n], C] == \mathbb{T}$$

Alternatively

$$f[n] \text{ is } convergent = convergent[f[n]] = a_o \in \{\mathbb{T}, \mathbb{F}\} =$$

$$\exists C :$$

$$f[n]\ converges\ to\ C$$

17.3 Definition of Divergence

$$diverges[f[n]] = \neg converges[f[n]] = d_o;\ d_o \in \{\mathbb{T}, \mathbb{F}\}$$

$$:= \nexists C : convergent[f[n], C] == \mathbb{T}$$

17.4 Alternate Definition of Divergence

$$diverges[f[n]] = \neg converges[f[n]] = d_o;\ d_o \in \{\mathbb{T}, \mathbb{F}\}$$

$$= convergent[f[n], C] == \mathbb{F} \quad \forall C$$

17.4.1 Proof of Equivalence; Alternate Definition of Divergence

17.5 ?Necessary or Sufficient? Criteria 1 For Divergence

? The derivative as a function of K ? Function f[n] diverges if the Kth derivative of f[n] is strictly increasing

$$diverges[f[n]] := \nexists C : convergent[f[n], C] == \mathbb{T}$$

$$\Longleftrightarrow$$

$$\Delta_n^{K+1} f[n] > \Delta_n^K f[n] \quad \forall K$$

Alternatively

$$\Delta_n^{K+1} f[n] - \Delta_n^K f[n] > 0 \quad \forall K$$

$$\Delta_n^{K+2} > 0 \;\; \forall K$$

17.5.1 Criteria 1; Proof of Necessity and Sufficiency

$$diverges[f[n]] = d_o;\; d_o \in \{\mathbb{T}, \mathbb{F}\}$$

$$= \nexists C : convergent[f[n], C] == \mathbb{T}$$

Let

$$f[n] :$$
$$\Delta_n^{K+2} > 0 \quad \forall K$$

17.6 ?Necessary or Sufficient? Criteria 2 For Divergence

Function f[n] diverges if the Derivative as a function of K does not Converge

$$f[n] \; is \; Divergent = Divergent[f[n]] = a_o \in \{\mathbb{T}, \mathbb{F}\} :=$$

$$\nexists c : lim_{n \to \infty} \Delta_n^K f[n] = c$$

17.6.1 Criteria 2; Proof of Necessity and Sufficiency

$$diverges[f[n]] = d_o;\; d_o \in \{\mathbb{T}, \mathbb{F}\}$$

$$= \nexists C : convergent[f[n], C] == \mathbb{T}$$

17.7 Verbal Expressions

$$f[n] \; diverges = f[n] \; is \; divergent =$$
$$f[n] \; is \; not \; convergent = f[n] \; does \; not \; converge$$

17.8 Definition of Divergent Function

17.8.1 Definition 1

Define Divergent Function f[n] having strictly increasing K^{th} derivative

$$f[n]\ is\ Divergent = Divergent[f[n]] = a_o \in \{\mathbb{T}, \mathbb{F}\} :=$$

$$\Delta_n^{K+1} f[n] > \Delta_n^K f[n] \quad \forall K$$

Alternatively

$$\Delta_n^{K+1} f[n] - \Delta_n^K f[n] > 0 \quad \forall K$$

$$\Delta_n^{K+2} > 0 \ \ \forall K$$

17.8.2 Definition 2

$$f[n]\ is\ Divergent = Divergent[f[n]] = a_o \in \{\mathbb{T}, \mathbb{F}\} :=$$

$$\nexists c : lim_{n\to\infty} \Delta_n^K f[n] = c$$

17.8.3 Proof of Equivalence Definition 1 $\Leftrightarrow$ Definition 2

17.8.4 Sufficient Proof

$$f[n]\ is\ Divergent \Longleftrightarrow$$

$$\nexists c : lim_{n\to\infty} \Delta_n^K f[n] = c$$

17.8.5 Necessary Proof

$$f[n]\ is\ Divergent \Longleftrightarrow$$

$$\nexists c : lim_{n\to\infty} \Delta_n^K f[n] = c$$

Proof by contradiction of definition of limit Using the definition of increasing convergence for a discrete function*

17.9 Defintion

Decision problem $\hat{D}$ with solution s^+ has divergent total complexity $O[n]$ if

$$lim_{n\to\infty} \frac{O[n+1]}{O[n]} \ \ diverges$$

17.10 Divergent Problems

$$\hat{\mathcal{D}} := \{\hat{D}_1, \hat{D}_2, ...\} :$$
$$lim_{n\to\infty} \frac{O[n+1]}{O[n]} \ \ diverges \ \ \ \forall s^+ \in S_i^+, \ \hat{D}_i \in \hat{\mathcal{D}}$$

17.11 The Set of Polynomial Solutions and the Set of Divergent Solutions are disjoint

$$\mathbb{P} \cap \hat{D} = \varnothing$$

17.12 Proof

Let D $\in \hat{\mathcal{D}}$

$$lim_{n\to\infty} \frac{O[n+1]}{O[n]} \ \ diverges \text{ by definition}$$

Assume D $\in \mathbb{P}$

$$lim_{n\to\infty} \frac{O[n+1]}{O[n]} = 1$$

$$lim_{n\to\infty} \frac{O[n+1]}{O[n]} = 1 \text{ contradicts the definition of Divergent Problems}$$

$$\therefore D \in \hat{\mathcal{D}} \Rightarrow D \notin \mathbb{P}$$

Let D $\in \mathbb{P}$

$$lim_{n\to\infty} \frac{O[n+1]}{O[n]} = 1 \text{ by property of Polynomial complexity}$$

Assume D $\in \hat{D}$

$$lim_{n\to\infty} \frac{O[n+1]}{O[n]} \text{ diverges}$$

$$lim_{n\to\infty} \frac{O[n+1]}{O[n]} \text{ diverges contradicts a property of Polynomial complexity}$$

$$\therefore D \in \mathbb{P} \Rightarrow D \notin \hat{\mathcal{D}}$$

$$\therefore \mathbb{P} \cap \hat{\mathcal{D}} = \varnothing$$

18 Subprograms

18.1 Definition of Subprogram

Define a Subprogram of Program P; a subset of Program P

$$P := \{s_1, s_2, ..., s_N, b_1, b_2, ..., b_M, y_o\} = \{\mathcal{L}, \mathcal{M}, y_o\}$$
$$P_{sub} := \tilde{P} \mid$$
$$\tilde{P} \subseteq P$$

18.2 Identity Subprogram

18.2.1 Definition

18.2.2 Prove the Identity Subprogram is a Subprogram of P

18.3 Restate the subprogram condition of general solutions

Recall the definition of general solution s^+

$$X_i = \{x_1, ..., x_n, C\}; \;\; \hat{X}_i = \{x_1, ..., x_{n+1}, C\}$$
$$s^+ = s^+[X_i] := P :$$
$$(P[X_i] \rightarrow y_o == a_o \;\; \forall X_i) \;\; \cap \;\; (P[\hat{X}_i] \supseteq P[X_i] \;\; \forall X_i, \hat{X}_i)$$

The subprogram condition is one of two conditions for a general solution

$$P[\hat{X}_i] \supseteq P[X_i] \;\; \forall X_i, \hat{X}_i$$

The term subprogram is used interchangeably with the term subfunction

18.4 Prove O[n] is a non-decreasing function

Consider solution s^+ with complexity $O[n]$

$$X_i = \{x_1, ..., x_n, C\}; \;\; \hat{X}_i = \{x_1, ..., x_{n+1}, C\}$$
$$s^+ = s^+[X_i] := P :$$
$$(P[X_i] \rightarrow y_o == a_o \;\; \forall X_i) \;\; \cap \;\; (P[\hat{X}_i] \supseteq P[X_i] \;\; \forall X_i, \hat{X}_i)$$

$$s^+ = \{s_1, s_2, ..., s_N | b_1, b_2, ..., b_M, y_o\} = \{s_1, s_2, ..., s_{O_T[n]}, b_1, b_2, ..., b_{O_S[n]}, y_o\}$$
$$= \{\mathcal{L}, \mathcal{M}, y_o\}$$

$$O[n] := O_T[n] + O_S[n]$$
$$O_T[n] := |\mathcal{L}| = N$$
$$O_S[n] := |\mathcal{M}| + |y_o| = M + 1$$

O[n+1] denotes the total complexity for solution $s^+[\hat{X}_i]$

$$s^+[\hat{X}_i] = \hat{s}^+$$

Let

$$O[n+1] < O[n]$$
$$\Rightarrow \hat{N} + \hat{M} < N + M$$
$$\hat{s}^+ = \{s_1, s_2, ..., s_{\hat{N}} | b_1, b_2, ..., b_{\hat{M}}, y_o\}$$

$$\Rightarrow \hat{s}^+ \not\equiv s^+$$
$$P[\hat{X}_i] \not\equiv P[X_i] \quad \forall X_i, \hat{X}_i$$

$$\therefore O[n+1] < O[n] \text{ contradicts the definition of solution } s^+$$
$$O[n+1] \geqslant O[n]$$

18.5 Definition of Subfunction

$$X_i = \{x_1, ..., x_n, C\}; \;\; \hat{X}_i = \{x_1, ..., x_{n+1}, C\}$$
$$s^+ = s^+[X_i] := P :$$
$$(P[X_i] \rightarrow y_o == a_o \quad \forall X_i) \;\; \cap \;\; (P[\hat{X}_i] \supseteq P[X_i] \quad \forall X_i, \hat{X}_i)$$

$$s^+ = \{s_1, s_2, ..., s_N | b_1, b_2, ..., b_M, y_o\} = \{s_1, s_2, ..., s_{O_T[n]}, b_1, b_2, ..., b_{O_S[n]}, y_o\}$$
$$= \{\mathcal{L}, \mathcal{M}, y_o\}$$

$$Sub[X_i] := S = \{s_j, ... | b_k, ..., y_o\} :$$
$$s_j, b_k \in s^+ \quad \forall s_j, b_k \in S$$

18.5.1 $s^+[X_i]$ is a subfunction of $s^+[\hat{X}_i]$

$$s^+ = \{s_1, s_2, ..., s_N | b_1, b_2, ..., b_M, y_o\} = \{s_1, s_2, ..., s_{O_T[n]}, b_1, b_2, ..., b_{O_S[n]}, y_o\}$$
$$\hat{s}^+ = \{s_1, s_2, ..., s_N, ..., s_{\hat{N}} | b_1, b_2, ..., b_M, ..., b_{\hat{M}}, y_o\}; \;\; \hat{N} + \hat{M} \geqslant N + M$$

By definition of solution

$$\hat{s}^+ = P[\hat{X}_i] \supseteq P[X_i] = s^+ \quad \forall X_i, \hat{X}_i$$
$$\Rightarrow s_j, b_k \in \hat{s}^+ \;\; \forall s_j, b_k \in s^+$$

18.6 Subfunction Decomposition of Solutions

FIX Double check conditions!!! Solutions s^+ can be written as the union of subfunctions $Sub_k[X_i]$

$$X_i = \{x_1, ..., x_n, C\}; \;\; \hat{X}_i = \{x_1, ..., x_{n+1}, C\}$$
$$s^+ = s^+[X_i] := P :$$
$$(P[X_i] \rightarrow y_o == a_o \quad \forall X_i) \;\; \cap \;\; (P[\hat{X}_i] \supseteq P[X_i] \quad \forall X_i, \hat{X}_i)$$

$$s^+ = \{s_1, s_2, ..., s_N | b_1, b_2, ..., b_M, y_o\} = \{s_1, s_2, ..., s_{O_T[n]}, b_1, b_2, ..., b_{O_S[n]}, y_o\}$$
$$= \{\mathcal{L}, \mathcal{M}, y_o\}$$

$$s^+ = Sub_1[X_i] \cup Sub_2[X_i] \cup ... \cup Sub_z[X_i]$$
$$= \{\mathcal{L}_1 | \mathcal{M}_1, y_o\} \cup \{\mathcal{L}_2 | \mathcal{M}_2, y_o\} \cup ... \cup \{\mathcal{L}_z | \mathcal{M}_z, y_o\} :$$
$$\mathcal{L}_j \cap \mathcal{L}_k = \varnothing \;\; \forall j, k \neq j$$

$$s^+ = \{s_1^1, ..., s_{N_1}^1 | b_1^1, ..., y_o\} \cup \{s_1^2, ..., s_{N_2}^2 | b_1^2, ..., y_o\} \cup ... \cup \{s_1^z, ..., s_{N_z}^z | b_1^z, ..., y_o\} :$$
$$\textstyle\sum_{l=1}^{z} N_l = N = O_T[n]$$

19 Subfunction Complexity

19.1 Disjoint Subfunction Operations

$$\mathcal{L}_i \cap \mathcal{L}_j = \varnothing \;\; \forall i, j \neq i$$

19.2 Shared Subfunction Memory

$$|\mathcal{M}_i \cap \mathcal{M}_j| \geqslant 0 \;\; \forall i, j \neq i$$

19.2.1 Time Complexity of Subfunctions

Subfunction time complexity is additive

$$s^+ = \{\mathcal{L}, \mathcal{M}, y_o\}$$

$$Sub_i[X] := S_i = \{s_j, ...|b_k, ..., y_o\} :$$
$$s_j, b_k \in s^+ \quad \forall s_j, b_k \in S_i$$

$$s^+ = \{\mathcal{L}_1|\mathcal{M}_1, y_o\} \cup \{\mathcal{L}_2|\mathcal{M}_2, y_o\} \cup ... \cup \{\mathcal{L}_z|\mathcal{M}_z, y_o\} :$$
$$\mathcal{L}_i \cap \mathcal{L}_j = \varnothing \;\; \forall i, j \neq i$$

$$\mathcal{L} = \cup_{i=1}^{z} \mathcal{L}_i$$
$$\mathcal{L}_i \cap \mathcal{L}_j = \varnothing \;\; \forall i, j \neq i$$

$$O_T[n] = |\mathcal{L}| = N$$
$$O_T[n] = |\cup_{i=1}^{z} \mathcal{L}_i| = \sum_{i=1}^{z} |\mathcal{L}_i|^* = |\mathcal{L}_1| + |\mathcal{L}_2| + ... + |\mathcal{L}_z|$$
$$= O_{T_1}[n] + O_{T_2}[n] + ... + O_{T_z}[n] = N_1 + N_2 + ... + N_z$$

*Due to the disjoint condition of subfunction operations $\mathcal{L}_i \cap \mathcal{L}_j = \varnothing \;\; \forall i, j \neq i$

19.2.2 Space Complexity of Subfunctions

Subfunctions can access the full memory $\mathcal{M}$ with no added space complexity

$$s^+ = \{\mathcal{L}, \mathcal{M}, y_o\}$$

$$Sub_i[X] := S_i = \{s_j, ...|b_k, ..., y_o\} :$$
$$s_j, b_k \in s^+ \quad \forall s_j, b_k \in S_i$$

$$s^+ = \{\mathcal{L}_1|\mathcal{M}_1, y_o\} \cup \{\mathcal{L}_2|\mathcal{M}_2, y_o\} \cup ... \cup \{\mathcal{L}_z|\mathcal{M}_z, y_o\} :$$
$$\mathcal{L}_i \cap \mathcal{L}_j = \varnothing \;\; \forall i, j \neq i$$

$$s^+ = \{\mathcal{L}_1|\mathcal{M}, y_o\} \cup \{\mathcal{L}_2|\mathcal{M}, y_o\} \cup ... \cup \{\mathcal{L}_z|\mathcal{M}, y_o\} :$$
$$\mathcal{L}_i \cap \mathcal{L}_j = \varnothing \;\; \forall i, j \neq i$$

$$\mathcal{M} = \cup_{i=1}^{z} \mathcal{M}_i = \cup_{i=1}^{z} \mathcal{M}$$

$$O_S[n] = |\mathcal{M}| = M$$

$$O_S[n] = | \cup_{i=1}^{z} \mathcal{M}_i| = M$$

19.2.3 Shared State Notation

$$s^+ = \{\mathcal{L}, \mathcal{M}, y_o\}$$

$$Sub_i[X] := S_i = \{s_j, ...|b_k, ..., y_o\} :$$
$$s_j, b_k \in s^+ \quad \forall s_j, b_k \in S_i$$

$$s^+ = \{\mathcal{L}_1|\mathcal{M}, y_o\} \cup \{\mathcal{L}_2|\mathcal{M}, y_o\} \cup ... \cup \{\mathcal{L}_z|\mathcal{M}, y_o\} :$$
$$\mathcal{L}_i \cap \mathcal{L}_j = \varnothing \;\; \forall i, j \neq i$$

20 Polynomial Solution Subfunction Properties

20.1 Restate Definition of Subfunction

$$X_n = \{x_1, ..., x_n, C\}; \;\; \hat{X}_i = \{x_1, ..., x_{n+1}, C\}$$

$$s^+ = s^+[X_n] := P :$$
$$(P[X_n] \to y_o == a_o \quad \forall X_i) \quad \cap \quad (P[\hat{X}_n] \supseteq P[X_n] \quad \forall X_n, \hat{X}_n)$$

$$s^+ = \{s_1, s_2, ..., s_N | b_1, b_2, ..., b_M, y_o\} = \{s_1, s_2, ..., s_{O_T[n]}, b_1, b_2, ..., b_{O_S[n]}, y_o\}$$
$$= \{\mathcal{L}, \mathcal{M}, y_o\}$$

$$Sub[X_n] := S = \{s_j, ... | b_k, ..., y_o\} :$$
$$s_j, b_k \in s^+ \quad \forall s_j, b_k \in S$$

20.2 Property of Polynomial Solution Subfunctions

Let

$$D \in \mathbb{P}$$

$$X_n = \{x_1, ..., x_n, C\}; \;\; \hat{X}_n = \{x_1, ..., x_{n+1}, C\}$$

$$s^+ = s^+[X_n] := P :$$
$$(P[X_i] \to y_o == a_o \quad \forall X_n) \quad \cap \quad (P[\hat{X}_n] \supseteq P[X_n] \quad \forall X_n, \hat{X}_n)$$

$$\exists K, C, \lambda_1 ... \lambda_K \quad :$$
$$O[n] = (\lambda_K n)^K + (\lambda_{K-1} n)^{K-1} ... + \lambda_1 n + C \quad \forall n$$

$$s^+ = Sub_1[X_n] \cup Sub_2[X_n] \cup ... \cup Sub_z[X_n]$$

$$limit_{n \to \infty} \frac{O[n+1]}{O[n]} = 1$$
$$= limit_{n \to \infty} \frac{O_T^1[n+1] + O_T^2[n+1] + ... + O_T^z[n+1] + O_S[n+1]}{O_T^1[n] + O_T^2[n] + ... + O_T^z[n] + O_S[n]}$$

$$= limit_{n \to \infty} \frac{O_T^1[n] + O_T^2[n] + ... + O_T^z[n] + O_S[n] + f_{T_{n+1}}^1[n] + f_{T_{n+1}}^2[n+1] + ... + f_{T_{n+1}}^z[n] + f_{S_{n+1}}[n]}{O_T^1[n] + O_T^2[n] + ... + O_T^z[n] + O_S[n]}$$

$$= limit_{n\to\infty} 1 + \frac{f^1_{T_{n+1}}[n] + f^2_{T_{n+1}}[n+1] + ... + f^z_{T_{n+1}}[n] + f_{S_{n+1}}[n]}{O^1_T[n] + O^2_T[n] + ... + O^z_T[n] + O_S[n]} = 1$$

$$\Rightarrow limit_{n\to\infty} \frac{f^1_{T_{n+1}}[n] + f^2_{T_{n+1}}[n+1] + ... + f^z_{T_{n+1}}[n] + f_{S_{n+1}}[n]}{O^1_T[n] + O^2_T[n] + ... + O^z_T[n] + O_S[n]} = 0^*$$

$$\Rightarrow limit_{n\to\infty} \frac{f^i_{T_{n+1}}[n] + f_{S_{n+1}}[n]}{O^1_T[n] + O^2_T[n] + ... + O^z_T[n] + O_S[n]} = 0 \ \ \forall i$$

$$limit_{n\to\infty} \frac{f^i_{n+1}[n]}{O[n]} = 0 \ \ \forall i$$

* O[n] is a positive, non-decreasing function

20.3 Theorem of Polynomial Subfunctions

The Theorem of Polynomial Subfunctions states a solution has polynomial complexity if and only if all of its subfunctions have polynomial complexity

$$|s^+[X_n]| = O[n] = (\lambda_K n)^K + (\lambda_{K-1} n)^{K-1} ... + \lambda_1 n + C \quad \forall n$$

$$s^+ = Sub_1[X_n] \cup Sub_2[X_n] \cup ... \cup Sub_z[X_n]$$

$$O[n] = (\lambda_K n)^K + (\lambda_{K-1} n)^{K-1} ... + \lambda_1 n + C \quad \forall n$$

$$\Longleftrightarrow$$

$$|Sub_i[X_n]| = O_i[n] = (\hat{\lambda}_M n)^M + (\hat{\lambda}_{M-1} n)^{M-1} + ... + \hat{\lambda}_1 n + C \quad \forall i, n$$

20.3.1 Sufficient Proof

Solution s^+ having polynomial complexity implies all of its subfunctions Sub_i have polynomial complexity

Let

$$|s^+[X_n]| = O[n] = (\lambda_K n)^K + (\lambda_{K-1} n)^{K-1} ... + \lambda_1 n + C \quad \forall n$$
$$O[n] = \sum_{i=1}^{z} O_i[n] = O_1[n] + O_2[n] + ... + O_z[n]$$

Since $O[n], O_i[n]$ is positive, non-decreasing

$$O_i[n] = (\hat{\lambda}_{M_i} n)^{M_i} + (\hat{\lambda}_{M_i - 1} n)^{M_i - 1} ... + \hat{\lambda}_1 n + C \quad M_i \leqslant K \ \ \forall i, n$$

$\Rightarrow Sub_i$ has polynomial complexity by definition of polynomial complexity

20.3.2 Necessary Proof

Every subfunction Sub_i having polynomial complexity implies solution s^+ has polynomial complexity

Let

$$O_i[n] = (\hat{\lambda}_{M_i} n)^{M_i} + (\hat{\lambda}_{M_i - 1} n)^{M_i - 1} + ... + \hat{\lambda}_{1_i} n + \hat{\lambda}_{0_i} \quad \forall i, n$$

$$O_{max}[n]^* := \tilde{O}[n] \in \{O_1[n], O_2[n], ...O_z[n]\} :$$

$$lim_{n \to \infty} \frac{\tilde{O}[n]}{\sum_{i=1}^{z} O_i[n]} = c \neq 0$$

$$O_{max}[n] = (\hat{\lambda}_{M_{max}} n)^{M_{max}} + (\hat{\lambda}_{M_{max} 1} n)^{M_{max} - 1} + ... + \hat{\lambda}_{1_{max}} n + \hat{\lambda}_{0_{max}} \quad \forall i, n$$

$$O[n] = \sum_{i=1}^{z} O_i[n] = O_1[n] + O_2[n] + ... + O_z[n]$$

$$= (\tilde{\lambda}_L n)^L + (\tilde{\lambda}_{L-1} n)^{L-1} + ... + \tilde{\lambda}_1 n + C \quad L = M_{max} \ \ \forall n$$

$\Rightarrow s^+$ has polynomial complexity by definition of polynomial complexity

* O_{max} is not necessarily unique, but necessarily exists. See appendix for proof

21 Divergent Solution Subfunction Properties

21.1 Restate Definition of Subfunction

$$X_i = \{x_1, ..., x_n, C\}; \;\; \hat{X}_i = \{x_1, ..., x_{n+1}, C\}$$
$$s^+ = s^+[X_i] := P :$$
$$(P[X_i] \rightarrow y_o == a_o \;\; \forall X_i) \;\; \cap \;\; (P[\hat{X}_i] \supseteq P[X_i] \;\; \forall X_i, \hat{X}_i)$$

$$s^+ = \{s_1, s_2, ..., s_N | b_1, b_2, ..., b_M, y_o\} = \{s_1, s_2, ..., s_{O_T[n]}, b_1, b_2, ..., b_{O_S[n]}, y_o\}$$
$$= \{\mathcal{L}, \mathcal{M}, y_o\}$$

$$Sub[X_i] := S = \{s_j, ... | b_k, ..., y_o\} :$$
$$s_j, b_k \in s^+ \;\; \forall s_j, b_k \in S$$

21.2 Property of Divergent Subfunctions

Let

$$D \in \hat{\mathcal{D}}$$

$$X_n = \{x_1, ..., x_n\}; \;\; \hat{X}_n = \{x_1, ..., x_{n+1}\}$$

$$s^+ = s^+[X_n] := P :$$
$$(P[X_i] \rightarrow y_o == a_o \;\; \forall X_n) \;\; \cap \;\; (P[\hat{X}_n] \supseteq P[X_n] \;\; \forall \hat{X}_n : \hat{X}_n \supset X_n)$$

By Definition of Divergent Problem

$$\nexists c : limit_{n \rightarrow \infty} \frac{O[n+1]}{O[n]} = c$$
$$= limit_{n \rightarrow \infty} \frac{O_T^1[n+1] + O_T^2[n+1] + ... + O_T^z[n+1] + O_S[n+1]}{O_T^1[n] + O_T^2[n] + ... + O_T^z[n] + O_S[n]}$$

$$= limit_{n \rightarrow \infty} \frac{O_T^1[n] + O_T^2[n] + ... + O_T^z[n] + O_S[n] + f_{T_{n+1}}^1[n] + f_{T_{n+1}}^2[n+1] + ... + f_{T_{n+1}}^z[n] + f_{S_{n+1}}[n]}{O_T^1[n] + O_T^2[n] + ... + O_T^z[n] + O_S[n]}$$

$$= limit_{n \rightarrow \infty} 1 + \frac{f_{T_{n+1}}^1[n] + f_{T_{n+1}}^2[n+1] + ... + f_{T_{n+1}}^z[n] + f_{S_{n+1}}[n]}{O_T^1[n] + O_T^2[n] + ... + O_T^z[n] + O_S[n]} \neq c$$

$$\Rightarrow limit_{n \rightarrow \infty} \frac{f_{T_{n+1}}^1[n] + f_{T_{n+1}}^2[n+1] + ... + f_{T_{n+1}}^z[n] + f_{S_{n+1}}[n]}{O_T^1[n] + O_T^2[n] + ... + O_T^z[n] + O_S[n]} \neq c^*$$

Prove

$$\exists i : limit_{n\to\infty} \frac{f^i_{T_{n+1}}[n] + f_{S_{n+1}}[n]}{O^1_T[n] + O^2_T[n] + ... + O^z_T[n] + O_S[n]} \;\; diverges$$

* O[n] is a positive, non-decreasing function

21.3 Theorem of Divergent Subfunctions

The Theorem of Divergent Subfunctions states a divergent subfunction implies divergent total complexity

$$lim_{n\to\infty} \frac{O[n+1]}{O[n]} \;\; diverges$$

$$\Longleftrightarrow$$

$$\exists i : lim_{n\to\infty} \frac{O_i[n+1]}{O[n]} \;\; diverges$$

21.3.1 Sufficient Direction

See 18.2

21.3.2 Necessary Direction

22 Computational Basis

22.1 Definition of a Computational Basis of Program P

Define a Computational Basis B of Program P

$$X_n = \{x_1, x_2, ..., x_n\}$$
$$P[X_n] \to Y_o := \{s_1, s_2, ..., s_N, b_1, b_2, ..., Y_o\} \to Y_o$$

$$B :=$$

For the remainder of this document, "computational basis" is denoted as "basis"

22.2 Definition of the Identity Basis of Program P

22.3 Prove the Identity Basis of Program P is a basis of Program P

22.4 Definition of Canonical Program

22.5 Definition of a Canonical Basis of Program P

22.6 Prove Canonical Basis $\mathbb{B}$ of Program P is a basis of Program P

22.7 Subprogram and Canonical Basis

Prove a subprogram is a canonical basis if and only if it's basis decomposition is the identity subprogram

22.8 Basis of Boolean Program P

23 Fundamental Theorem of Computation

The Fundamental Theorem of Computation states every program P has a canonical basis $\mathbb{B}$

23.1 Proof

24 Input Spaces

24.1 Definition of Input Space

Define the Input Space $\mathbb{I}$ of Program P

24.2 Define the Cardinality Function $C[n]$ of Input Space $\mathbb{I}$

24.3 Existence, Uniqueness, etc.

24.4 Worst Case

24.5 Prove $|\mathbb{B}| = C[n]$

25 Theorem of Solution Complexity

The Theorem of Solution Complexity relates the complexity of solution s^+ to a basis B of solution s^+

$$X_n = \{x_1, x_2, ..., x_n\}$$

$$\mathcal{Q} := f[X_n] \rightarrow A_o \subseteq \Omega \quad \forall X_n \in D_{\mathcal{Q}}$$

$$s^+ = s^+[X_n] := P[X_n] \rightarrow Y_o :$$

$$(Y_o = A_o \quad \forall X_n \in D_{\mathcal{Q}}) \quad \cap \quad (P[X_{n+1}] \supseteq P[X_n] \quad \forall X_n \in D_{\mathcal{Q}} \quad \forall X_{n+1} \in D_{\mathcal{Q}})$$

26 Theorem of Optimal Complexity

27 Theorem of Divergent Complexity

28 Sum to N Problem with 2 integers

28.1 State formal definition of Sum to N : $x_i + x_j == N$

$$X_n = \{x_1, ..., x_n\}$$

$$D := f[X_i, N] = a_o \in \{\mathbb{T}, \mathbb{F}\} \quad \forall X_i$$

$$s^+[X_n] = P[X_n] :$$
$$(P[X_i] = y_o == a_o \quad \forall X_i) \quad \cap \quad (P[X_{n+1}] \supseteq P[X_n] \quad \forall X_{n+1})$$

$$s^+ = \{s_1, s_2, ..., s_{O_T[n]}, b_1, b_2, ..., b_{O_S[n]}, y_o\} = \{\mathcal{L}, \mathcal{M}, y_o\}$$

$$D = f[X_i] = \quad \exists x_j, x_k \in X_n \quad j \neq k :$$
$$x_j + x_k == N$$

28.2 Express a formal solution : $O_S[n] \sim n^0$

$$s^+ = \{s_1, s_2, ..., s_{O_T[n]}, b_1, b_2, ..., b_{O_S[n]}, y_o\} = \{\mathcal{L}, \mathcal{M}, y_o\}$$
$$s_1 = y_o \leftarrow \mathbb{F};$$
$$\forall i < n \ , \ n \geqslant j > i$$
$$s_2, s_3, s_8, s_9, ..., s_{3ij-4}, s_{3ij-3}..., s_{3n(n-1)-4}, s_{3n(n-1)-3} = b_1 \leftarrow x_i + x_j$$
$$s_4, s_5, s_{10}, s_{11}, ..., s_{3ij-2}, s_{3ij-1}..., s_{3n(n-1)-2}, s_{3n(n-1)-1} = b_1 \leftarrow b_1 == N$$
$$s_6, s_7, s_{12}, s_{13}..., s_{3ij}, s_{3ij+1}..., s_{3n(n-1)}, s_{3n(n-1)+1} = y_o \leftarrow y_o \vee b_1$$
$$s^+ = \{y_o \leftarrow \mathbb{F}, y_o \leftarrow y_o \vee (x_i + x_j == N) \quad \forall i, j > i \mid b_1, y_o\}$$

28.3 Prove s^+ satisfies the subfunction condition of solutions: $P[X_{n+1}] \supseteq P[X_n] \quad \forall X_{n+1}$

$$X_n = \{x_1, x_2, ..., x_n\}; \quad X_{n+1} = \{x_1, x_2, ..., x_n, x_{n+1}\}$$
$$s^+ = \{s_1, s_2, ..., s_{O_T[n]}, b_1, b_2, ..., b_{O_S[n]}, y_o\} = \{\mathcal{L}, \mathcal{M}, y_o\}$$
$$s^+_{n+1} = s^+ \cup \hat{s}^+$$
$$s_1 = y_o \leftarrow \mathbb{F};$$
$$\forall i < n \ , \ n \geqslant j > i$$

$$s_2, s_3, s_8, s_9, ..., s_{3ij-4}, s_{3ij-3}..., s_{3n(n-1)-4}, s_{3n(n-1)-3} = b_1 \leftarrow x_i + x_j$$
$$s_4, s_5, s_{10}, s_{11}, ..., s_{3ij-2}, s_{3ij-1}..., s_{3n(n-1)-2}, s_{3n(n-1)-1} = b_1 \leftarrow b_1 == N$$
$$s_6, s_7, s_{12}, s_{13}..., s_{3ij}, s_{3ij+1}..., s_{3n(n-1)}, s_{3n(n-1)+1} = y_o \leftarrow y_o \vee b_1$$

$$\forall k < n+1$$
$$s... = b_1 \leftarrow x_k + x_{n+1}$$
$$s... = b_1 \leftarrow b_1 == N$$
$$s... = y_o \leftarrow y_o \vee b_1$$

$$s^+ = \{y_o \leftarrow \mathbb{F}, y_o \leftarrow y_o \vee (x_i + x_j == N) \quad \forall i, j > i \mid b_1, y_o\}$$
$$\hat{s}^+ = \{y_o \leftarrow y_o \vee (x_k + x_{n+1} == N) \ \ \forall k < n+1 | \, b_1, y_o\}$$
$$s^+_{n+1} = \{y_o \leftarrow \mathbb{F}, y_o \leftarrow y_o \vee (x_i + x_j == N) \quad \forall i, j > i \mid b_1, y_o\} \ \cup$$
$$\{y_o \leftarrow y_o \vee (x_k + x_{n+1} == N) \ \ \forall k < n+1 | \, b_1, y_o\}$$

$$s^+_{n+1} = s^+ \cup \hat{s}^+ = P[X_{n+1}] \supseteq P[X_n] = s^+$$

28.4 Determine $O[n], O_S[n], O_T[n], f_{n+1}[n], f^T_{n+1}[n], f^S_{n+1}[n]$ for the above solution

$$O_S[n] = |y_o| + |b_1| = 2$$
$$O_T[n] = 3n(n-1) + 1 = 3n(n-1) - 1 + O_S[n]$$
$$O[n] = 3n(n-1) + 3 = 3n^2 - 3n + 3$$

$$f^S_{n+1}[n] = 0$$
$$f^T_{n+1}[n] = 6n$$
$$f_{n+1}[n] = f^S_{n+1}[n] + f^T_{n+1}[n]$$

28.5 Verify $O[n+1] = O[n] + f_{n+1}[n]$

$$O[n+1] = O[n] + \hat{O}[n]$$
$$3(n+1)^2 - 3(n+1) + 3 = 3n^2 - 3n + 3 + 6n$$
$$3n^2 + 6n + 3 - 3n - 3 + 3 = 3n^2 + 3n + 3$$

$$3n^2 + 3n + 3 = 3n^2 + 3n + 3$$

28.6 Show s^+ has Polynomial Complexity by the definition of Total Polynomial Complexity

$$O[n] = 3n^2 - 3n + 3$$

28.7 Show the limit $_{n\to\infty}\frac{O[n+1]}{O[n]}$ does not Diverge

$$limit_{n\to\infty}\frac{O[n+1]}{O[n]} =$$

$$limit_{n\to\infty}\frac{3n^2+3n+3}{3n^2-3n+3} =$$

$$limit_{n\to\infty}(\frac{3n^2-3n+3}{3n^2-3n+3} + \frac{6n}{3n^2-3n+3}) =$$

$$limit_{n\to\infty}(1 + \frac{6n}{3n^2-3n+3}) = 1$$

29 The Knapsack Problem

29.1 The Knapsack Problem

The Knapsack Problem is a famous problem in computer science which asks if objects can be stored in a knapsack. Typically the problem is designed with two constraints, weight and value. Given objects x_i, each with a respective weight w_i and value v_i, does there exist a combination of objects lighter than input weight W and more valuable than input value V?

29.2 Formal Definition

$$X_n = \{x_1, x_2, ..., x_n\} = \{\{w_1, v_1\}, \{w_2, v_2\}, ..., \{w_n, v_n\}\}$$

$$I = \{i_1, i_2, ..., i_n\} : i_l \in \{0, 1\} \ \ \forall i_l \in I$$

$$D := f[X_n, W, V] = a_o \in \{\mathbb{T}, \mathbb{F}\} = \exists I :$$
$$(\textstyle\sum_{j=1}^{n} i_j w_j < W) \wedge (\sum_{j=1}^{n} i_j v_j \geqslant V)$$

29.3 Solve for C[n]

29.3.1 Expressing I as a binary number

$$I = \{i_1, i_2, ..., i_n\} : i_l \in \{0, 1\} \ \ \forall i_l \in I$$

Valid combinations of I

$$I_{valid} =$$
$$\{\{0, 0, 0, ..., 0, 0, 1\}, \{0, 0, 0, ..., 0, 1, 0\}, \{0, 0, 0, ..., 0, 1, 1\}, ..., \{1, 1, 1, ..., 1, 1, 1\}\}$$

$$C[n] = |I_{valid}[n]| = 2^n - 1$$

29.3.2 Using a sum of combinations of inputs x_i

$$X_n = \{x_1, x_2, ..., x_n\} = \{\{w_1, v_1\}, \{w_2, v_2\}, ..., \{w_n, v_n\}\}$$

Valid combinations of x_i

$$X_{valid}[n] =$$

$$\{x_1\}\cup\{x_2\}\cup...\cup\{x_n\}\cup\{x_1,x_2\}\cup\{x_1,x_3\}\cup...\cup\{x_{n-1},x_n\}\cup...\cup\{x_1,x_2,...,x_n\}$$

$$= \ {}_{X_n}C_1 \cup \ {}_{X_n}C_2 \cup ... \cup \ {}_{X_n}C_n$$

$$C[n] = |X_{valid}[n]| = \sum_{j=1}^{n} \ {}_nC_j$$

29.3.3 Verify consistency

$$C[n] = |X_{valid}[n]| = |I_{valid}[n]|$$

$$= 2^n - 1 = \sum_{j=1}^{n} \ {}_nC_j = {}_nC_1 + {}_nC_2 + ... + {}_nC_n$$

$$= 2^n - 1 = 2^n - 1$$

29.4 Express a solution s^+ of the Knapsack Problem

29.5 Prove s^+ satisfies the subfunction condition of solutions

29.6 Determine $O[n], O_T[n], O_S[n], f_{n+1}[n]$

29.7 Show $s^+ \notin \mathbb{P}$

29.8 Express the Solution Space $\mathbb{S}$ for The Knapsack Problem

29.9 Prove a lower bound for all solutions $s^+ \in S^+ := O_{lower}[n]$

29.10 Prove D has Divergent Complexity

Appendix

30 Criticism of Overloaded Equivalence

In computer science, it is convention to overload equivalence =

$$a_i = a_i \quad \forall a_i \in \Omega$$

Consider standard C++ syntax
int x = 3;
int y = 4;
int z = x + y;

Int x is not inherently equal to 3. Rather, we are creating an open space "x" for a value and setting the value to 3. Similarly, z is not inherently equal to the value of x + y. Rather, we are creating an open space "z" for a value and setting the value to the sum of x and y which have already been set.

$$x \leftarrow 3$$

$$y \leftarrow 4$$

$$z \leftarrow x + y$$

31 Existence of $O_{max}[n]$

31.1 Proof

31.1.1 Alternate Definition; Left Hand Derivative

Some sources define

$$\Delta_n^1 f[n] = f[n] - f[n-1]$$

Citations

[1] $https://kapeli.com/cheat_sheets/LaTeX_Math_Symbols.docset/Contents/Resources/Docum$
[2] $chat.openai.com$
[3] $google.com$
[4] $wikipedia.org$
[3] $wolframalpha.com$
[4] $youtube.com$
[5] $https://stackoverflow.com/questions/3518973/floating-point-exponentiation-without-power-function$
[6] $https://stackoverflow.com/questions/27086195/linear-index-upper-triangular-matrix$