

# YAKEEN NEET 2.0

**2026**

Electric Charges and Fields

**Physics**

**Assignment Solution 01**

**By- Manish Raj (MR Sir)**



## QUESTION



Match the facts given in Column I with the Column II:

**1** I-b, II-a, III-d, IV-c

**2** I-b, II-a, III-c, IV-d

**3** I-b, II-d, III-a, IV-c

**4** I-b, II-c, III-d, IV-a


	Column-I		Column-II
I.	Charge cannot exist	a.	Without charge
II.	Mass can exist (new)	b.	Without mass
III.	Charge is	c.	Not conserved
IV.	Mass is	d.	conserved

## QUESTION



If a body is charged by rubbing it with another body then weight of the system:

- (A) may increase slightly
- (B) may decrease slightly
- (C) must increase slightly
- ☒ (D) remains precisely constant



1 Only A

2 AB


3 D ✓

4 C



## QUESTION

$$\text{Ring} \rightarrow E = \frac{kQx}{(R^2 + x^2)^{3/2}}$$




A uniformly charged ring is placed in  $x$ - $z$  plane as shown in figure. A point charge  $(+q)$  is placed on  $y$ -axis and it is in equilibrium. Choose the correct statement.

1 If  $y = \frac{R}{\sqrt{2}}$ , nature of equilibrium is stable

2 If  $y > \frac{R}{\sqrt{2}}$ , nature of equilibrium is unstable





3 If  $y < \frac{R}{\sqrt{2}}$ , nature of equilibrium is unstable


4 If  $y = \frac{R}{\sqrt{2}}$  is the position where electric force experienced by  $+q$  is minimum





# Ring symmetry






$$E = \frac{K 4Qx}{(R^2 + x^2)^{3/2}}$$

$$= \frac{K 4Qx}{\left(\left(\frac{l}{\sqrt{2}}\right)^2 + x^2\right)^{3/2}}$$

E will be max of  $x = \frac{RadR}{\sqrt{2}} = \frac{l}{\sqrt{2} \times \sqrt{2}}$

$x = \frac{l}{2}$



$$2R = \sqrt{2}l$$

$R = \frac{\sqrt{2}l}{2\sqrt{2}} = \frac{l}{\sqrt{2}}$

## QUESTION

Two equal point charges A and B are R distance apart. A third point charge placed on the perpendicular bisector at a distance 'x' from the centre the experience maximum electrostatic force when: ✓

1  $x = \frac{R}{2\sqrt{2}}$  ✓


2  $x = \frac{R}{\sqrt{2}}$

3  $x = R\sqrt{2}$

4  $x = 2\sqrt{2}R$

$$= \frac{\text{Radius}}{\sqrt{2}}$$

$$= \frac{R}{2\sqrt{2}}$$





## QUESTION




Find magnitude of net electric force on charge  $Q$  placed on X-axis:

1  $\frac{\sqrt{2}Q^2}{8\pi \epsilon_0 R^2}$

2  $\frac{\sqrt{2}Q^2}{4\pi \epsilon_0 R^2}$

3  $\frac{\sqrt{2}Q^2}{8\pi \epsilon_0 R}$

4  $\frac{2\sqrt{2}Q^2}{8\pi \epsilon_0 R^2}$



$$f = QE$$
$$= Q \left( \frac{kQR}{(R^2 + R^2)^{3/2}} \right)$$

✓





## QUESTION




**Assertion (A):** If a spherical cavity is made inside a uniformly charged sphere, then electric field inside the cavity is uniform.

**Reason (R):** The electric field inside the cavity is independent of radius of cavity, but depends on radius of sphere.

- 1 Both (A) and (R) are true and (R) is the correct explanation of (A).
- 2 Both (A) and (R) are true and (R) is NOT the correct explanation of (A).
- 3 (A) is true but (R) is false.
- 4 (A) is false but (R) is true.



$$\begin{aligned}
 (E_p)_{net} &= E_{comp} - E_{cavity} \\
 &= \frac{\rho \vec{r}}{3\epsilon_0} - \frac{\rho \vec{r}_1}{3\epsilon_0} \\
 &= \frac{\rho}{3\epsilon_0} (\vec{r} - \vec{r}_1) = \frac{\rho \vec{r}_0}{3\epsilon_0}
 \end{aligned}$$



Uniform

$$\begin{aligned}
 \vec{r}_0 + \vec{r}_1 &= \vec{r} \\
 \vec{r}_0 &= \vec{r} - \vec{r}_1
 \end{aligned}$$



# QUESTION

100%

$$\begin{aligned} 2\pi r &\rightarrow \text{circumference} \\ 10^r &\rightarrow \frac{2}{2\pi r} \\ 2\theta &\rightarrow \frac{2}{2\pi r} \end{aligned}$$

Force B/w Ring &  $Q$   
charge at centre = zero




A thin ring of radius ' $r$ ' made up of a conducting material having charge ' $q$ '. Find the increase in the tension the ring if a point charge  $Q$  is placed at centre of ring:

1  $\frac{Qq}{8\pi^2\epsilon_0 r^2}$  ✓ As


2  $\frac{Qq}{4\pi r^2 \epsilon}$

3  $\frac{Qq}{8\pi r^2 \epsilon_0}$

4  $\frac{Qq}{16\pi^2 r^2 \epsilon_0}$



$$dq = \frac{q\theta}{\pi}$$



$$F_c = \frac{kQq\theta}{\pi r^2} \quad (1)$$


$$\begin{aligned} T &= \frac{kQq}{2\pi r^2} \\ T &= \frac{Qq}{4\pi\epsilon_0 \times 2\pi r^2} \end{aligned}$$




## QUESTION



Consider the charge configuration shown in the figure. Variation of electric field along x-axis is




1




2




3





4





None of these









## QUESTION

JEE Adv - 2021



Three large non-conducting plane sheets of charge with charge densities  $\sigma$ ,  $2\sigma$  and  $3\sigma$  are arranged as shown. The net electric field in region III will be


1  $\frac{3\sigma}{\epsilon_0}$  (towards left)

2  $\frac{2\sigma}{\epsilon_0}$  (towards left)

3 0 ✓

4  $\frac{3\sigma}{\epsilon_0}$  (towards right)

$\vec{E}_I \Rightarrow$



$$\begin{aligned} E_{\text{net}} &= E_2 + E_3 - E_1 \\ &= \frac{2\sigma}{2\epsilon_0} + \frac{\sigma}{2\epsilon_0} - \frac{3\sigma}{2\epsilon_0} \\ &= \frac{3\sigma - 3\sigma}{2\epsilon_0} = 0 \end{aligned}$$



## QUESTION



Electric field of a region varies as  $\vec{E} = bx \hat{i} \text{ NC}^{-1}$ . Find the net charge enclosed by the cube as shown in figure.

non uniform

1  $\frac{bl^3}{2\epsilon_0}$

2  $\frac{bl^3}{3\epsilon_0}$


3  $bl^3\epsilon_0$

4  $\frac{bl^3\epsilon_0}{2}$

$$\oint_{\text{enc sur}} = \frac{q_{\text{in}}}{\epsilon_0}$$

$$bq^3 = \frac{q_{\text{in}}}{\epsilon_0}$$

$$q_{\text{in}} = \epsilon_0 b l^3$$



$E = 0$   
(0, y, z)  
 $\phi_A = EA = 0$

$E = bl\hat{i}$   
 $\phi_B = EA = bl \times l^2 = bl^3$

## QUESTION



A short electric dipole is placed along a radial line of a uniformly charged sphere as shown. If the force experienced by dipole at  $r = 2R$  and  $r = 4R$  and  $F_1$  and  $F_2$  respectively, then  $F_2/F_1$  is


1 ✓  $1/8$

2  $1/4$

3  $1/16$


4  $1/32$

$$F \propto \frac{1}{r^2} \times$$



$$\vec{F} = qE = Q \frac{2kP}{r^3}$$

$$\boxed{F \propto \frac{1}{r^3}}$$





## QUESTION

H/W



An electron moving with the speed  $5 \times 10^6$  m/s is projected parallel to the electric field of intensity  $1 \times 10^3$  N/C. Field is responsible for the retardation of motion of electric. Find the distance travelled by the electron before coming to rest for an instant ( $m_e = 9 \times 10^{-31}$  kg,  $e = 1.6 \times 10^{-19}$  C)

- 1 7 m
- 2 0.7 mm
- 3 7 cm
- 4 0.7 cm

## QUESTION

જાન નિશ્ચય



A conducting charged spherical shell has a tiny hole cut into its surface. If surface charge density near the hole is  $\sigma$  then electric field in the hole is:-

- 1  $\frac{\sigma}{\epsilon_0}$ , directed inwards
- 2  $\frac{\sigma}{\epsilon_0}$ , directed outwards
- 3  $\frac{\sigma}{2\epsilon_0}$ , directed inwards
- 4  $\frac{\sigma}{2\epsilon_0}$ , directed outwards ✓



## QUESTION

If linear density of a wire of length  $L$  depends on distance  $x$  from one end as  $\lambda = \frac{\lambda_0 x}{L}$ .  
Find total charge of wire.

**1**  $\lambda_0 L$

**2**  $\frac{\lambda_0 L}{2}$

**3**  $\frac{\lambda_0 L}{4}$

**4**  $2\lambda_0 L$






## QUESTION

A long charged wire is kept on  $+y$ -axis, with one end at origin. Its linear charge density is given as  $\lambda = \lambda_0 y$ . Calculate electric flux passing through the cylinder.

- 1  $\frac{\lambda_0 L^2}{\epsilon_0}$
- 2  $\frac{\lambda_0 L^2}{2 \epsilon_0}$
- 3  $\frac{2\lambda_0 L^2}{\epsilon_0}$
- 4 None



## QUESTION

Match the column:

**Code:**









**P      Q      R      S**

**1**    2    1    4    3

**2**    3    1    2    4

**3**    3    2    1    4

**4**    1    4    3    2


	List-I		List-II
(P)		(1)	
(Q)		(2)	
(R)		(3)	
(S)		(4)	




# QUESTION

Two metal spheres of same mass are suspended from a common point by a light insulating string. The length of each string is same. The spheres are given electric charges  $+q$  on one and  $+2q$  on the other. Which of the following diagram best shows the resulting positions of spheres?


1




2



3



4



## QUESTION

don't do it  
Proton



Two identical charged spheres suspended from a common point by two massless strings of length  $l$  are initially at a distance  $d$  ( $d \ll l$ ) apart because of their mutual repulsion. The charge begins to leak from both the spheres at a constant rate. As a result the charge approach each other with a velocity  $v$ . Then as a function of distance  $x$  between them.

**[NEET-2016]**

**1**  $v \propto x^{-1}$

**2**  $v \propto x^{\frac{1}{2}}$

**3**  $v \propto x$

**4**  $v \propto x^{-\frac{1}{2}}$



## QUESTION



A charged particle  $q$  of mass  $m$  is in equilibrium at a height  $h$  from a horizontal infinite line charge with uniform linear charge density  $\lambda$ . The charge lies in the vertical plane containing the line charge. If the particle is displaced slightly (vertically) prove that the motion of the charged particle will be simple harmonic. Also find its time period.



## QUESTION

Two point charges  $2\ \mu\text{C}$  and  $8\ \mu\text{C}$  are placed at position  $x = 0$  and  $x = 4\ \text{m}$  respectively. The distance of the neutral point from the smaller charge is found to be at  $x = \alpha/\beta$ . Find the value of  $(\alpha + \beta)$ .

1 7

2 5

3 4

4 3



## QUESTION

**Statement-I:** In a given situation of arrangement of charges, an extra charge is placed outside the Gaussian surface. In the Gauss Theorem

$$\oint \vec{E} \cdot d\vec{s} = \frac{Q_{\text{in}}}{\epsilon_0}$$

$Q_{\text{in}}$  remains unchanged where  $\vec{E}$  as electric field at the site of the element is changed.

**Statement-II:** Electric field  $E$  at any point on the Gaussian surface is due to inside charge only.

- 1 Statement-I is true, statement-II is true and statement-II is correct explanation for statement-I.
- 2 Statement-I is true, statement-II is true and statement-II is NOT the correct explanation for statement-I.
- 3 Statement-I is true, statement-II is false.
- 4 Statement-I is false, statement-II is true.

**Statement-I:** Coulombic force follows the principle of superposition.  
**and**

**Statement-II:** Coulombic force is directly proportional to the charges on both particle.

- 1** Statement-I is true, statement-II is true and statement-II is correct explanation for statement-I.
- 2** Statement-I is true, statement-II is true and statement-II is NOT the correct explanation for statement-I.
- 3** Statement-I is true, statement-II is false.
- 4** Statement-I is false, statement-II is true.



## QUESTION



**Assertion (A):** If a positive point charge is brought near an isolated neutral metal cube, the cube becomes negatively charged.

**Reason (R):** If a positive point charge is brought near an isolated neutral metal cube, the cube remains neutral.

- 1 Both (A) and (R) are true and (R) is the correct explanation of (A).
- 2 Both (A) and (R) are true and (R) is NOT the correct explanation of (A).
- 3 (A) is true but (R) is false.
- 4 (A) is false but (R) is true.

## QUESTION



**Assertion (A):** If a charge is released from rest in an electric field, it will always start to move along an electric field line.

**Reason (R):** Force on a charged particle is always in the direction of electric field.

- 1 Both (A) and (R) are true and (R) is the correct explanation of (A).
- 2 Both (A) and (R) are true and (R) is NOT the correct explanation of (A).
- 3 (A) is true but (R) is false.
- 4 (A) is false but (R) is true.




## QUESTION



A few electric field lines for a system of two charges  $Q_1$  and  $Q_2$  fixed at two different points on their axis are shown in the figure. With the figure what information about charges and electric field is incorrect.

- 1  $|Q_1| > |Q_2|$
- 2  $Q_1 \rightarrow +ve, Q_2 \rightarrow -ve$
- 3 At a finite distance to the left of  $Q_1$  the electric field is zero
- 4 At a finite distance to the right of  $Q_2$  the electric field is zero




## QUESTION



For the arrangement shown in the Fig., the two positive charges,  $+Q$  each, are fixed. Mark out the correct statement regarding a third charged particle  $-q$  placed at mid point  $P$  that can be displaced along or perpendicular to the line connecting the charges,

- 1 The particle will perform *SHM* for  $x \ll a$ .
- 2 The particle will undergo oscillatory motion but not *SHM* about  $P$  for  $x$  comparable to  $a$ .
- 3 The particle will perform *SHM* for  $y \ll a$ .
- 4 The particle will undergo oscillatory motion but not *SHM* about  $P$  for  $y \ll a$ .





## QUESTION



Match Column-I with Column-II

**1** I-c, II-d, III-a, IV-b

**2** I-b, II-d, III-c, IV-a

**3** I-b, II-d, III-a, IV-c

**4** I-b, II-a, III-d, IV-c

	Column-I		Column-II
I.	Electric field due to infinite plane sheet of charge	a.	0
II.	Electric field due to infinite conducting plane sheet of uniform thickness	b.	$\frac{\sigma}{2\epsilon_0}$
III.	Electric field due to Non-conducting charged solid sphere at its surface	c.	$\frac{R\rho}{3\epsilon_0}$
IV.	Electric field due to Non-conducting charged solid sphere at its centre	d.	$\frac{\rho}{\epsilon_0}$

## QUESTION



Two charges  $(-q, m)$  and  $(+q, m)$  are connected by a massless rod of length  $l$  and released in a region of uniform electric field as shown. Find the maximum speed of the charges:

**1**  $\sqrt{\frac{qE\ell}{2m}}$

**2**  $\sqrt{\frac{3qE\ell}{2m}}$


**3**  $\sqrt{\frac{3qE\ell}{m}}$

**4**  $\sqrt{\frac{qE\ell}{3m}}$

C.O.M.E ✓

find  $\omega$

$V = r\omega$



**THANK**  
**YOU**