

Bayesian Network Independencies

测验, 5 个问题

✓ 恭喜！您通过了！

下一项




1 / 1
分数

1.

Independencies in a graph.

Which pairs of variables are independent in the graphical model below, given that none of them have been observed? You may select 1 or more options.



A, B



正确

There are no active trails between A and B, so they are independent.



A, E



未选择的是正确的



D, E



未选择的是正确的



A, C



Bayesian Network Independencies

测验, 5 个问题



None - there are no pairs of independent variables.



未选择的是正确的




1 / 1
分数

2.

***Independencies in a graph.** (An asterisk marks a question that is more challenging. Congratulations if you get it right!)

Now assume that the value of E is known. (E is observed. A, B, C, and D are not observed.) Which pairs of variables (not including E) are independent in the same graphical model, given E? You may select 1 or more options.



None - given E, there are no pairs of variables that are independent.



正确

Observing E activates the V-structures around C and E, giving rise to active trails between every pair of variables in the network.



A, B



未选择的是正确的



A, C



未选择的是正确的



A, D



Bayesian Network Independencies

未选择的是正确的
测验, 5 个问题



B, D



未选择的是正确的



D, C



未选择的是正确的



B, C



未选择的是正确的



1 / 1
分数

3.

I-maps. I-maps can also be defined directly on graphs as follows. Let $I(G)$ be the set of independencies encoded by a graph G . Then G_1 is an I-map for G_2 if $I(G_1) \subseteq I(G_2)$.

Which of the following statements about I-maps are true? You may select 1 or more options.



A graph K is an I-map for a graph G if and only if all of the independencies encoded by K are also encoded by G .



正确

K is an I-map for G if K does not make independence assumptions that are not true in G . An easy way to remember this is that the complete graph, which has no independencies, is an I-map of all distributions.



A graph K is an I-map for a graph G if and only if K encodes all of the independencies that G has and more.



未选择的是正确的



An I-map is a function f that maps a graph G to itself, i.e., $f(G) = G$.



未选择的是正确的



The graph K that is the same as the graph G , except that all of the edges are oriented in the opposite direction as the corresponding edges in G , is always an I-map for G , regardless of the structure of G .



未选择的是正确的

Bayesian Network Independencies

测验, 5个问题

未选择的是正确的




1 / 1
分数

4.

*Naive Bayes.

Consider the following Naive Bayes model for flu diagnosis:



Assume a population size of 10,000. Which of the following statements are true in this model? You may select 1 or more options.



Say we observe that 1000 people have the flu, out of which 500 people have a headache (and possibly other symptoms) and 500 have a fever (and possibly other symptoms).

We would expect that approximately 250 people with the flu also have both a headache and fever.

正确

Given that someone has the flu, whether he has a headache is independent of whether he has a fever. We can thus calculate:

$$\begin{aligned}
 P(\text{Headache} = 1, \text{Fever} = 1 | \text{Flu} = 1) &= P(\text{Headache} = 1 | \text{Flu} = 1) \times P(\text{Fever} = 1 | \text{Flu} = 1) \\
 &\approx 0.5 * 0.5 \\
 &= 0.25.
 \end{aligned}$$

Since 1000 people have the flu, we can estimate that 250 of these people will have both a headache and fever.

Note that this is only an estimate: we can assert with high confidence that

$P(\text{Headache} = 1, \text{Fever} = 1 | \text{Flu} = 1)$ is near to 0.25, but in general it will not be exactly 0.25.

Moreover, even if it is exactly 0.25, the number of people with the flu, a headache and a fever need not

Bayesian Network Independencies

be exactly 250 all the time. Think of this as analogous to flipping a fair coin: even though the probability of getting a head is exactly 0.5, in any given sequence of coin flips we need not see exactly half of the coin turning up heads.



Say we observe that 1000 people have the flu, out of which 500 people have a headache (and possibly other symptoms) and 500 people have a fever (and possibly other symptoms).

Without more information, we cannot estimate how many people with the flu also have both a headache and fever.



未选择的是正确的



Say we observe that 1000 people have a headache (and possibly other symptoms), out of which 500 people have the flu (and possibly other symptoms), and 500 people have a fever (and possibly other symptoms).

Without more information, we cannot estimate how many people with a headache also have both the flu and a fever.



正确

Even after observing the Headache variable, there is still an active trail from Flu to Fever. Thus, the probability of someone with a headache also having a flu is dependent on the probability of his having a fever as well. For example, if someone has a flu, he could be more likely to have a fever, irrespective of whether he has a headache or not.

We therefore cannot estimate $P(Flu = 1, Fever = 1 | Headache = 1)$ from the conditional marginal probabilities $P(Flu = 1 | Headache = 1)$ and $P(Fever = 1 | Headache = 1)$.



Say we observe that 500 people have a headache (and possibly other symptoms) and 500 people have a fever (and possibly other symptoms).

Without more information, we cannot estimate how many people have both a headache and fever.



正确

Without having observed the Flu variable, there is an active trail from Headache and Fever. Thus, the probability of someone having a headache (without observing flu status) is not independent of the probability of the same person having a fever. For example, if someone has a headache, he might be more likely to have the flu, which would correspondingly increase the probability that he has a fever as well.

We therefore cannot estimate $P(Headache = 1, Fever = 1)$ from the marginal probabilities $P(Headache = 1)$ and $P(Fever = 1)$.



1 / 1
分数

5.

I-maps.

Bayesian Network Independencies

测验 5 个问题
Suppose $(A \perp B) \in \mathcal{I}(P)$, and G is an I-map of P , where G is a Bayesian network and P is a probability distribution. Is it necessarily true that $(A \perp B) \in \mathcal{I}(G)$?

☐ Yes

☒ No

正确

Since G is an I-map of P , all independencies in G are also in P . However, this doesn't mean that all independencies in P are also in G . An easy way to remember this is that the complete graph, which has no independencies, is an I-map of all distributions.

