

Q-10.13.3.10

Yash Patil - EE22BTECH11058

Question: All the jacks, queens and kings are removed from a deck of 52 playing cards. The remaining cards are well shuffled and then one card is drawn at random. Giving ace a value 1 similar value for other cards, find the probability that the card has a value

- 1) 7
- 2) greater than 7
- 3) less than 7

Solution: Number of cards left after removing all jacks, queens and kings(=N)

$$= 52 - 4 \times 3 \quad (1)$$

$$= 40 \quad (2)$$

Parameter	Value	Description
X	1-10	Represents the value of the card picked

Finding pmf:

$$p_X(k) = \Pr(X = k) \quad (3)$$

$$= \frac{4 \times 1}{40} \quad (4)$$

$$= \frac{1}{10} \quad (5)$$

$$(6)$$

CDF for the following pmf is:

$$F_X(k) = \sum_{k=k_0}^{k=k_1} p_X(k) \quad (7)$$

$$= (k_1 - k_0 + 1) \times \frac{1}{10} \quad (8)$$

- 1) Probability that card has value equal to 7:

$$= F_X(k); k = 7 \quad (9)$$

$$= 1 \times \frac{1}{10} = \frac{1}{10} \quad (10)$$

- 2) Probability that card has value greater than 7

$$= F_X(k); 8 \leq k \leq 10 \quad (11)$$

$$= 3 \times \frac{1}{10} = \frac{3}{10} \quad (12)$$

- 3) Probability that card has value less than 7

$$= F_X(k); 1 \leq k \leq 6 \quad (13)$$

$$= 6 \times \frac{1}{10} = \frac{6}{10} \quad (14)$$