

Lecture 10: Single-Cell RNA Sequencing

BIOINF3005/7160: Transcriptomics Applications

Dr Stephen Pederson

Bioinformatics Hub,
The University of Adelaide

May 25th, 2020

Background

scRNA Protocols

Cell Isolation

Sequencing Protocols

Data Analysis

QC

Quantification

Normalisation

Clustering

DE Analysis

Trajectory Analysis

Spatial Transcriptomics

Background

Introduction

- scRNA-Seq is the 'latest and greatest' transcriptomic technique
- Previously all our analysis involved multiple cells per sample
- All were combined during tissue extraction, library preparation etc.
- Most experiments have **highly** heterogeneous cell populations, e.g.

Introduction

- scRNA-Seq is the ‘latest and greatest’ transcriptomic technique
- Previously all our analysis involved multiple cells per sample
- All were combined during tissue extraction, library preparation etc.
- Most experiments have **highly** heterogeneous cell populations, e.g.
 - Different regions of the brain contain highly specialised cells
 - The immune system is highly complex
 - Cancer samples have both infiltrating and tumour cells

Introduction

- If a gene is increased 2-fold in expression:
 - Is this 2-fold in 100% of cells?
 - Or is it 4-fold in 50% of cells?
 - Or is it down 2-fold in 25% and up 8-fold in 25% and unchanged in 50%?
- Changes in gene expression can be highly specific to individual cell-types
- In general, determining heterogeneity of our samples is challenging


Introduction

- The most intuitive solution is to obtain RNA from each cell and sequence
- Reality is much trickier than this

Introduction

- The most intuitive solution is to obtain RNA from each cell and sequence
- Reality is much trickier than this
- How do we characterise which cell is which cell-type?
- How do we capture as many transcripts from each cell as we can?
 - Missing values are a huge issue in scRNA-seq
- How do we compare within the same cell-types between experimental groups?
 - E.g., treated and untreated cell types may not be assigned to the same cluster/cell-type

Workflow Outline




scRNA Protocols


Isolating Individual Cells

- Early protocols used a dilution series or manual isolation with a microscope (*micromanipulation*)
- Laser Capture Micro-dissection (LCM)
- Fluorescence-Activated Cell Sorting (FACS)
 - Labelled antibodies to specific surface markers
 - MACS is a magnetic-based approach
- Microfluidics/Droplet-based approaches

Protocol Timeline




Protocol Timeline




IFC Capture

- Integrated Fluidic Circuit (IFC) chips
 - Most common is the Fluidigm C1
- Deliver tiny volumes into 'reaction chambers'
- Early chips had 96 chambers \implies multiple chips / experiment
- Recent chips handle ~ 800 cells


Protocol Timeline



Droplet-based Approaches



Droplet-based Approaches



Flow rate is modelled as a *Poisson* process to minimise doublets

Sequencing Overview

- Individual cells are isolated \implies how do we sequence?
- Need a method to track which reads come from which cell
- Sequencing is performed on a standard Illumina machine, i.e. multiplexed
- Each cell is essentially an individual library prep
 - Barcodes / UMIs are used to identify individual cells

Data Analysis

Spatial Transcriptomics