

MATH 738
HW 2
Due 11/29

- (1) Let \mathcal{C} be an additive category. An ideal of \mathcal{C} is a collection of subspaces

$$I(X, Y) \subseteq \text{Hom}(X, Y)$$

for each pair of objects X, Y such that for any $f : X \rightarrow X'$ and $g : Y \rightarrow Y'$ we have

$$g \circ I(X, Y) \circ f \subseteq I(X', Y').$$

Show that $\bar{\mathcal{C}}$ with the same objects as \mathcal{C} and with

$$\text{Hom}_{\bar{\mathcal{C}}}(X, Y) := \text{Hom}_{\mathcal{C}}(X, Y) / I(X, Y)$$

has the structure of an additive category such that

$$\begin{aligned} X &\mapsto X \\ f &\mapsto \bar{f} \end{aligned}$$

is an essentially surjective and full additive functor.

- (2) Let P, Q be a bounded above complexes of projectives. Directly show that any quasi-isomorphism

$$Q \rightarrow P$$

has an inverse up to homotopy.

- (3) Assume we have a commutative diagram

$$\begin{array}{ccccccc} \cdots & \longrightarrow & A^i & \longrightarrow & B^i & \longrightarrow & C^i & \longrightarrow & A^{i+1} & \longrightarrow & \cdots \\ & & \downarrow f^i & & \downarrow g^i & & \downarrow h^i & & \downarrow f^{i+1} & & \\ \cdots & \longrightarrow & D^i & \longrightarrow & E^i & \longrightarrow & F^i & \longrightarrow & D^{i+1} & \longrightarrow & \cdots \end{array}$$

in an abelian category \mathcal{A} with exact rows. Show that if any two of $f^\bullet, g^\bullet, h^\bullet$ are all isomorphisms, then so is the third.

- (4) Let \mathcal{T} a compactly generated triangulated category and let $\mathcal{P}(X)$ be a predicate depending on objects of \mathcal{T} . Assume that

- if X and Y are isomorphic and $\mathcal{P}(X)$ is true, then $\mathcal{P}(Y)$ is true for Y
- \mathcal{P} is true for a set of compact generators,
- if $X \rightarrow Y \rightarrow Z \rightarrow X[1]$ is a triangle and \mathcal{P} is true for two of the three X, Y , and Z , then \mathcal{P} is true for the third.
- if \mathcal{P} is true for X_a for $a \in A$, then \mathcal{P} is true for $\bigoplus_a X_a$.

Show that \mathcal{P} is true for all objects of \mathcal{T} .

- (5) An *cohomological functor* $H : \mathcal{T}^{op} \rightarrow \text{Ab}$ is a functor that takes triangles to long exact sequences and for which the natural map

$$H \left(\bigoplus_{a \in A} X_a \right) \rightarrow \prod_{a \in A} H(X_a)$$

is an isomorphism. Show that if \mathcal{T} is compactly generated, then any cohomological functor is representable. (Hint: use the resolution constructed in the notes replacing $\text{Hom}(-, X)$ with H .)