

HOMEWORK

- (1) Translate the statement and proof of problem1 in Hw10.lean to a pen-and-paper statement and proof.
- (2) Translate the statement and proof of problem2 in Hw10.lean to a pen-and-paper statement and proof.
- (3) Translate the statement and proof of problem3 in Hw10.lean to a pen-and-paper statement and proof.
- (4) Below is a theorem and proof. Translate the statement as a theorem problem4 and give a proof of it in Lean in Hw10.lean.

Theorem. Let $f : A \rightarrow B$ and $g : B \rightarrow C$ be functions. If $d : B \rightarrow A$ is a left inverse to f and $e : C \rightarrow B$ is a right inverse to g , then $d \circ e$ is a left inverse to $g \circ f$.

Proof. To be a left inverse we need to show that

$$(d \circ e) \circ (g \circ f) = \text{id}_A$$

for all a . Using the fact that composition is associative, we can rewrite the left-hand side as

$$(d \circ e) \circ (g \circ f) = d \circ (e \circ g) \circ f$$

By definition of a right inverse, we know that $e \circ g = \text{id}_B$ so we can rewrite again

$$d \circ (e \circ g) \circ f = d \circ \text{id}_B \circ f$$

Since composition with the identity function is the identity we have

$$d \circ \text{id}_B \circ f = d \circ f$$

Finally, as d is a left inverse to f , we have

$$d \circ f = \text{id}_A$$

□

- (5) Below is a theorem and proof. Translate the statement as a theorem labeled problem5 give a proof of it in Lean in Hw10.lean.

Theorem. Being in bijection is symmetric. More precisely, if there exists a bijection $A \cong B$, then there also exists a bijection $B \cong A$.

Proof. Let $f : A \rightarrow B$ be a bijection. Since f is a bijection, it has an inverse $f^{-1} : B \rightarrow A$ which satisfies $f \circ f^{-1} = \text{id}_B$ and $f^{-1} \circ f = \text{id}_A$. We claim that f^{-1} is also a bijection. But the conditions we just listed imply that f is the inverse of f^{-1} . Since f^{-1} has an inverse, it is a bijection also. □