

# Sampling: Stratified sampling

```
$ echo "Data Sciences Institute"
```

# Learning Outcomes

*How might our study be impacted if we divide our population into groups by shared characteristics before sampling? How do we effectively study a sample selected in this manner?*

- Identify benefits of using stratified random sampling
- Compute sample statistics for stratified random samples
- Design a study using stratified random sampling
- Distinguish between stratified random sampling and quota sampling

# What is stratified sampling?

1. Divide the whole population into non-overlapping subpopulations based on shared characteristics. These subpopulations are called **strata** .
  2. Take independent probability samples (often SRS) from each stratum.
  3. Pool individual samples together to calculate overall population estimates.
- Stratified sampling often requires supplemental information about a population in order to divide it into separate groups.
    - For example if you have a list of all student emails from a university and you want to stratify by gender, this list will need to be linked with a data source that includes each student's gender

# Why stratify?

- Preventing a non-representative sample
- Seeking estimates with known precisions for certain subpopulations
- Convenience and lower cost
- Higher precision (lower variance) estimates for population means and totals

# Sample and Population Sizes

- Suppose we have a population of size  $N$  divided into  $H$  strata. Let  $N_h$  be the number of population units in stratum  $h$ . Then we must have,

$$N_1 + N_2 + \dots + N_H = N$$

- Suppose we then take an SRS from each stratum. Let  $n_h$  represent the size of the sample selected from stratum  $h$ . The total sample size is,

$$n_1 + n_2 + \dots + n_H = n$$

- Sample and population sizes do not have to be equal across all strata

# Sample Mean

- The sample mean for stratum  $h$  can be calculated,

$$\bar{y}_h = \frac{1}{n_h} \sum_{i=1}^{n_h} x_{hi}$$

- To estimate the population mean, the sample mean for the whole sample (from all strata combined) can be calculated,

$$\bar{y} = \sum_{h=1}^H \frac{N_h}{N} \bar{y}_h$$

- This is a weighted mean of all sample strata means.

# Stratum Sample Variance

- The sample variance for the the sample from each stratum can be computed the same way as an SRS:

$$s_h^2 = \sum_{i=1}^{n_h} \frac{(y_{hj} - \bar{y}_h)^2}{n_h - 1}$$

# Estimator Variance and Error

- The variance of the sample mean can be computed,

$$\hat{V}(\bar{y}) = \sum_{h=1}^H \frac{s_h^2}{n_h} \left(1 - \frac{n_h}{N_h}\right) \left(\frac{N_h}{N}\right)^2$$

(how much our mean will vary across samples)

- The standard error (SE) and coefficient of variation (CV) remain the same as for an SRS:

$$SE(\bar{y}) = \sqrt{\hat{V}(\bar{y})}$$

$$CV(\bar{y}) = \frac{SE(\bar{y})}{\bar{y}}$$

# Weights

# Weights

- Remember from SRS: a sample weight is how much a sample is 'worth', how much of our original population it represents
- When using stratified sampling, weights may differ by stratum.
- The inclusion probability for unit  $i$  of stratum  $h$  is,

$$\pi_{hi} = \frac{n_h}{N_h}$$

- Where  $n_h$  is the size of SRS from stratum  $h$  and  $N_h$  is the total number of units in stratum  $h$ .
- As previous, the sampling weight for unit  $i$  of stratum  $h$  is then,

$$w_{hi} = \frac{1}{\pi_{hi}}$$

# Using Sample Weights

- The population mean can be estimated directly using a weighted mean of recorded observations:

$$\bar{y} = \frac{\sum_{h=1}^H \sum_{i=1}^{n_h} w_{hi} y_{hi}}{\sum_{h=1}^H \sum_{i=1}^{n_h} w_{hi}}$$

- In stratified sampling, we need to sum over the weights and units in each stratum, and then sum over all strata.

# Defining Strata & Allocating Observations

# How do you divide your population into strata?

- Mean values should differ greatly between strata
  - Stratify by a variable that is closely related to the variable(s) you are trying to estimate
  - For example, if you wish to estimate average height, you might stratify by age or sex instead of geographic location
- Data availability
  - Is there existing survey data to help you define appropriate strata? If not, are you able to collect preliminary data for this purpose?
  - More supplementary data often means more strata

# How do you divide your population into strata?

- Difficulty and cost
  - More strata may mean a higher cost or effort involved
  - Is this additional cost worthwhile for the precision you wish to achieve or the type of analysis you wish to conduct?

# How many units should you sample from each stratum?

- Proportional Allocation
  - Sample the same proportion of units from each stratum
  - Sample weights ( $\pi_{hi}$ ) are the same for each sampled unit regardless of stratum
- Optimal Allocation
  - Variation among larger sampling units may be greater than variation among smaller sampling units, so a higher proportion of large units should be sampled
  - Useful for businesses, cities, and institutions like schools or hospitals

# How many units should you sample from each stratum?

- Allocation for Precision with Strata
  - Sample to reduce the variation in stratum-level estimates, not population-level estimates
  - Useful when the goal is comparing estimates between strata

# Quota Sampling

# Quota Sampling

- Population is divided into subpopulations like strata
- **Non-probability sampling** is conducted within each subpopulation
  - Often convenience sampling is used
- Specified amounts (**quotas**) of types of units are selected

## Why use quota sampling?

- Probability sampling may be expensive or impractical
- May give better results than a pure convenience sample due to enforced quotas
- Cheaper than probability samples

## Why *not* use quota sampling?

- Prone to selection bias
- Methods of analysis for probability samples do not apply

# Next

- Cluster sampling
- Resampling