

Exercise 1 - RNN for Sentiment Analysis

Suppose we are training a vanilla RNN like below to determine whether a sentence expresses positive or negative sentiment. This RNN will be a character-level RNN where $x^{(1)}, \dots, x^{(T)}$ is the sequence of input characters. The RNN is given as follows:

$$h^{(t)} = \tanh(Ux^{(t)} + Wh^{(t-1)} + b)$$

$$y = \sigma(Vh^{(T)} + d)$$

- How many times do we need to apply the weight matrix U , W , and V ?
- What are the shapes of the matrices U , W , and V ?
- How many addition and multiplication operations are required to make a prediction? You can assume that no addition and multiplications are performed when applying the tanh and sigmoid activation functions.

Solution

- We will need to compute $h^{(t)}$ for $t = 1, \dots, T$. Each of this computation requires applying the weight matrices W and T once. The matrix V is only applied once at the end. Therefore, we need to apply W and U T times each and V once.
- The shape of U is $d_h \times d_x$, the shape of W is $d_h \times d_h$, and the shape of V is $d_y \times d_h$, where d_h is the dimensionality of the $h^{(i)}$ (i.e. $h^{(i)} \in \mathbb{R}^{d_h}$), d_x is the dimensionality of the inputs $x^{(i)}$, and d_y is the dimensionality of the output y .
- For each of the T steps, we need to perform two matrix-vector multiplications (one for $Ux^{(i)}$ and one for $Wh^{(i)}$) and two vector additions. To compute the output, we need one additional matrix-vector multiplication and one vector addition.

Exercise 2 - Scalar RNN

Suppose we have the following vanilla RNN network, where the inputs and hidden units are scalars.

$$h^{(t)} = \tanh(w \cdot h^{(t-1)} + u \cdot x^{(t-1)} + b_h)$$

$$y = \sigma(v \cdot h^{(T)} + b_y)$$

- Show that if $|w| < 1$, and the number of time steps T is large, then the gradient $\frac{\partial y}{\partial x^{(0)}}$ vanishes.
- Why is the result from Part (a) troubling?

Solution

- To make the sequence length T explicit in the notation, we will write y instead of y_T . Formally, what we have to show is

$$|w| < 1 \implies \lim_{T \rightarrow \infty} \frac{\partial y_T}{\partial x^{(0)}} = 0.$$