

STAT565 Final Exam

Winter 2019

Question 1 (Show work to get full points)

Suppose there are 4 levels of a single treatment factor to be compared in a balanced incomplete block design. There are only 6 blocks available each with only 2 experimental units.

(a). Compute the number of times each treatment level appears in the whole design.

4 levels of treatment means $a = 4$; 6 blocks means $b = 6$; 2 experimental units means $k = 2$;

$$N = kb = ra \implies r = kb/a = 2 \times 6 \div 4 = 3$$

(b). Compute the number of times each pair of treatment levels appears in the whole design.

$$\lambda = \frac{r(k-1)}{a-1} = \frac{3(2-1)}{4-1} = 1$$

(c). Provide a random assignment of treatment levels so that columns represent blocks and rows represent treatment levels.

	block1	block2	block3	block4	block5	block6
unit1	1	2	3	4	1	2
unit2	2	3	4	1	3	4


Question 2 (Use software to analyze the given data)

Big sagebrush is often planted in range restoration projects. An experiment is performed to determine the effects of storage length and relative humidity on the viability of seeds. Sixty-three batches of 300 seeds each are randomly divided into 21 groups of three. These 21 groups each receive a different treatment, namely the combinations of storage length (0, 60, 120, 180, 240, 300, or 360 days) and storage relative humidity (0, 32, or 45). After the storage time, the seeds are planted, and the response is the percentage of seeds that sprout. The data are shown above and given in Sagebrush Excel file.

Part I: Analyze these data for the effects of the factors on viability. Do not report code here. Your analysis should include the following:

(a). Plot of the data (a plot with data and/or means of treatment combinations). Do not report code here.

```
## Classes 'tbl_df', 'tbl' and 'data.frame':    63 obs. of  5 variables:
## $ Humidity: num  0 0 0 0 0 0 0 0 0 0 0 0 ...
## $ Days    : num  0 0 0 60 60 60 120 120 120 180 ...
## $ Percent : num  82.1 79 81.9 78.6 80.8 80.5 79.8 79.1 78.2 82.3 ...
## $ h       : Factor w/ 3 levels "0%","32%","45%": 1 1 1 1 1 1 1 1 1 1 ...
## $ d       : Factor w/ 7 levels "0","60","120",...: 1 1 1 2 2 2 3 3 3 4 ...
```



(b). Description of the observed relationship between two factors based on the graph.

The graph show that, when relative humidity is 45%, the average percentage of viability of seeds have obvious difference between different storage length. The longer storage length have lower average percentage of viability of seeds.

When humidity is 0 or 32%, the average percentage have not obvious difference between different storage length.

When storage length is 0 day, the average percentage of viability of seeds has not obvious difference between different relative humidity.

When storage length is 60 to 360 days, the average percentage of viability have significant difference between 45% humidity and other levels of humidity. The higher humidity has lower average percentage of viability of seeds.

The average percentage have not obvious difference between 0 and 32% humidity on any storage length.

The line plot shows that not all lines are parallel. Difference in average percentage between humidity is not same for different storage length. There could be an interaction effect.

(c). Tabular form of the numerical summaries for each treatment combination and factor levels separately.

h=relative humidity; d=storage length (days); h.d=interaction items

h	min	Q1	median	Q3	max	mean	sd	n	missing
0%	75.5	79.1	81.1	82.1	87.9	81.13333	2.670268	21	0
32%	77.6	78.9	81.0	83.1	83.8	80.99524	2.128491	21	0
45%	21.0	40.6	52.9	61.2	83.1	52.43333	17.196114	21	0

* h=relative humidity

d	min	Q1	median	Q3	max	mean	sd	n	missing
0	78.9	80.5	81.9	82.4	83.1	81.33333	1.599219	9	0
60	61.2	66.5	78.3	80.5	83.6	74.33333	8.767554	9	0
120	52.9	59.3	79.1	80.4	83.8	72.68889	11.984203	9	0
180	48.7	54.3	77.8	79.1	82.3	69.97778	13.708462	9	0
240	48.8	52.2	81.1	81.7	83.8	71.64444	15.580606	9	0
300	37.9	40.6	80.3	82.1	87.9	68.00000	21.921451	9	0
360	21.0	25.8	81.0	82.7	84.6	62.66667	29.070690	9	0

* d=storage length (days)

h.d	min	Q1	median	Q3	max	mean	sd	n	missing
0%.0	79.0	80.45	81.9	82.00	82.1	81.00000	1.7349352	3	0
32%.0	80.5	81.45	82.4	82.75	83.1	82.00000	1.3453624	3	0
45%.0	78.9	79.95	81.0	82.05	83.1	81.00000	2.1000000	3	0
0%.60	78.6	79.55	80.5	80.65	80.8	79.96667	1.1930353	3	0
32%.60	78.1	78.20	78.3	80.95	83.6	80.00000	3.1192948	3	0
45%.60	61.2	61.30	61.4	63.95	66.5	63.03333	3.0038864	3	0
0%.120	78.2	78.65	79.1	79.45	79.8	79.03333	0.8020806	3	0
32%.120	80.4	81.10	81.8	82.80	83.8	82.00000	1.7088007	3	0
45%.120	52.9	55.90	58.9	59.10	59.3	57.03333	3.5851546	3	0
0%.180	75.5	77.30	79.1	80.70	82.3	78.96667	3.4019602	3	0
32%.180	77.8	78.30	78.8	79.60	80.4	79.00000	1.3114877	3	0
45%.180	48.7	50.80	52.9	53.60	54.3	51.96667	2.9143324	3	0
0%.240	80.1	80.60	81.1	81.40	81.7	80.96667	0.8082904	3	0
32%.240	81.5	82.60	83.7	83.75	83.8	83.00000	1.3000000	3	0
45%.240	48.8	50.35	51.9	52.05	52.2	50.96667	1.8823744	3	0
0%.300	82.1	83.55	85.0	86.45	87.9	85.00000	2.9000000	3	0
32%.300	77.6	78.95	80.3	81.15	82.0	79.96667	2.2188586	3	0
45%.300	37.9	38.25	38.6	39.60	40.6	39.03333	1.4011900	3	0
0%.360	81.7	82.20	82.7	83.65	84.6	83.00000	1.4730920	3	0
32%.360	78.9	79.95	81.0	82.05	83.1	81.00000	2.1000000	3	0
45%.360	21.0	23.10	25.2	25.50	25.8	24.00000	2.6153394	3	0

* h.d=interaction items

(d). The complete (theoretical, not the estimated) linear model and explain the terms for this experiment.

This is a two-factor factorial model with fixed effect.

$$y_{ijk} = \mu + \tau_i + \beta_j + (\tau\beta)_{ij} + \varepsilon_{jik}$$

for $i = 1, 2, \dots, a$ represents 0, 32%, and 45% humidity; $j = 1, 2, \dots, b$ represents 7 levels of storage length; $k = 1, 2, \dots, n$ represents 3 experimental units.

y_{ijk} is the the average percentage of viability of seeds for the k^{th} Exerimental Unit when i^{th} level of relative humidity and j^{th} level of storage length are applied.

a is the number of levels of relative humidity being compared;

b is the number of levels of storage length being compared;

τ_i is fixed main effect of i^{th} level of relative humidity (Treatment effect of relative humidity);

β_j is fixed main effect of j^{th} level of storage length (Treatment effect of storage length);

$(\tau\beta)_{ij}$ is fixed interaction effect of i^{th} level of relative humidity and j^{th} level of storage length (Interaction effect of relative humidity and storage length)

ε_{jik} is random error for the k^{th} Exerimental Unit when i^{th} level of relative humidity and j^{th} level of storage length are applied.

The model includes below assumptions:

$$\varepsilon_{ijk} \sim iidN(0, \sigma^2); \sum_i^a \tau_i = 0; \sum_j^b \beta_j = 0; \sum_i^a (\tau\beta)_{ij} = 0; \sum_j^b (\tau\beta)_{ij} = 0$$

(e). The complete ANOVA table for the fitted model you have in part (d). Do not report code here.

	Df	Sum Sq	Mean Sq	F value	Pr(>F)
humidity	2	11476	5738	1179	1.271e-37

	Df	Sum Sq	Mean Sq	F value	Pr(>F)
days	6	1789	298.1	61.25	3.504e-19
h:d	12	4154	346.2	71.12	6.511e-24
Residuals	42	204.4	4.867	NA	NA
Total	62	17623.78	284.2546	NA	NA

(f). Conclusion from the test of significant interaction effect along with the p value. What do you recommend as the next step in your analysis? (That is, should you test for main effects or simple effects? Why? Explain clearly and provide reasons).

According to ANOVA table, there is a significant interaction effect from relative humidity and storage length on the average percentage around 5% significance level ($P\text{-value}=6.511 \times e^{-24}$). That means, effect of humidity and effect of storage length on average percentage of viability is not independent. Therefore, simple effects examinations are recommended.

(g). According to your answer in part (f), conclusion(s) from the appropriate tests about the effect of factors along with the p-value(s).

The pairwise test show the simple effects:

- In comparisons of storage length Least Squares Means by humidity,

When 0 humidity applied, **ONLY** 120 versus 300 days, 180 versus 300 days **have** significant difference on the average percentage around 5% significance level ($P\text{-value}=0.0290, 0.0264$ but $P\text{-value}=0.0852, 0.0780$ by Tukey method).

When 32% humidity applied, **NONE** of comparisons of storage length have significant difference on the average percentage around 5% significance level (all $P\text{-value}>0.05$).

When 45% humidity applied, **ONLY** 120 versus 180 days, 180 versus 240 days **have not** significant difference on the average percentage around 5% significance level ($P\text{-value}=0.0966, 0.09977$). Moreover, by **Tukey method**, 60 versus 120 days, 120 versus 240 days also have not significant difference on the average percentage around 5% significance level ($P\text{-value}=0.0815, 0.0747$)


- In comparisons of humidity Least Squares Means by storage length,

When 0 day applied, **NONE** of comparisons of humidity have significant difference on the average percentage around 5% significance level (all $P\text{-value}>0.05$).

When 60 to 360 day applied, **NONE** of 0 versus 32% have significant difference on the average percentage around 5% significance level (all $P\text{-value}>0.05$). **ALL** of 0 versus 45%, 32% versus 45% have significant difference on the average percentage around 5% significance level (all $P\text{-value}<0.05$). The **only exception** is, when 300 day applied, 0 versus 32% have significant difference on the average percentage around 5% significance level ($P\text{-value}=0.0209$ but $P\text{-value}=0.2538$ by Tukey method).

The overall conclusion is that humidity lower than 45% can keep the average percentage of viability of seeds being around 80%. If we cannot control the high humidity, the shorter storage length can give higher average percentage of viability of seeds.

(h). Check the assumptions of the fitted model and if you find any violation of assumptions, recommend solutions to fix them. Clearly explain and provide all the graphs and/or tables used to answer this question here. (Do not provide any code here).



The histogram of residuals shows an acceptable normal distribution. The QQ plot didn't show obvious violation of normality except a few data points are not on the line and flattening at the extremes, which is acceptable.

The plot of studentized residual versus predicted (fitted) value shows that except few outliers, the residuals are evenly distributed about zero at each predicted value (zero mean) and vertical deviations of residuals from zero are about same for each predicted value (constant variance).

The plots of studentized residual versus humidity and storage length levels didn't show obvious violation of zero mean and constant variance.

Part II: The researchers are interested in testing whether the mean percentage of seeds that sprout is significantly different at humidity of 45% compared to average of percentage of seeds that sprout at humidity of 0 and 32% when the storage length is 60 days.

(a). To test this, write a contrast statement in terms of the treatment means.

$$\Gamma : \mu_{32} - \frac{1}{2}(\mu_{12} + \mu_{22}) - 0 \implies \tau_{32} + (\tau\beta)_{32} - \frac{1}{2}(\tau_{12} + (\tau\beta)_{12} + \tau_{22} + (\tau\beta)_{22}) = 0$$

where subscript "32" represents 45% humidity and 60-days storage; "12" represents 0 humidity and 60-days storage; "22" represents 32% humidity and 60-days storage.

(b). test your contrast using the software and provide the conclusion along with p value.

The H_0 for contrast is $\mu_{32} - \frac{1}{2}(\mu_{12} + \mu_{22}) = 0$ or $\tau_{32} + (\tau\beta)_{32} - \frac{1}{2}(\tau_{12} + (\tau\beta)_{12} + \tau_{22} + (\tau\beta)_{22}) = 0$.

The H_1 is they don't equal zero.

The estimated value of the contrast is -16.9 and the adjusted Tukey's p value for testing the above hypotheses for contrast is less than 0.0001, which is small enough. We can reject the H_0 and conclude that the average percentage of viability of seeds for 60-days storage at 45% humidity versus that at 0 and 32% humidity are different at 5% significance level.

Part III: Provide your full code and/or output (only the ones used to answer above questions) here.

```
## # I(a)
## table_sagebrush <- read_xlsx("Sagebrush.xlsx")
## table_sagebrush$h <- factor(table_sagebrush$Humidity, levels = c(0, 32, 45),
```

```

##      labels = c("0%", "32%", "45%"))
## table_sagebrush$d <- factor(table_sagebrush$Days, levels = c(0, 60, 120, 180,
##      240, 300, 360), labels = c("0", "60", "120", "180", "240", "300", "360"))
## str(table_sagebrush)
## ggline(table_sagebrush, "d", "Percent", add = c("mean", "jitter"), shape = "h",
##      color = "h", linetype = "h", ylab = "viability of seeds", xlab = "storage length (day)")
## ggline(table_sagebrush, "h", "Percent", add = c("mean", "jitter"), shape = "d",
##      color = "d", linetype = "d", ylab = "viability of seeds", xlab = "relative humidity")
## # I(c)
## favstats(Percent ~ h, data = table_sagebrush)
## favstats(Percent ~ d, data = table_sagebrush)
## favstats(Percent ~ h + d, data = table_sagebrush)
## # I(e)
## model_sagebrush <- aov(Percent ~ h * d, data = table_sagebrush)
## summary(model_sagebrush)
## # I(g)
h_d <- pairs(lsmeans(model_sagebrush, ~h | d))
d_h <- pairs(lsmeans(model_sagebrush, ~d | h))
kable(h_d, format = "latex", caption = "Humidity Applied; Storage Applied; Adjusted by Tukey M
      kable_styling("condensed", full_width = F, font_size = 6) %>% row_spec(16,
      bold = T, background = "#EAF AF1")
kable(d_h, format = "latex") %>% kable_styling("condensed", full_width = F,
      font_size = 6) %>% row_spec(c(14, 17, 43:63), bold = T) %>% row_spec(c(54,
      58), background = "#EAF AF1")
kable(rbind(d_h, h_d), format = "latex") %>% kable_styling("condensed", full_width = F,
      font_size = 6) %>% row_spec(c(43:63, 68, 69, 71, 72, 74, 75, 77, 78, 80,
      81, 83, 84), bold = T) %>% row_spec(c(49, 54, 55, 58), background = "#EAF AF1")
## # I(h)
## table_sagebrush$student_resid <- rstudent(model_sagebrush)
## ols_plot_resid_hist(lm(Percent ~ h * d, data = table_sagebrush))
## ols_plot_resid_qq(lm(Percent ~ h * d, data = table_sagebrush))
## ols_plot_resid_stud_fit(lm(Percent ~ h * d, data = table_sagebrush))
## ggplot(table_sagebrush, aes(h, student_resid)) + geom_dotplot(binaxis = "y",
##      stackdir = "center", binwidth = 0.1, alpha = 0.6) + theme_light()
## ggplot(table_sagebrush, aes(d, student_resid)) + geom_dotplot(binaxis = "y",
##      stackdir = "center", binwidth = 0.1, alpha = 0.6) + theme_light()
## # II(b)
table_sagebrush$hd <- interaction(table_sagebrush$h, table_sagebrush$d)
kable(summary(contrast(lsmeans(lm(Percent ~ hd, table_sagebrush), "hd"), list(c32_12.22 = c(re
      3), -0.5, -0.5, 1, rep(0, 15))))), adjust = "tukey")) %>% kable_styling("condensed",
      full_width = F, font_size = 6)

```

Table 2: Humidity Applied; Storage Applied; Adjusted by Tukey Method

contrast	d	estimate	SE	df	t.ratio	p.value
0% - 32%	0	-1.0000000	1.801352	42	-0.5551387	0.8444154
0% - 45%	0	0.0000000	1.801352	42	0.0000000	1.0000000
32% - 45%	0	1.0000000	1.801352	42	0.5551387	0.8444154
0% - 32%	60	-0.0333333	1.801352	42	-0.0185046	0.9998112
0% - 45%	60	16.9333333	1.801352	42	9.4003486	0.0000000
32% - 45%	60	16.9666667	1.801352	42	9.4188532	0.0000000
0% - 32%	120	-2.9666667	1.801352	42	-1.6469115	0.2376442
0% - 45%	120	22.0000000	1.801352	42	12.2130513	0.0000000
32% - 45%	120	24.9666667	1.801352	42	13.8599628	0.0000000
0% - 32%	180	-0.0333333	1.801352	42	-0.0185046	0.9998112
0% - 45%	180	27.0000000	1.801352	42	14.9887448	0.0000000
32% - 45%	180	27.0333333	1.801352	42	15.0072494	0.0000000
0% - 32%	240	-2.0333333	1.801352	42	-1.1287820	0.5019991
0% - 45%	240	30.0000000	1.801352	42	16.6541609	0.0000000
32% - 45%	240	32.0333333	1.801352	42	17.7829429	0.0000000
0% - 32%	300	5.0333333	1.801352	42	2.7941981	0.0208809
0% - 45%	300	45.9666667	1.801352	42	25.5178754	0.0000000
32% - 45%	300	40.9333333	1.801352	42	22.7236773	0.0000000
0% - 32%	360	2.0000000	1.801352	42	1.1102774	0.5131671
0% - 45%	360	59.0000000	1.801352	42	32.7531831	0.0000000
32% - 45%	360	57.0000000	1.801352	42	31.6429057	0.0000000

contrast	h	estimate	SE	df	t.ratio	p.value
0 - 60	0%	1.0333333	1.801352	42	0.5736433	0.9972487
0 - 120	0%	1.9666667	1.801352	42	1.0917728	0.9270715
0 - 180	0%	2.0333333	1.801352	42	1.1287820	0.9155098
0 - 240	0%	0.0333333	1.801352	42	0.0185046	1.0000000
0 - 300	0%	-4.0000000	1.801352	42	-2.2205548	0.3067175
0 - 360	0%	-2.0000000	1.801352	42	-1.1102774	0.9214250
60 - 120	0%	0.9333333	1.801352	42	0.5181294	0.9984403
60 - 180	0%	1.0000000	1.801352	42	0.5551387	0.9977063
60 - 240	0%	-1.0000000	1.801352	42	-0.5551387	0.9977063
60 - 300	0%	-5.0333333	1.801352	42	-2.7941981	0.1006614
60 - 360	0%	-3.0333333	1.801352	42	-1.6839207	0.6300978
120 - 180	0%	0.0666667	1.801352	42	0.0370092	1.0000000
120 - 240	0%	-1.9333333	1.801352	42	-1.0732681	0.9324512
120 - 300	0%	-5.9666667	1.801352	42	-3.3123276	0.0290470
120 - 360	0%	-3.9666667	1.801352	42	-2.2020502	0.3161924
180 - 240	0%	-2.0000000	1.801352	42	-1.1102774	0.9214250
180 - 300	0%	-6.0333333	1.801352	42	-3.3493368	0.0263962
180 - 360	0%	-4.0333333	1.801352	42	-2.2390594	0.2974152
240 - 300	0%	-4.0333333	1.801352	42	-2.2390594	0.2974152
240 - 360	0%	-2.0333333	1.801352	42	-1.1287820	0.9155098
300 - 360	0%	2.0000000	1.801352	42	1.1102774	0.9214250
0 - 60	32%	2.0000000	1.801352	42	1.1102774	0.9214250
0 - 120	32%	0.0000000	1.801352	42	0.0000000	1.0000000
0 - 180	32%	3.0000000	1.801352	42	1.6654161	0.6418876
0 - 240	32%	-1.0000000	1.801352	42	-0.5551387	0.9977063
0 - 300	32%	2.0333333	1.801352	42	1.1287820	0.9155098
0 - 360	32%	1.0000000	1.801352	42	0.5551387	0.9977063
60 - 120	32%	-2.0000000	1.801352	42	-1.1102774	0.9214250
60 - 180	32%	1.0000000	1.801352	42	0.5551387	0.9977063
60 - 240	32%	-3.0000000	1.801352	42	-1.6654161	0.6418876
60 - 300	32%	0.0333333	1.801352	42	0.0185046	1.0000000
60 - 360	32%	-1.0000000	1.801352	42	-0.5551387	0.9977063
120 - 180	32%	3.0000000	1.801352	42	1.6654161	0.6418876
120 - 240	32%	-1.0000000	1.801352	42	-0.5551387	0.9977063
120 - 300	32%	2.0333333	1.801352	42	1.1287820	0.9155098
120 - 360	32%	1.0000000	1.801352	42	0.5551387	0.9977063
180 - 240	32%	-4.0000000	1.801352	42	-2.2205548	0.3067175
180 - 300	32%	-0.9666667	1.801352	42	-0.5366341	0.9981014
180 - 360	32%	-2.0000000	1.801352	42	-1.1102774	0.9214250
240 - 300	32%	3.0333333	1.801352	42	1.6839207	0.6300978
240 - 360	32%	2.0000000	1.801352	42	1.1102774	0.9214250
300 - 360	32%	-1.0333333	1.801352	42	-0.5736433	0.9972487
0 - 60	45%	17.9666667	1.801352	42	9.9739919	0.0000000
0 - 120	45%	23.9666667	1.801352	42	13.3048241	0.0000000
0 - 180	45%	29.0333333	1.801352	42	16.1175268	0.0000000
0 - 240	45%	30.0333333	1.801352	42	16.6726655	0.0000000
0 - 300	45%	41.9666667	1.801352	42	23.2973206	0.0000000
0 - 360	45%	57.0000000	1.801352	42	31.6429057	0.0000000
60 - 120	45%	6.0000000	1.801352	42	3.3308322	0.0276928
60 - 180	45%	11.0666667	1.801352	42	6.1435349	0.0000049
60 - 240	45%	12.0666667	1.801352	42	6.6986736	0.0000008
60 - 300	45%	24.0000000	1.801352	42	13.3233287	0.0000000
60 - 360	45%	39.0333333	1.801352	42	21.6689138	0.0000000
120 - 180	45%	5.0666667	1.801352	42	2.8127027	0.0966242
120 - 240	45%	6.0666667	1.801352	42	3.3678414	0.0251553
120 - 300	45%	18.0000000	1.801352	42	9.9924965	0.0000000
120 - 360	45%	33.0333333	1.801352	42	18.3380816	0.0000000
180 - 240	45%	1.0000000	1.801352	42	0.5551387	0.9977063
180 - 300	45%	12.9333333	1.801352	42	7.1797938	0.0000002
180 - 360	45%	27.9666667	1.801352	42	15.5253789	0.0000000
240 - 300	45%	11.9333333	1.801352	42	6.6246551	0.0000010
240 - 360	45%	26.9666667	1.801352	42	14.9702402	0.0000000
300 - 360	45%	15.0333333	1.801352	42	8.3455851	0.0000000

h	d	contrast	estimate	SE	df	t.ratio	p.value
0%	.	0 - 60	1.0333333	1.801352	42	0.5736433	1.0000000
0%	.	0 - 120	1.9666667	1.801352	42	1.0917728	1.0000000
0%	.	0 - 180	2.0333333	1.801352	42	1.1287820	1.0000000
0%	.	0 - 240	0.0333333	1.801352	42	0.0185046	1.0000000
0%	.	0 - 300	-4.0000000	1.801352	42	-2.2205548	1.0000000
0%	.	0 - 360	-2.0000000	1.801352	42	-1.1102774	1.0000000
0%	.	60 - 120	0.9333333	1.801352	42	0.5181294	1.0000000
0%	.	60 - 180	1.0000000	1.801352	42	0.5551387	1.0000000
0%	.	60 - 240	-1.0000000	1.801352	42	-0.5551387	1.0000000
0%	.	60 - 300	-5.0333333	1.801352	42	-2.7941981	0.6557437
0%	.	60 - 360	-3.0333333	1.801352	42	-1.6839207	1.0000000
0%	.	120 - 180	0.0666667	1.801352	42	0.0370092	1.0000000
0%	.	120 - 240	-1.9333333	1.801352	42	-1.0732681	1.0000000
0%	.	120 - 300	-5.9666667	1.801352	42	-3.3123276	0.1604071
0%	.	120 - 360	-3.9666667	1.801352	42	-2.2020502	1.0000000
0%	.	180 - 240	-2.0000000	1.801352	42	-1.1102774	1.0000000
0%	.	180 - 300	-6.0333333	1.801352	42	-3.3493368	0.1444330
0%	.	180 - 360	-4.0333333	1.801352	42	-2.2390594	1.0000000
0%	.	240 - 300	-4.0333333	1.801352	42	-2.2390594	1.0000000
0%	.	240 - 360	-2.0333333	1.801352	42	-1.1287820	1.0000000
0%	.	300 - 360	2.0000000	1.801352	42	1.1102774	1.0000000
32%	.	0 - 60	2.0000000	1.801352	42	1.1102774	1.0000000
32%	.	0 - 120	0.0000000	1.801352	42	0.0000000	1.0000000
32%	.	0 - 180	3.0000000	1.801352	42	1.6654161	1.0000000
32%	.	0 - 240	-1.0000000	1.801352	42	-0.5551387	1.0000000
32%	.	0 - 300	2.0333333	1.801352	42	1.1287820	1.0000000
32%	.	0 - 360	1.0000000	1.801352	42	0.5551387	1.0000000
32%	.	60 - 120	-2.0000000	1.801352	42	-1.1102774	1.0000000
32%	.	60 - 180	1.0000000	1.801352	42	0.5551387	1.0000000
32%	.	60 - 240	-3.0000000	1.801352	42	-1.6654161	1.0000000
32%	.	60 - 300	0.0333333	1.801352	42	0.0185046	1.0000000
32%	.	60 - 360	-1.0000000	1.801352	42	-0.5551387	1.0000000
32%	.	120 - 180	3.0000000	1.801352	42	1.6654161	1.0000000
32%	.	120 - 240	-1.0000000	1.801352	42	-0.5551387	1.0000000
32%	.	120 - 300	2.0333333	1.801352	42	1.1287820	1.0000000
32%	.	120 - 360	1.0000000	1.801352	42	0.5551387	1.0000000
32%	.	180 - 240	-4.0000000	1.801352	42	-2.2205548	1.0000000
32%	.	180 - 300	-0.9666667	1.801352	42	-0.5366341	1.0000000
32%	.	180 - 360	-2.0000000	1.801352	42	-1.1102774	1.0000000
32%	.	240 - 300	3.0333333	1.801352	42	1.6839207	1.0000000
32%	.	240 - 360	2.0000000	1.801352	42	1.1102774	1.0000000
32%	.	300 - 360	-1.0333333	1.801352	42	-0.5736433	1.0000000
45%	.	0 - 60	17.9666667	1.801352	42	9.9739919	0.0000000
45%	.	0 - 120	23.9666667	1.801352	42	13.3048241	0.0000000
45%	.	0 - 180	29.0333333	1.801352	42	16.1175268	0.0000000
45%	.	0 - 240	30.0333333	1.801352	42	16.6726655	0.0000000
45%	.	0 - 300	41.9666667	1.801352	42	23.2973206	0.0000000
45%	.	0 - 360	57.0000000	1.801352	42	31.6429057	0.0000000
45%	.	60 - 120	6.0000000	1.801352	42	3.3308322	0.1522207
45%	.	60 - 180	11.0666667	1.801352	42	6.1435349	0.0000207
45%	.	60 - 240	12.0666667	1.801352	42	6.6986736	0.0000033
45%	.	60 - 300	24.0000000	1.801352	42	13.3233287	0.0000000
45%	.	60 - 360	39.0333333	1.801352	42	21.6689138	0.0000000
45%	.	120 - 180	5.0666667	1.801352	42	2.8127027	0.6248936
45%	.	120 - 240	6.0666667	1.801352	42	3.3678414	0.1370256
45%	.	120 - 300	18.0000000	1.801352	42	9.9924965	0.0000000
45%	.	120 - 360	33.0333333	1.801352	42	18.3380816	0.0000000
45%	.	180 - 240	1.0000000	1.801352	42	0.5551387	1.0000000
45%	.	180 - 300	12.9333333	1.801352	42	7.1797938	0.0000007
45%	.	180 - 360	27.9666667	1.801352	42	15.5253789	0.0000000
45%	.	240 - 300	11.9333333	1.801352	42	6.6246551	0.0000042
45%	.	240 - 360	26.9666667	1.801352	42	14.9702402	0.0000000
45%	.	300 - 360	15.0333333	1.801352	42	8.3455851	0.0000000
.	0	0% - 32%	-1.0000000	1.801352	42	-0.5551387	1.0000000
.	0	0% - 45%	0.0000000	1.801352	42	0.0000000	1.0000000
.	0	32% - 45%	1.0000000	1.801352	42	0.5551387	1.0000000
.	60	0% - 32%	-0.0333333	1.801352	42	-0.0185046	1.0000000
.	60	0% - 45%	16.9333333	1.801352	42	9.4003486	0.0000000
.	60	32% - 45%	16.9666667	1.801352	42	9.4188532	0.0000000
.	120	0% - 32%	-2.9666667	1.801352	42	-1.6469115	1.0000000
.	120	0% - 45%	22.0000000	1.801352	42	12.2130513	0.0000000
.	120	32% - 45%	24.9666667	1.801352	42	13.8599628	0.0000000
.	180	0% - 32%	-0.0333333	1.801352	42	-0.0185046	1.0000000
.	180	0% - 45%	27.0000000	1.801352	42	14.9887448	0.0000000
.	180	32% - 45%	27.0333333	1.801352	42	15.0072494	0.0000000
.	240	0% - 32%	-2.0333333	1.801352	42	-1.1287820	1.0000000
.	240	0% - 45%	30.0000000	1.801352	42	16.6541609	0.0000000
.	240	32% - 45%	32.0333333	1.801352	42	17.7829429	0.0000000
.	300	0% - 32%	5.0333333	1.801352	42	2.7941981	0.6557437
.	300	0% - 45%	45.9666667	1.801352	42	25.5178754	0.0000000
.	300	32% - 45%	40.9333333	1.801352	42	22.7236773	0.0000000
.	360	0% - 32%	2.0000000	1.801352	42	1.1102774	1.0000000
.	360	0% - 45%	59.0000000	1.801352	42	32.7531831	0.0000000
.	360	32% - 45%	57.0000000	1.801352	42	31.6429057	0.0000000

contrast	estimate	SE	df	t.ratio	p.value
c32_12.22	-16.95	1.560016	42	-10.86527	0

Question 3 (Show main steps of your work to get full points)

Let $y_{ijk} = \mu + \tau_i + \beta_j + (\tau\beta)_{ij} + \delta_k + \varepsilon_{ijk}$; $\varepsilon_{ijk} \sim iidN(0, \sigma^2)$; for $i = 1, \dots, a$; $j = 1, \dots, b$; $k = 1, \dots, n$;
Derive the normal equations from the least squares parameter estimation method and estimate the model parameters using the following constraints. $\sum_i^a \hat{\tau}_i = 0$; $\sum_j^b \hat{\beta}_j = 0$; $\sum_i^a \widehat{(\tau\beta)}_{ij} = 0$; $\sum_j^b \widehat{(\tau\beta)}_{ij} = 0$; $\sum_k^n \hat{\delta}_k = 0$

This is a two-factor factorial in RCBD model with fixed effect.

$$SSE = \sum_i^a \sum_j^b \sum_k^n (y_{ijk} - \mu - \tau_i - \beta_j - (\tau\beta)_{ij} - \delta_k)^2$$

Derive

$$\left. \frac{\partial SSE}{\partial \mu} \right|_{\hat{\mu}, \hat{\tau}_i, \hat{\beta}_j, \widehat{(\tau\beta)}_{ij}, \hat{\delta}_k} = 2 \sum_i^a \sum_j^b \sum_k^n (y_{ijk} - \hat{\mu} - \hat{\tau}_i - \hat{\beta}_j - \widehat{(\tau\beta)}_{ij} - \hat{\delta}_k)(-1) = 0$$

For $i = 1, \dots, a$,

$$\left. \frac{\partial SSE}{\partial \tau_i} \right|_{\hat{\mu}, \hat{\tau}_i, \hat{\beta}_j, \widehat{(\tau\beta)}_{ij}, \hat{\delta}_k} = 2 \sum_j^b \sum_k^n (y_{ijk} - \hat{\mu} - \hat{\tau}_i - \hat{\beta}_j - \widehat{(\tau\beta)}_{ij} - \hat{\delta}_k)(-1) = 0$$

For $j = 1, \dots, b$,

$$\left. \frac{\partial SSE}{\partial \beta_j} \right|_{\hat{\mu}, \hat{\tau}_i, \hat{\beta}_j, \widehat{(\tau\beta)}_{ij}, \hat{\delta}_k} = 2 \sum_i^a \sum_k^n (y_{ijk} - \hat{\mu} - \hat{\tau}_i - \hat{\beta}_j - \widehat{(\tau\beta)}_{ij} - \hat{\delta}_k)(-1) = 0$$

For $i = 1, \dots, a$; $j = 1, \dots, b$,

$$\left. \frac{\partial SSE}{\partial (\tau\beta)_{ij}} \right|_{\hat{\mu}, \hat{\tau}_i, \hat{\beta}_j, \widehat{(\tau\beta)}_{ij}, \hat{\delta}_k} = 2 \sum_k^n (y_{ijk} - \hat{\mu} - \hat{\tau}_i - \hat{\beta}_j - \widehat{(\tau\beta)}_{ij} - \hat{\delta}_k)(-1) = 0$$

For $k = 1, \dots, n$,

$$\left. \frac{\partial SSE}{\partial \delta_k} \right|_{\hat{\mu}, \hat{\tau}_i, \hat{\beta}_j, \widehat{(\tau\beta)}_{ij}, \hat{\delta}_k} = 2 \sum_i^a \sum_j^b (y_{ijk} - \hat{\mu} - \hat{\tau}_i - \widehat{(\tau\beta)}_{ij} - \hat{\beta}_j - \hat{\delta}_k)(-1) = 0$$

$$\begin{cases} y_{...} = abn\hat{\mu} + bn \sum_i^a \hat{\tau}_i + an \sum_j^b \hat{\beta}_j + n \sum_i^a \sum_j^b \widehat{(\tau\beta)}_{ij} + ab \sum_k^n \hat{\delta}_k \\ y_{i..} = bn\hat{\mu} + bn\hat{\tau}_i + n \sum_j^b \hat{\beta}_j + n \sum_j^b \widehat{(\tau\beta)}_{ij} + b \sum_k^n \hat{\delta}_k \\ y_{.j.} = an\hat{\mu} + n \sum_i^a \hat{\tau}_i + an\hat{\beta}_j + n \sum_i^a \widehat{(\tau\beta)}_{ij} + a \sum_k^n \hat{\delta}_k \\ y_{ij.} = n\hat{\mu} + n\hat{\tau}_i + n\hat{\beta}_j + n\widehat{(\tau\beta)}_{ij} + \sum_k^n \hat{\delta}_k \\ y_{..k} = ab\hat{\mu} + b \sum_i^a \hat{\tau}_i + a \sum_j^b \hat{\beta}_j + \sum_i^a \sum_j^b \widehat{(\tau\beta)}_{ij} + ab\hat{\delta}_k \end{cases}$$

For $\sum_i^a \hat{\tau}_i = 0$; $\sum_j^b \hat{\beta}_j = 0$; $\sum_i^a \widehat{(\tau\beta)}_{ij} = 0$; $\sum_j^b \widehat{(\tau\beta)}_{ij} = 0$; $\sum_k^n \hat{\delta}_k = 0$,

and for $\hat{\mu}$ is constant, $\hat{\tau}_i, \hat{\beta}_j, \widehat{(\tau\beta)}_{ij}, \hat{\delta}_k$ are constants for summations on other parameters,

and for $y_{...} = abn\bar{y}_{...}$, $y_{i..} = bn\bar{y}_{i..}$, $y_{.j.} = an\bar{y}_{.j.}$, $y_{ij.} = n\bar{y}_{ij.}$, $y_{..k} = ab\bar{y}_{..k}$, then

$$\begin{cases} abn\bar{y}_{...} = abn\hat{\mu} + 0 + 0 + 0 + 0 \\ bn\bar{y}_{i..} = bn\hat{\mu} + bn\hat{\tau}_i + 0 + 0 + 0 \\ an\bar{y}_{.j.} = an\hat{\mu} + 0 + an\hat{\beta}_j + 0 + 0 \\ n\bar{y}_{ij.} = n\hat{\mu} + n\hat{\tau}_i + n\hat{\beta}_j + n(\widehat{\tau\beta})_{ij} + 0 \\ ab\bar{y}_{..k} = ab\hat{\mu} + 0 + 0 + 0 + ab\hat{\delta}_k \end{cases} \implies \begin{cases} \hat{\mu} = \bar{y}_{...} \\ \hat{\tau}_i = \bar{y}_{i..} - \bar{y}_{...} \\ \hat{\beta}_j = \bar{y}_{.j.} - \bar{y}_{...} \\ (\widehat{\tau\beta})_{ij} = \bar{y}_{ij.} - \bar{y}_{i..} - \bar{y}_{.j.} + \bar{y}_{...} \\ \hat{\delta}_k = \bar{y}_{..k} - \bar{y}_{...} \end{cases}$$

Question 4 (Show main steps of your work to get full points)

Consider the two-factor factorial model with fixed effects

$$y_{ijk} = \mu + \tau_i + \beta_j + (\tau\beta)_{ij} + \varepsilon_{ijk}, \text{ for } i = 1, 2, \dots, a; j = 1, 2, \dots, b; k = 1, 2, \dots, n. \quad \varepsilon_{ijk} \sim iidN(0, \sigma^2); \sum_{i=1}^a \tau_i = 0 \text{ and } \sum_{j=1}^b \beta_j = 0; \sum_{i=1}^a (\tau\beta)_{ij} = 0, \sum_{j=1}^b (\tau\beta)_{ij} = 0. \quad \bar{y}_{i..} = \frac{1}{bn} \sum_{j=1}^b \sum_{k=1}^n y_{ijk}; \bar{y}_{.j.} = \frac{1}{an} \sum_{i=1}^a \sum_{k=1}^n y_{ijk}; \bar{y}_{...} = \frac{1}{abn} \sum_{i=1}^a \sum_{j=1}^b \sum_{k=1}^n y_{ijk}; \bar{y}_{...} = \frac{\bar{y}_{...}}{abn}. \quad SS_A = bn \sum_{i=1}^a (\bar{y}_{i..} - \bar{y}_{...})^2; SS_B = an \sum_{j=1}^b (\bar{y}_{.j.} - \bar{y}_{...})^2; SS_{AB} = n \sum_{i=1}^a \sum_{j=1}^b (\bar{y}_{ij.} - \bar{y}_{i..} - \bar{y}_{.j.} + \bar{y}_{...})^2$$

- Method 1: Using the estimation of model parameters

In problem 3, we know $\hat{\mu} = \bar{y}_{...}$, $\hat{\tau}_i = \bar{y}_{i..} - \bar{y}_{...}$, $\hat{\beta}_j = \bar{y}_{.j.} - \bar{y}_{...}$, $(\widehat{\tau\beta})_{ij} = \bar{y}_{ij.} - \bar{y}_{i..} - \bar{y}_{.j.} + \bar{y}_{...}$ for two-factor factorial in RCBD model with fixed effect. It is also true for two-factor factorial model with fixed effect by the same method.

For $\varepsilon_{ijk} \sim iidN(0, \sigma^2)$, $\widehat{\varepsilon}_{ijk} = 0$, $\bar{y}_{...} = \hat{\mu}$.

$$\hat{y}_{ijk} = \hat{\mu} + \hat{\tau}_i + \hat{\beta}_j + (\widehat{\tau\beta})_{ij} + \widehat{\varepsilon}_{ijk} = \bar{y}_{...} + (\bar{y}_{i..} - \bar{y}_{...}) + (\bar{y}_{.j.} - \bar{y}_{...}) + (\bar{y}_{ij.} - \bar{y}_{i..} - \bar{y}_{.j.} + \bar{y}_{...}) + 0 = \bar{y}_{ij.}.$$

Therefore, $\bar{y}_{ij.} - \bar{y}_{...} = \hat{\mu} + \hat{\tau}_i + \hat{\beta}_j + (\widehat{\tau\beta})_{ij} - \hat{\mu} = \hat{\tau}_i + \hat{\beta}_j + (\widehat{\tau\beta})_{ij}$

$$n \sum_{i=1}^a \sum_{j=1}^b (\bar{y}_{ij.} - \bar{y}_{...})^2 = n \sum_{i=1}^a \sum_{j=1}^b (\hat{\tau}_i + \hat{\beta}_j + (\widehat{\tau\beta})_{ij})^2$$

For the two-factor factorial model with fixed effects, all the cross product terms are zero, then

$$\begin{aligned} n \sum_{i=1}^a \sum_{j=1}^b (\hat{\tau}_i + \hat{\beta}_j + (\widehat{\tau\beta})_{ij})^2 &= bn \sum_{i=1}^a \hat{\tau}_i^2 + an \sum_{j=1}^b \hat{\beta}_j^2 + n \sum_{i=1}^a \sum_{j=1}^b (\widehat{\tau\beta})_{ij}^2 \\ &= bn \sum_{i=1}^a (\bar{y}_{i..} - \bar{y}_{...})^2 + an \sum_{j=1}^b (\bar{y}_{.j.} - \bar{y}_{...})^2 + n \sum_{i=1}^a \sum_{j=1}^b (\bar{y}_{ij.} - \bar{y}_{i..} - \bar{y}_{.j.} + \bar{y}_{...})^2 = SS_A + SS_B + SS_{AB} \end{aligned}$$

Therefore,

$$SS_{AB} = n \sum_{i=1}^a \sum_{j=1}^b (\bar{y}_{ij.} - \bar{y}_{...})^2 - SS_A - SS_B$$

- Method 2

$$\bar{y}_{ij.} = \frac{1}{n} \sum_{k=1}^n y_{ijk} = \frac{1}{n} \sum_{k=1}^n (\mu + \tau_i + \beta_j + (\tau\beta)_{ij} + \varepsilon_{ijk}) = \mu + \tau_i + \beta_j + (\tau\beta)_{ij} + \bar{\varepsilon}_{ij.}$$

For $\varepsilon_{ijk} \sim iidN(0, \sigma^2)$, $\sum_{i=1}^a \sum_{j=1}^b \sum_{k=1}^n \varepsilon_{ijk} = 0$, $\sum_{i=1}^a \tau_i = 0$ and $\sum_{j=1}^b \beta_j = 0$; $\sum_{i=1}^a (\tau\beta)_{ij} = 0$, $\sum_{j=1}^b (\tau\beta)_{ij} = 0$

$$\begin{aligned}
\bar{y}_{...} &= \frac{1}{abn} \sum_{i=1}^a \sum_{j=1}^b \sum_{k=1}^n y_{ijk} = \frac{1}{abn} \sum_{i=1}^a \sum_{j=1}^b \sum_{k=1}^n (\mu + \tau_i + \beta_j + (\tau\beta)_{ij} + \varepsilon_{ijk}) \\
&= \mu + \frac{1}{abn} \left[\sum_{j=1}^b \sum_{k=1}^n \left(\sum_{i=1}^a \tau_i \right) + \sum_{i=1}^a \sum_{k=1}^n \left(\sum_{j=1}^b \beta_j \right) + \sum_{k=1}^n \sum_{i=1}^a \sum_{j=1}^b (\tau\beta)_{ij} + \sum_{i=1}^a \sum_{j=1}^b \sum_{k=1}^n \varepsilon_{ijk} \right] = \mu \\
\bar{y}_{i..} &= \frac{1}{bn} \sum_{j=1}^b \sum_{k=1}^n y_{ijk} = \frac{1}{bn} \sum_{j=1}^b \sum_{k=1}^n (\mu + \tau_i + \beta_j + (\tau\beta)_{ij} + \varepsilon_{ijk}) \\
&= \mu + \tau_i + \frac{1}{bn} \sum_{k=1}^n \sum_{j=1}^b \beta_j + \frac{1}{bn} \sum_{k=1}^n \sum_{j=1}^b (\tau\beta)_{ij} + \frac{1}{bn} \sum_{k=1}^n \sum_{j=1}^b \varepsilon_{ijk} = \mu + \tau_i + \bar{\varepsilon}_{i..} \\
\bar{y}_{.j.} &= \frac{1}{an} \sum_{i=1}^a \sum_{k=1}^n y_{ijk} = \frac{1}{an} \sum_{i=1}^a \sum_{k=1}^n (\mu + \tau_i + \beta_j + (\tau\beta)_{ij} + \varepsilon_{ijk}) \\
&= \mu + \frac{1}{an} \sum_{k=1}^n \sum_{i=1}^a \tau_i + \beta_j + \frac{1}{an} \sum_{k=1}^n \sum_{i=1}^a (\tau\beta)_{ij} + \frac{1}{an} \sum_{k=1}^n \sum_{i=1}^a \varepsilon_{ijk} = \mu + \beta_j + \bar{\varepsilon}_{.j.}
\end{aligned}$$

For $\tau_i, \beta_j, (\tau\beta)_{ij}, \varepsilon_{ijk}$ are independent, all the cross product terms are zero. Then,

$$SS_A = bn \sum_{i=1}^a (\bar{y}_{i..} - \bar{y}_{...})^2 = bn \sum_{i=1}^a (\mu + \tau_i + \bar{\varepsilon}_{i..} - \mu)^2 = bn \sum_{i=1}^a (\tau_i + \bar{\varepsilon}_{i..})^2 = n \sum_{i=1}^a \sum_{j=1}^b (\tau_i^2 + \bar{\varepsilon}_{i..}^2)$$

$$SS_B = an \sum_{j=1}^b (\bar{y}_{.j.} - \bar{y}_{...})^2 = an \sum_{j=1}^b (\mu + \beta_j + \bar{\varepsilon}_{.j.} - \mu)^2 = n \sum_{i=1}^a \sum_{j=1}^b (\beta_j^2 + \bar{\varepsilon}_{.j.}^2)$$

$$n \sum_{i=1}^a \sum_{j=1}^b (\bar{y}_{ij.} - \bar{y}_{...})^2 = n \sum_{i=1}^a \sum_{j=1}^b (\mu + \tau_i + \beta_j + (\tau\beta)_{ij} + \bar{\varepsilon}_{ij.} - \mu)^2 = n \sum_{i=1}^a \sum_{j=1}^b (\tau_i^2 + \beta_j^2 + (\tau\beta)_{ij}^2 + \bar{\varepsilon}_{ij.}^2)$$

$$\begin{aligned}
n \sum_{i=1}^a \sum_{j=1}^b (\bar{y}_{ij.} - \bar{y}_{...})^2 - SS_A - SS_B &= n \sum_{i=1}^a \sum_{j=1}^b (\tau_i^2 + \beta_j^2 + (\tau\beta)_{ij}^2 + \bar{\varepsilon}_{ij.}^2) - n \sum_{i=1}^a \sum_{j=1}^b (\tau_i^2 + \bar{\varepsilon}_{i..}^2) - \\
&n \sum_{i=1}^a \sum_{j=1}^b (\beta_j^2 + \bar{\varepsilon}_{.j.}^2) = n \sum_{i=1}^a \sum_{j=1}^b ((\tau\beta)_{ij}^2 + \bar{\varepsilon}_{ij.}^2 - \bar{\varepsilon}_{i..}^2 - \bar{\varepsilon}_{.j.}^2)
\end{aligned}$$

$$\bar{y}_{ij.} - \bar{y}_{i..} - \bar{y}_{.j.} + \bar{y}_{...} = \mu + \tau_i + \beta_j + (\tau\beta)_{ij} + \bar{\varepsilon}_{ij.} - (\mu + \tau_i + \bar{\varepsilon}_{i..}) - (\mu + \beta_j + \bar{\varepsilon}_{.j.}) + \mu = (\tau\beta)_{ij} + \bar{\varepsilon}_{ij.} - \bar{\varepsilon}_{i..} - \bar{\varepsilon}_{.j.}$$

$$SS_{AB} = n \sum_{i=1}^a \sum_{j=1}^b (\bar{y}_{ij.} - \bar{y}_{i..} - \bar{y}_{.j.} + \bar{y}_{...})^2 = n \sum_{i=1}^a \sum_{j=1}^b [(\tau\beta)_{ij} + \bar{\varepsilon}_{ij.} - \bar{\varepsilon}_{i..} - \bar{\varepsilon}_{.j.}]^2 = n \sum_{i=1}^a \sum_{j=1}^b ((\tau\beta)_{ij}^2 + \bar{\varepsilon}_{ij.}^2 - \bar{\varepsilon}_{i..}^2 - \bar{\varepsilon}_{.j.}^2)$$

Therefore,

$$SS_{AB} = n \sum_{i=1}^a \sum_{j=1}^b (\bar{y}_{ij.} - \bar{y}_{i...})^2 - SS_A - SS_B$$

- Method 3: Using the cross products of being zero

Proof three cross-product terms equal 0

$$n \sum_{i=1}^a \sum_{j=1}^b (\bar{y}_{i..} - \bar{y}_{...}) = \sum_{i=1}^a y_{i..} - y_{...} = y_{...} - y_{...} = 0$$

$$n \sum_{i=1}^a \sum_{j=1}^b (\bar{y}_{.j.} - \bar{y}_{...}) = \sum_{j=1}^b y_{.j.} - y_{...} = y_{...} - y_{...} = 0$$

$$n \sum_{i=1}^a \sum_{j=1}^b (\bar{y}_{ij.} - \bar{y}_{i..} - \bar{y}_{.j.} + \bar{y}_{...}) = \sum_{i=1}^a \sum_{j=1}^b y_{ij.} - \sum_{i=1}^a y_{i..} - \sum_{j=1}^b y_{.j.} + y_{...} = y_{...} - y_{...} - y_{...} + y_{...} = 0$$

$$\begin{aligned} 2n \sum_{i=1}^a \sum_{j=1}^b (\bar{y}_{ij.} - \bar{y}_{i..} - \bar{y}_{.j.} + \bar{y}_{...})(\bar{y}_{i..} - \bar{y}_{...}) &= 2n \sum_{i=1}^a \sum_{j=1}^b (\bar{y}_{ij.} - \bar{y}_{i..} - \bar{y}_{.j.} + \bar{y}_{...})(\bar{y}_{.j.} - \bar{y}_{...}) = \\ 2n \sum_{i=1}^a \sum_{j=1}^b (\bar{y}_{i..} - \bar{y}_{...})(\bar{y}_{.j.} - \bar{y}_{...}) &= 0 \end{aligned}$$

Expand the sum-of-squares terms

$$\begin{aligned} n \sum_{i=1}^a \sum_{j=1}^b (\bar{y}_{ij.} - \bar{y}_{...})^2 &= n \sum_{i=1}^a \sum_{j=1}^b (\bar{y}_{ij.} - \bar{y}_{i..} - \bar{y}_{.j.} + \bar{y}_{...} + \bar{y}_{i..} - \bar{y}_{...} + \bar{y}_{.j.} - \bar{y}_{...})^2 \\ &= n \sum_{i=1}^a \sum_{j=1}^b (\bar{y}_{ij.} - \bar{y}_{i..} - \bar{y}_{.j.} + \bar{y}_{...})^2 + n \sum_{i=1}^a \sum_{j=1}^b (\bar{y}_{i..} - \bar{y}_{...})^2 + n \sum_{i=1}^a \sum_{j=1}^b (\bar{y}_{.j.} - \bar{y}_{...})^2 + 0 + 0 + 0 \\ &= SS_{AB} + bn \sum_{i=1}^a (\bar{y}_{i..} - \bar{y}_{...})^2 + an \sum_{j=1}^b (\bar{y}_{.j.} - \bar{y}_{...})^2 = SS_{AB} + SS_A + SS_B \end{aligned}$$

Therefore,

$$SS_{AB} = n \sum_{i=1}^a \sum_{j=1}^b (\bar{y}_{ij.} - \bar{y}_{...})^2 - SS_A - SS_B$$