

$U \sim Geom(p = \frac{1}{2}), u = 1, 2..$ the number of trials needed to get the first head.
 $V \sim NBin(p = \frac{1}{2}, r = 2), v = 2, 3..$ the number of trials needed to get two heads in repeated tosses of a fair coin.
the distribution of (U, V) is $\{(u, v) : u = 1, 2, ..; v = u + 1, u + 2, ..\}$ is not a cross-product set. U and V are not independent.

Conditional expectations and variances

$$f(x_2|x_1) = \frac{f(x_1,x_2)}{f_1(x_1)}; p(x_2|x_1) = \frac{p(x_1,x_2)}{p_1(x_1)}$$

$$4.2.1 \; f(y|x) = P(Y = y|X = x) = \frac{f(x,y)}{f_X(x)}$$

$$4.2.3 \; E(g(x_2)|x_1) = \sum_{all \; x_2} g(x_2)p(x_2|x_1), E(g(x_2)|x_1) = \int_{all \; x_2} g(x_2)f(x_2|x_1)dx_2$$

$$4.2.4 \; Var(y|x) = E(Y^2|x) - (E(Y|x))^2,$$

$$V[X|Y] = E[(X - E[X|Y])^2|Y]$$

$$4.2.7 \; f(x,y) = g(x)h(y) \iff \text{indep}$$

$$4.2.10 \; E(g(X)h(Y)) = \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} g(x)h(y)f(x,y)dxdy =$$

$$\int_{-\infty}^{\infty} h(y)f_Y(y) \left[\int_{-\infty}^{\infty} g(x)f_X(x)dx \right] dy = E[g(x)]E[h(y)] = (Eg(X))(Eh(Y))$$

$$F_{XY}(x,y) = P(X \leq x, Y \leq y)$$

Covariance and Correlation

Transformation of pairs of r.v.s

$$f_{Y_1,Y_2}(y_1,y_2) = \sum_{i=1}^k f_{X_1,X_2}(h_1(y_1,y_2), h_2(y_1,y_2))|J_i|$$

$$J_{1,2} = \begin{vmatrix} \frac{\partial h_1}{\partial y_1} & \frac{\partial h_2}{\partial y_1} \\ \frac{\partial h_1}{\partial y_2} & \frac{\partial h_2}{\partial y_2} \end{vmatrix} = \frac{\partial h_1}{\partial y_1} \frac{\partial h_2}{\partial y_2} - \frac{\partial h_2}{\partial y_1} \frac{\partial h_1}{\partial y_2}$$

Properties of \bar{X} and S^2

$$\bar{X}_{n+1} = \frac{\sum_{i=1}^{n+1} X_i}{n+1} = \frac{\sum_{i=1}^n X_i + X_{n+1}}{n+1} = \frac{n\bar{X}_n + X_{n+1}}{n+1}$$

$$nS_{n+1}^2 = (n-1)S_n^2 + (\frac{n}{n+1})(X_{n+1} - \bar{X}_n)^2$$

$$\text{when } X_1,.., X_n \text{ be iid } n(\mu, \sigma^2),$$

$$\frac{(n-1)S^2}{\sigma^2} \sim \chi_{n-1}^2, Var \chi_{n-1}^2 = 2(n-1)$$

$$Var[\frac{(n-1)S^2}{\sigma^2}] = \frac{(n-1)}{\sigma^4} Var[S^2] = 2(n-1) \implies Var[S^2] = \frac{2\sigma^4}{n-1}$$

Connection between N, χ^2, t, F

| | | | |
|------------|---|-----------------------------------|---|
| W Normal | $\bar{X} \sim N(\mu, \frac{\sigma^2}{n})$ | $\bar{X}, S^2 \text{ indep}$ | $\frac{(n-1)S^2}{\sigma^2} \sim \chi_{n-1}^2$ |
| W/o Normal | $E[\bar{X}] = \mu$ | $V[\bar{X}] = \frac{\sigma^2}{n}$ | $E[S^2] = \sigma^2$ |

5.2.8 Distribution of the mean: $\bar{X} \sim gamma(n\alpha, \beta/n)$ 5.2.9

Convolution formula $Z = X + Y \; f_Z(z) = \int_{-\infty}^{\infty} f_X(w)f_Y(z-w)dw$

5.3.2 $Z \sim n(0, 1), Z^2 \sim \chi_1^2, X_1..X_n \text{ indep}$

$X_i \sim \chi_{p_i}^2, X_1 = .. = X_n \sim \chi_{p_1+..p_n}^2$

$\chi_p^2 \sim gamma(p/2, 2)$

5.3.4 T

5.3.5 $X_1..X_n, Y_1..Y_m \text{ indep.}$

$X_i \sim n(\mu_X, \sigma_X^2) Y_j \sim n(\mu_Y, \sigma_Y^2), \frac{S_X^2}{\sigma_X^2} \sim \chi^2_F: \frac{S_X^2/\sigma_X^2}{S_Y^2/\sigma_Y^2}$

Order statistics

5.4.6 $f_{X(i), X(j)}(u, v) =$

$$\frac{n!}{(i-1)!(j-1-i)!(n-j)!} f_X(u)f_X(v)[F_X(u)]^{i-1}[F_X(v) - F_X(u)]^{j-1-i}[1 - F_X(v)]^{n-j}$$

$$f_{X(1),...,X(n)}(x_1,..,x_n) =$$

$$\begin{cases} n!f_X(x_1) \cdot ... \cdot f_X(x_n) & -\infty < x_1 < .. < x_n < \infty \\ 0 & \text{otherwise} \end{cases}$$

Conditional/Baye's: $P(A|B) = \frac{P(A \cap B)}{P(B)} = \frac{P(B|A)P(A)}{P(B)}$;disjoint

$$P(B) = P(B|A_1)P(A_1) + ..P(B|A_k)P(A_k) = \sum_{i=1}^k P(B|A_i)P(A_i)$$

Independence:

$$\begin{array}{l|l} P(A \cap B) = P(A)P(B) & P(X \leq x, Y \leq y) = P(X \leq x)P(Y \leq y) \\ E(XY) = E(X)E(Y) & F_{X,Y}(x,y) = F_X(x)F_Y(y) \\ V(X \pm Y) = VX + VY & f_{X,Y}(s,t) = f_X(s)f_Y(t) \end{array}$$

pdf/pmf

$$\text{CDF: } F(x) = P(X \leq x) = \sum_{y: y \leq x} p(y) = \int_{-\infty}^x f(y)dy$$

probabilities: $a \leq b$:

PMF: $p(x) = P(X = x) = P(\forall w \in \mathcal{W} : X(w) = x)$

$P(a \leq X \leq b) = F(b) - F(a^-)$; $P(a < X \leq b) = F(b) - F(a)$;

$P(a \leq X \leq a) = p(a)$; $P(a < X < b) = F(b^-) - F(a)$; (where a^- is the largest possible X value strictly less than a); Taking $a = b$ yields $P(X = a) = F(a) - F(a - 1)$ as desired.

PDF: $P(\forall w \in \mathcal{W} : a \leq X(w) \leq b) = \int_a^b f(x)dx$

$P(a \leq X \leq b) = P(a < X \leq b) = P(a < X < b)$;

$P(X > a) = 1 - F(a)$; $P(a \leq X \leq b) = F(b) - F(a)$

CDF

Condition: $f(x) \geq 0 \forall x$ pmf $\sum_x f_X(x) = 1, pdf \int_{-\infty}^{\infty} f(x)dx = 1$

$$\begin{array}{l|l} c \sin x \geq 0, 0 < x < \pi/2 & \left| \int_0^{\frac{\pi}{2}} c \sin x dx = 1 \right. \\ ce^{-|x|} \geq 0, -\infty < x < \infty & \left| c \int_{-\infty}^0 e^x dx + c \int_0^{\infty} e^{-x} dx = 1 \right. \end{array} \quad \begin{array}{l} c = 1 \\ c = \frac{1}{2} \end{array}$$

mean/variance

$$E[g(x)] = \mu = \sum_{x \in D} h(x)p(x) = \int_{-\infty}^{\infty} g(x)f(x)dx$$

$$E[(X - \mu)^n] = \mu_n = \sum (x - \mu)^n p(x) = \int (x - \mu)^n f(x)dx$$

$$\begin{array}{l|l} E(aX + b) = aE(X) + b & E(X) = \int_{-\infty}^0 F_X(t)dt + \int_0^{\infty} F_X(t)dt \\ V(aX + b) = a^2\sigma^2 & V(X \pm Y) = VX + VY \pm 2Cov(X, Y) \\ \sigma_{ax+b} = |a| \cdot \sigma_x & Cov(x, y) = E(XY) - E(X)E(Y) \end{array}$$

$$V(X) = \sigma_x^2 = E(X^2) - [E(X)]^2$$

| $f_X(x)$ $\frac{1}{n}$ | EX $\sum_1^n x \frac{1}{n}$ $\frac{1}{n} \frac{n(n+1)}{2}$ | EX^2 $\sum_1^n x^2 \frac{1}{n}$ $\frac{1}{n} \frac{n(n+1)(2n+1)}{6}$ | $V(X)$ $\frac{n^2-1}{12}$ |
|---------------------------|---|--|---|
| ax^{a-1} | $\int_0^1 axa^{a-1}dx$ $\frac{a}{a+1}x^{a+1} _0^1$ | $\int_0^1 x^2ax^{a-1}dx$ $\frac{a}{a+2}x^{a+2} _0^1$ | $\frac{a}{a+2} - (\frac{a}{a+1})^2$ $\frac{a}{(a+2)(a+1)^2}$ |
| $\frac{3}{2}(x-1)^2$ | $\int_0^2 x\frac{3}{2}(x-1)^2dx$ $\frac{3}{2}(\frac{x^4}{4} _0^2..)$ | $\int_0^2 x^2\frac{3}{2}(x-1)^2dx$ $\frac{3}{2}(\frac{x^5}{5} _0^2..)$ | $\frac{8}{5} - 1^2$ $\frac{3}{5}$ |

transform

$$\begin{array}{l|l} g(x) \uparrow & F_Y(y) = F_X(g^{-1}(y)) \\ g(x) \downarrow & F_Y(y) = 1 - F_X(g^{-1}(y)) \end{array}$$

not monotone $\because X \leq 0 \text{ is } \emptyset \therefore P(X \leq -\sqrt{y}) = 0$

$$F_Y(y) = P(Y \leq y) = P(X^2 \leq y) = P(-\sqrt{y} \leq X \leq \sqrt{y}) = P(X \leq \sqrt{y}) - P(X \leq -\sqrt{y}) = F_X(\sqrt{y}) = \sqrt{y}, \; 0 < \sqrt{y} < 1$$

$$\begin{array}{l|l|l|l} Y & X & f_X(x) & F_Y(y) \\ & \sqrt{Y} & \frac{e^{-x^2/2}}{\sqrt{2\pi}} & F_X(\sqrt{y}) - F_X(-\sqrt{y}) \end{array} \quad \begin{array}{l} f_Y(y) = \frac{dF_Y(y)}{dy} \\ \frac{e^{-y/2}}{2\sqrt{2\pi y}}, y > 0 \end{array}$$

monotone: $f_Y(y) = f_X(g^{-1}(y))|\frac{d(g^{-1}(y))}{dy}|$

| | Y | X | $f_X(x)$ |
|---|------------|-----------------------|--|
| 1 | $-\log X$ | e^{-Y} | $x^n(1-x)^m, 0 < x < 1$ |
| 2 | e^X | $\ln Y$ | $\frac{1}{\sigma^2}xe^{-(x/\sigma)^2/2}, 0 < x < \infty$ |
| 3 | $-2\ln X$ | $e^{-Y/2}$ | $1, 0 < x < 1$ |
| 4 | X^2 | \sqrt{Y} | $1, 0 < x < 1$ |
| 5 | $d \tan X$ | $\arctan \frac{Y}{d}$ | $\frac{1}{b-a} = \frac{2}{\pi}$ |

$$1 \; (e^{-y})^n(1 - e^{-y}m)|\frac{d(e^{-y})}{dy}| = e^{-(n+1)y}(1 - e^{-y})^m, \; 0 < y < \infty$$

$$2 \; \frac{1}{\sigma^2} \ln ye^{-(\ln y/\sigma)^2/2}|\frac{d(\ln y)}{dy}| = \frac{\ln(y)}{\sigma^2y}e^{-(\ln(y)/\sigma)^2/2}, 1 < y < \infty$$

$$3 \; 1|\frac{de^{-Y/2}}{dy}| = \frac{1}{2}e^{-Y/2}, 0 < y < \infty$$

$$4 \; 1|\frac{d\sqrt{y}}{dy}| = \frac{1}{2\sqrt{y}}, 0 < y < 1$$

$$5 \; \frac{2}{\pi}|\frac{d(\arctan \frac{y}{d})}{dy}| = \frac{2}{\pi} \frac{d(\frac{y}{d})}{[(\frac{y}{d})^2+1]} = \frac{2}{\pi d[(\frac{y}{d})^2+1]} 0 < y < \infty$$

find mean/variance by mgf

$$E(e^{tx}) = \int e^{tx}f(x)dx = \sum e^{tx}f(x); M_{aX+b}(t) = e^{tb}M_{aX}(t)$$

$$\begin{array}{l|l} M_X(t) = E(e^{tx}) & M_X(0) = 1 \\ M_X'(t) = E(xe^{tx}) & M_X'(0) = E(X) \\ M_X''(t) = E(x^2e^{tx}) & M_X''(0) = E(X^2) \\ M_X^n(t) = E(x^ne^{tx}) & M_X^n(0) = E(X^n) \end{array}$$

Series

| Finite | Binomial |
|---|---|
| $\sum_{k=1}^n k = \frac{n(n+1)}{2}$ | $\sum_{k=0}^n \binom{n}{k} = 2^n$ |
| $\sum_{k=1}^n (2k-1) = n^2$ | $\sum_{k=0}^n \binom{r+k}{k} = \binom{r+n+1}{n}$ |
| $\sum_{k=1}^n k^2 = \frac{n(n+1)(2n+1)}{6}$ | $\sum_{k=0}^n \binom{k}{m} = \binom{n+1}{m+1}$ |
| $\sum_{k=1}^n k^3 = (\frac{n(n+1)}{2})^2$ | $\sum_{k=0}^r \binom{m}{k} \binom{n}{r-k} = \binom{m+n}{r}$ |
| $\sum_{k=0}^n c^k = \frac{c^{n+1}-1}{c-1}$ | $\sum_{k=0}^n \binom{n}{k} a^{n-k}b^k = (a+b)^n$ |
| Infinite $ p < 1$ | $\sum_{k=0}^{\infty} kp^{k-1} = \frac{d}{dp} (\sum_{k=0}^{\infty} p^k) = \frac{1}{(1-p)^2}$ |
| $\sum_{k=0}^{\infty} p^k = \frac{1}{1-p}$ | $\sum_{k=0}^{\infty} \binom{r+k-1}{k} x^k = (1-x)^{-r} r \in \mathbb{N}^+$ |
| $\sum_{k=1}^{\infty} p^k = \frac{p}{1-p}$ | $\sum_{k=0}^{\infty} \binom{\alpha}{k} p^k = (1+p)^\alpha \quad p < 1, \alpha \in \mathbb{C}$ |
| $\Gamma(n) = (n-1)!$ | $\sum_{n=0}^{\infty} \frac{x^n}{n!} = e^x$ |
| $\Gamma(a+1) = a\Gamma(a)$ | $\sum_{n=0}^{\infty} ar^n = \frac{a}{1-r}, r < 1$ |
| $\Gamma(1/2) = \sqrt{\pi}$ | $\sum_{k=0}^{n-1} ar^n = \frac{a(1-r^n)}{1-r}$ |
| $\Gamma(0) = \Gamma(-1) = \infty$ | $\int_0^1 x^{a-1}(1-x)^{b-1}, dx = \frac{\Gamma(a)\Gamma(b)}{\Gamma(a+b)}$ |
| $\Gamma(-1/2) = -2\Gamma(1/2)$ | $\int_0^{\infty} x^{t-1}e^{-x}dx = \Gamma(t)$ |

Integrals+c

$$\begin{array}{l|l|l} \int kdx = kx & \int_a^b cdx = c(b-a) & \left| \int_a^b f(x)dx \right| \leq \int_a^b |f(x)|dx \\ \int e^udu = e^u & \int \ln udu = u\ln(u) - u & \int \frac{1}{a^2+u^2}du = \frac{1}{a}\tan^{-1}(\frac{u}{a}) \\ \int x^n = \frac{x^{n+1}}{n+1} & \int \frac{1}{ax+b} = \frac{1}{a}\ln|ax+b| & \int \frac{1}{\sqrt{a^2-u^2}}du = \frac{1}{\sin(\frac{u}{a})} \\ \int \ln aa^x = a^x & & \end{array}$$

Substitution

$$\begin{array}{l|l} u = g(x) & \begin{array}{l} du = g'(x)dx \quad \int_a^b f(g(x))g'(x)dx = \int_{f(a)}^{f(b)} f(u)du \\ x^3 \quad \quad \quad du = 3x^2dx \quad \int_1^2 5x^2\cos(x^3)dx = \int_1^8 \frac{5}{3}\cos(u)du \end{array} \end{array}$$

Integration by parts

$$\begin{array}{l|l|l|l|l} u & du & dv & v & \int_a^b u dv \\ x & dx & e^{-x} & -e^{-x} & \int xe^{-x}dx \\ \ln x & \frac{1}{x}dx & dx & x & \int_3^5 x \ln x dx \end{array}$$

$$= uv|_a^b - \int_a^b vdu$$

$$= -xe^{-x} + \int xe^{-x}dx = -xe^{-x} - e^{-x} + c$$

$$= x\ln x|_3^5 - \int_3^5 dx = (x\ln x - x)|_3^5 = 5\ln 5 - 3\ln 3 - 2$$

$$\int_a^b f(x)dx = F(b) - F(a) = -\int_b^a f(x)dx = \int_a^c f(x)dx + \int_c^b f(x)dx;$$

$$\frac{d}{dx} \int_{v(x)}^{u(x)} f(t)dt = u'(x)f[u(x)] - v'(x)f[v(x)]$$

Derivatives

$$\begin{array}{l|l|l} (cf)' = cf'(x) & (fg)' = f'g + fg' & (f \pm g)' = f'(x) \pm g'(x) \\ \frac{dx}{dx} = 1 & (\frac{f}{g})' = \frac{(fg'-fg'g)}{g^2} & (f(g(x)))' = f'(g(x))g'(x) \\ \frac{de^x}{dx} = e^x & \frac{d(x^n)}{dx} = nx^{n-1} & \frac{d\ln(x)}{dx} = \frac{1}{x}, x > 0 \\ \frac{da^x}{dx} = a^x \ln(a) & \frac{d\ln^{-1}x}{dx} = \frac{1}{1+x^2} & \frac{d\log_a(x)}{dx} = \frac{1}{x\ln a}, x > 0 \\ \frac{d\cos x}{dx} = -\sin x & \frac{d\tan x}{dx} = \sec^2 x & \frac{d x(\ln x-1)}{dx} = \ln x \end{array}$$

| Distribution | CDF | P(X=x),f(x) | μ | EX^2 | Var | MGF | M'(t) | M''(t) | $M^n(t)$ |
|-------------------------------|--|---|--------------------------------|-------------------------------|---|--|----------------------------|--|---|
| Bern(p) | | $p^x q^{1-x}, x \in \{1, 0\}$ | p | p | pq | $pe^t + q$ | | | |
| Bin(n, p) | $I_{1-p}(n-x, x+1)$ | $\binom{n}{x} p^x q^{n-x}; x \in \{0, 1..n\}$ | np | $\mu(\mu + q)$ | $q\mu$ | $(pe^t + q)^n$ | | | |
| Geom(p) | $1 - q^x$ | $pq^{x-1}, x \in 1, 2, ..$ | $\frac{1}{p}$ | $\frac{p+2q}{p^2}$ | $\frac{q}{p^2}$ | $\frac{pe^t}{1-qe^t}, t < -\ln q$ | | | |
| | $1 - q^{x+1}$ | $pq^x, x \in 0, 1, ..$ | $\frac{q}{p}$ | $\frac{q^2+q}{p^2}$ | $\frac{q}{p^2}$ | $\frac{p}{1-qe^t}, qe^t < 1$ | $\frac{pqe^t}{(1-qe^t)^2}$ | $\frac{2pqe^t}{(1-qe^t)^3} - M'(t)$ | |
| NBin(r, p) | | $\binom{x-1}{r-1} p^r q^{x-r}, x \in r, r+1..$ | $\frac{r}{p}$ | $\frac{r}{p}$ | $\frac{rq}{p^2}$ | $(\frac{pe^t}{1-qe^t})^r$ | | | |
| | | $\binom{x+r-1}{r-1} p^r q^x, x \in 0, 1..$ | $\frac{rq}{p}$ | $\frac{rq}{p^2}$ | $\frac{rq}{p^2}$ | $(\frac{p}{1-qe^t})^r, qe^t < 1$ | | | |
| HGeom(N, m, k) | | $\frac{\binom{m}{x} \binom{N-m}{k-x}}{\binom{N}{k}}$ | $\frac{km}{N}$ | | $\mu \frac{(N-m)(N-k)}{N(N-1)}$ | | | | |
| HGeom(w, b, k) | | $\frac{\binom{w}{x} \binom{b}{k-x}}{\binom{w+b}{k}}$ | $\frac{k w}{w+b}$ | | $\mu \frac{b(w+b-k)}{(w+b)(w+b-1)}$ | | | | |
| Pois(μ) | $e^{-\mu} \sum_{i=0}^x \frac{\mu^i}{i!}$ | $\frac{\mu^x}{x!} e^{-\mu}, x \in 0, 1..$ | μ | $\mu^2 + \mu$ | μ | $e^{\mu(e^t-1)}$ | $\mu e^t M(t)$ | $\mu e^t (1 + \mu e^t) M(t)$ | |
| Unif(n) | | $\frac{1}{n}, x \in 1, 2..n$ | $\frac{n+1}{2}$ | $\frac{(n+1)(2n+1)}{6}$ | $\frac{(n^2-1)}{12}$ | $\frac{\sum_{i=1}^n e^{ti}}{n}$ | | | |
| Unif(a, b) | $\frac{x-a}{b-a}$ | $\frac{1}{b-a}, x \in (a, b)$ | $\frac{a+b}{2}$ | | $\frac{(b-a)^2}{12}$ | $\frac{e^{tb}-e^{ta}}{t(b-a)}$ | | | |
| $\mathcal{N}(\mu, \sigma^2)$ | | $\frac{1}{\sigma\sqrt{2\pi}} e^{-\frac{(x-\mu)^2}{2\sigma^2}}$ | μ | $\mu^2 + \sigma^2$ | σ^2 | $e^{\mu t + \frac{\sigma^2 t^2}{2}}$ | $(\mu + \sigma^2 t) M(t)$ | $[(\mu + \sigma^2 t)^2 + \sigma^2] M(t)$ | |
| $\mathcal{N}(0, 1)$ | | $\frac{1}{\sqrt{2\pi}} e^{-\frac{x^2}{2}}$ | 0 | 1 | 1 | $e^{\frac{t^2}{2}}$ | | | |
| $\mathcal{LN}(\mu, \sigma^2)$ | | $\frac{1}{x\sigma\sqrt{2\pi}} e^{-\frac{(\ln x - \mu)^2}{2\sigma^2}}$ | $e^{\mu + \frac{\sigma^2}{2}}$ | $e^{2\mu + 2\sigma^2}$ | $\theta^2(e^{\sigma^2} - 1)$ | \times | | | |
| Cauchy(θ, σ^2) | | $\frac{1}{\pi\sigma} \frac{1}{1+(\frac{x-\theta}{\sigma})^2}$ | \times | \times | \times | | | | |
| DExpo(μ, σ^2) | | $\frac{1}{2\pi\sigma} e^{- \frac{x-\mu}{\sigma} }$ | μ | $\mu^2 + 2\sigma^2$ | $2\sigma^2$ | $\frac{e^{\mu t}}{1-\sigma^2 t^2}$ | | | |
| Expo(λ) | $1 - e^{-\lambda x}$ | $\lambda e^{-\lambda x}, x \in (0, \infty)$ | $\frac{1}{\lambda}$ | | $\frac{1}{\lambda^2}$ | $\frac{\lambda}{\lambda-t}, t < \lambda$ | | | |
| Expo(β) | | $\frac{1}{\beta} e^{-\frac{x}{\beta}}$ | β | | β^2 | $\frac{1}{1-\beta t}$ | $\beta(1-\beta t)^{-2}$ | $2\beta^2(1-\beta t)^{-3}$ | |
| Gamma(a, λ) | | $\frac{\lambda^a}{\Gamma(a)} x^{a-1} e^{-\lambda x}, x \in (0, \infty)$ | $\frac{a}{\lambda}$ | $\frac{a}{\lambda^2}$ | $\left(\frac{\lambda}{\lambda-t}\right)^a, t < \lambda$ | | | | |
| Gamma(α, β) | | $\frac{1}{\Gamma(a)\beta^\alpha} x^{a-1} e^{-x/\beta}$ | $\alpha\beta$ | $\alpha\beta^2$ | $\left(\frac{1}{1-\beta t}\right)^a, t < \frac{1}{\beta}$ | | | | |
| Beta(a, b) | | $\frac{\Gamma(a+b)}{\Gamma(a)\Gamma(b)} x^{a-1} (1-x)^{b-1}$ | $\frac{a}{a+b}$ | | $\frac{\mu(1-\mu)}{(a+b+1)}$ | | | | $\frac{\Gamma(\alpha+n)\Gamma(\alpha+\beta)}{\Gamma(\alpha+\beta+n)\Gamma(\alpha)}$ |
| $B(\alpha, \beta) =$ | | $\frac{\Gamma(\alpha)\Gamma(\beta)}{\Gamma(\alpha+\beta)}, x \in (0, 1)$ | | $\frac{a(a+1)}{(a+b)(a+b+1)}$ | $\frac{ab}{(a+b)^2(a+b+1)}$ | | | | |
| χ_p^2 | | $\frac{1}{2^{p/2}\Gamma(p/2)} x^{p/2-1} e^{-x/2}$ | p | $2p + p^2$ | $2p$ | $(1-2t)^{-p/2}, t < \frac{1}{2}$ | | | |
| t_n | | $\frac{\Gamma(\frac{n+1}{2})}{\sqrt{n\pi}\Gamma(\frac{n}{2})} (1 + \frac{x^2}{n})^{-\frac{n+1}{2}}$ | $0, n > 1$ | | $\frac{n}{n-2}, n > 2$ | \times | | | |