What do I need to know after Urethroplasty?

DR JUSTIN CHEE
RECONSTRUCTIVE UROLOGIST

- What to expect?
- Wound care
- Catheter care
- Medications
- DO'S and DON'T'S
- Follow up plan

What to expect?

- Length of Stay in hospital: 3-Day 2-Night
- Dr Chee will visit you daily
- Catheter (IDC) stays in bladder until 4 weeks after surgery
- Day 1 he will remove outer supportive dressing
- Get out of bed/ walk around/ shower after Dr Chee reviewed
- You will be educated to use leg bag

Mouth Wound care

- If your operation required a Buccal Mucosa Graft (BGM)
- Avoid hard to chew food eg steak/ hamburgers
- Stitches are dissolvable, avoid touching or pulling
- If stitches come apart or break, don't be alarmed, can happen; try to limit degree of mouth opening
- May gargle with warm salt water twice a day

Perineal Wound care

- Dressing is waterproof
- Pat dry with clean towel after shower to avoid rubbing off
- Please remove dressing in the shower 1 day prior to your post- op appointment (7-10 days after)

Catheter (IDC) care

- IDC always taped onto abdomen with Mefix
- Reapply everyday after shower
- Leg bag: Attach to thigh, empty regularly, change weekly
- Night bag: Put on floor when sleeping
- Your Nurse will show you how to care IDC before discharge

Medications

- Regular pain relief will be given in hospital, can ask for more if needed
- Adequate pain relief will be given for discharge
- Common side effect: constipation
- Antibiotics for 5 days
- Important to take laxatives (coloxyl/ lactulose- which can be purchased over the counter), prunes or pears to keep stool soft

DO's

- Walk as much as you want/ feel comfortable with
- Take laxatives if needed
- Salt water to gargle if there is mouth wound
- Sit leaning backwards instead of sitting up to avoid pressure on wound
- Can sit on doughnut cushion if needed
- Do let Dr Chee knows if increased pain/ fever/chills/ feeling unwell/ increase bleeding
- Do ignore erections if there is any, can be quite painful if IDC is in place, can take a cool shower or walk around to reduce sensation
- Do let Dr Chee knows if you develop bladder spasm or urine leaking around IDC, we can organise some medication to assist

DON'T'S

- No lifting more then 10kg
- No strenuous activity for 4-6 weeks after surgery
- No driving until IDC removed and discontinued pain relief
- No intercourse for 6 weeks
- No bike riding for minimum 6 months- no direct pressure to wound

Follow up plan

- Will have appointment booked in rooms around 1 week post discharge for wound review
- 4 weeks after surgery for IDC removal
- 4- 6 weeks after removal of IDC for flow test and review

Material to take home with

- 4 x leg bags
- 4 x night bags
- 4- 6 leggy fix
- Multiple combines (to wear in your underwear when the dressing is removed in 1 week)
- Multiple stripes of Mefix, cut to length to make it easy to apply

Placement of IDC



Contacting us

Nurses Contact:
nurse@drjustinchee.com
Available Tue & Fri.

Reception:
recption@drjustinchee.com or 9088 5138
Tue - Fri 0900- 1630

Pager service- 03 8679 6579
Only for URGENT matters during
non- business hours