

LABORATORY REPORT

REMARKS

: URVISH SUKHADIA Sex/Age : Male J. 22 Years Casse ID: : 20200203135

Ref. By . Dr. Self Dis. At : LAB Pt ID -1877302

Pt. Loc : OPD Collection Bill. Loc. : Labcore spec lab baroda

: 12-Feb-2022 13:53 Sample Type Reg Date and Time : Nasopharyangeal + Mobile No.

RESULTS

Oropharyngeal Swab

BIOLOGICAL REF RANGE

Sample Date and Time : 12-Feb-2022 13:53 Sample Coll. By : non Refild1 TRAVEL Report Date and Time : 13-Feb-2022 20:46 Acc. Remarks Ref Id2 : 7984297731

UNIT

Genomics

COVID19 Qualitative by Real time PCR (ICMR No. SUPRASSIF)

COVID19 Interpretation NEGATIVE

N gene (Ct) Negative

Orf gene (Ct) Negative

Test: Qualitative test of COVID19 RNA by standard procedure on rt Real-time PCR.

Methodology: Reverse transcriptase Real-time Polymerase chain reaction.

Interpretations:

TEST

Real time PCI

Cycle threshold (Ct value) Value ranges from 15-40 cycle. Lower the Ct value higher is the viral load (Inversely proportional).

Kindly correlate with the clinical presentation and findings.

According to latest CDC guidelines, Ct cutoff of more than 33 is not considered as infective as it is extremely difficult to detect any live virus in a sample above the threshold of 33 cycles. Clinical Significance:

- Coronaviruses are a family of large RNA viruses with size ranging from 26 to 32 kb.
- As the coronavirus is anRNA virus it has a relatively high mutation rate resulting in rapid evolution.
- In December 2019, a new deadly coronavirus known as 2019-nCoV, which has a high sequence similarity to SARS-CoV, was identified and has caused a pneumonia outbreak in Wuhan, China and spread globally.

- The results of this test are highly dependent on the sampling technique employed, sample type, cold-chain maintenance andclinical condition. There is poor standardization between commercially available PCR tests, and results from different institutions should not be directly compared. Results are best monitored using a single institution.
- b. Presence of PCR inhibitors (cannot be traced by technologist), specimen collected very early/late in infection or viral load lesser than the assay lower limit of detection as wellas presence of rare genotypes or mutations may result in false-negative report.
- False-positive report may be obtained in cases where there is possibility of background RNA contamination from pre analyticalor in lab environment.
- d. The assay performance characteristics for this test are determined by STMPL which is used for clinical diagnosis. This test isnot approved by FDA nor accredited by NABL or CAP.
- RT-PCR kits used for this assay are approved by ICMR (Supratech Micropath Laboratory & Research Institute Pvt. Ltd. ICMR No. SUPRA001f). Test performed on Quantstudio 5 Real-time PCR machine.

--- End Of Report --

Note:(LL-VeryLow,L-Low,H-High,HH-VeryHigh ,A-Abnormal)

Dr. Krutarth Shah

M.D. Microbiologist

Dr. Sandip Shah M.D. (Path, & Bact.)

Consultant Pathologist

Page 1 of 1

Printed On: 12-Feb-20