

the 1990s, the number of people with a mental health problem has increased by 50% (Mental Health Foundation 2000). The prevalence of mental health problems in the UK is estimated to be 10% (Mental Health Foundation 2000).

There is a growing awareness of the need to address the needs of people with mental health problems. The Department of Health (2000) has set out a vision for mental health care in the UK, which is based on the principles of recovery, self-help, and community care. The vision is to ensure that people with mental health problems are able to live full and meaningful lives, and that they are able to participate in the community.

One of the key challenges in achieving this vision is to ensure that people with mental health problems are able to access the services they need. This is particularly true for people who are homeless, as they often face significant barriers to accessing mental health services. The Department of Health (2000) has identified homelessness as one of the key areas where mental health services need to be improved.

One of the ways in which mental health services can be improved for people who are homeless is by providing them with a safe and stable home. This is because homelessness is often a major factor in the development of mental health problems, and it can make it difficult for people to access mental health services. The Department of Health (2000) has identified housing as one of the key areas where mental health services need to be improved.

Another way in which mental health services can be improved for people who are homeless is by providing them with support and advice. This is because people who are homeless often face significant challenges, and they may need help to access mental health services. The Department of Health (2000) has identified support and advice as one of the key areas where mental health services need to be improved.

One of the ways in which support and advice can be provided to people who are homeless is by providing them with a dedicated mental health service. This is because people who are homeless often face significant barriers to accessing mental health services, and they may need a dedicated service to help them. The Department of Health (2000) has identified a dedicated mental health service as one of the key areas where mental health services need to be improved.

Another way in which support and advice can be provided to people who are homeless is by providing them with a dedicated housing service. This is because people who are homeless often face significant challenges, and they may need help to access housing services. The Department of Health (2000) has identified a dedicated housing service as one of the key areas where mental health services need to be improved.


One of the ways in which a dedicated housing service can be provided to people who are homeless is by providing them with a dedicated housing officer. This is because people who are homeless often face significant challenges, and they may need help to access housing services. The Department of Health (2000) has identified a dedicated housing officer as one of the key areas where mental health services need to be improved.

























S1


S

2

Quiz










































2

3











the 1990s, the number of people in the world who are undernourished has increased from 600 million to 800 million. The number of people who are malnourished has increased from 1.2 billion to 1.5 billion. The number of people who are obese has increased from 100 million to 300 million.

The World Bank's *World Development Indicators* (2004) report that the number of people who are undernourished has increased from 600 million in 1990 to 800 million in 2000. The number of people who are malnourished has increased from 1.2 billion in 1990 to 1.5 billion in 2000. The number of people who are obese has increased from 100 million in 1990 to 300 million in 2000.

The World Bank's *World Development Indicators* (2004) report that the number of people who are undernourished has increased from 600 million in 1990 to 800 million in 2000. The number of people who are malnourished has increased from 1.2 billion in 1990 to 1.5 billion in 2000. The number of people who are obese has increased from 100 million in 1990 to 300 million in 2000.

The World Bank's *World Development Indicators* (2004) report that the number of people who are undernourished has increased from 600 million in 1990 to 800 million in 2000. The number of people who are malnourished has increased from 1.2 billion in 1990 to 1.5 billion in 2000. The number of people who are obese has increased from 100 million in 1990 to 300 million in 2000.

The World Bank's *World Development Indicators* (2004) report that the number of people who are undernourished has increased from 600 million in 1990 to 800 million in 2000. The number of people who are malnourished has increased from 1.2 billion in 1990 to 1.5 billion in 2000. The number of people who are obese has increased from 100 million in 1990 to 300 million in 2000.

The World Bank's *World Development Indicators* (2004) report that the number of people who are undernourished has increased from 600 million in 1990 to 800 million in 2000. The number of people who are malnourished has increased from 1.2 billion in 1990 to 1.5 billion in 2000. The number of people who are obese has increased from 100 million in 1990 to 300 million in 2000.

The World Bank's *World Development Indicators* (2004) report that the number of people who are undernourished has increased from 600 million in 1990 to 800 million in 2000. The number of people who are malnourished has increased from 1.2 billion in 1990 to 1.5 billion in 2000. The number of people who are obese has increased from 100 million in 1990 to 300 million in 2000.


The World Bank's *World Development Indicators* (2004) report that the number of people who are undernourished has increased from 600 million in 1990 to 800 million in 2000. The number of people who are malnourished has increased from 1.2 billion in 1990 to 1.5 billion in 2000. The number of people who are obese has increased from 100 million in 1990 to 300 million in 2000.






Quiz

S1



S2




c 1



c 2



H 3



ESEMPIO: HttpClient, Angular

http.get(...).subscribe();

- Sappiamo che esegue una chiamata Http
- Sappiamo che la chiamata viene effettuata solo al subscribe
- Sappiamo che emette al massimo un valore, e poi completa

Quindi, presumiamo che:

- Sia un observable cold
- Due subscribe allo stesso observable generino **due** chiamate distinte
- Dobbiamo evitare di usare una async pipe direttamente sull'observable, perché ogni ciclo di Change Detection farebbe ripartire la chiamata
- Se vogliamo condividere il risultato, dobbiamo usare 1 subscribe e salvare il risultato, oppure dobbiamo renderlo tiepido.