Quick Demo Chart for Community Health Workers

Publisher: Demo Community Health Programme
Edition: 1.0, November 2026
For software demonstration only. Not for clinical use.

WHO THIS CHART IS FOR

Use this chart for a child age 2 months up to 5 years.
If the child is younger than 2 months, or 5 years or older, do not use this chart.
Refer the child to the health facility with a referral note.
Follow up on return to the health worker.

ASK AND LOOK: do every step for every child

1. Ask: Does the child have a cough?
2. Ask: Has the child had convulsions?
3. Ask: Is the child not able to drink or breastfeed?
4. Look: Does the child have chest indrawing?
5. Count the breaths in one minute.
6. Ask the child's age in months (this chart is for age 2 months up to 5 years).

Ask every question above for every child, in this order, whatever the earlier answers.

DANGER SIGNS AND FAST BREATHING

Danger signs: convulsions, not able to drink or breastfeed, chest indrawing.
Fast breathing:
- Age 2 months up to 12 months: 50 breaths per minute or more.
- Age 12 months up to 5 years: 40 breaths per minute or more.

Go to page 2 to decide and treat.

DECIDE AND TREAT

Check the boxes in this order. Use the first box that applies to the child.

1. ANY DANGER SIGN

If the child has any danger sign:
Refer the child to the health facility immediately.
Follow up on return from the health facility.

2. FAST BREATHING, NO DANGER SIGN

If the child has fast breathing and no danger sign:
Treat at home: give oral amoxicillin 250 mg tablet twice daily for 5 days, the dose by age being
Age 2 months up to 12 months – 1 tablet; Age 12 months up to 5 years – 2 tablets.
Follow up in 3 days.

3. COUGH, NO FAST BREATHING, NO DANGER SIGN

If the child has a cough, no fast breathing and no danger sign:
Give home care advice: keep giving fluids and food, soothe the throat with warm drinks.
Advise the caregiver to come back if the child gets worse.

4. NO COUGH, NO FAST BREATHING, NO DANGER SIGN

If none of the boxes above applies:
No treatment is needed.
Advise the caregiver to come back if the child gets sick.