

Representations of Lie Groups

Main goal: discuss representations of compact/complex semisimple Lie groups. (At the end we will also briefly discuss infinite-dimensional representations of noncompact groups.)

Note: These groups and their representations (in finite-dimensional case) were completely classified in the early 20th century by work of Killing, E. Cartan and Weyl.

References:

- Adams: "Lectures on Lie Groups"
- Fulton & Harris: "Representation Theory"

§1: Introduction

Mantra: examples are the key to this subject.

E.g.: S^1 , $O(3)$, $SO(3)$. More generally,

SKI) Pers. LI (2)

any closed subgroup of $GL(n, \mathbb{R})$.

The Lie algebra of a Lie group G can be defined in various equivalent ways. We take it to be the tangent space at the identity:

$$\mathfrak{g} = T_1 G$$

And define its Lie bracket via the following construction.

Recall the adjoint action $G \curvearrowright G$ (given by conjugation). Then for $g \in G$ we have a map $\text{Ad}_g: G \rightarrow G$.

Since $\text{Ad}_g(1) = 1$, we then obtain $d_1 \text{Ad}_g: \mathfrak{g} \rightarrow \mathfrak{g}$, and in fact $d_1 \text{Ad}_g \in GL(\mathfrak{g})$. Thus we get a map:

$$G \rightarrow GL(\mathfrak{g})$$

which is by defⁿ the adjoint action of

SKD Reps. L1 (3)

G on its Lie algebra. Again this maps $I \in G$ to $\text{Id} \in \text{GL}(\mathfrak{g})$, so differentiating at I again gives a map:

$$\mathfrak{g} \rightarrow \text{Lie } \text{GL}(\mathfrak{g}) = \text{End } \mathfrak{g} = \mathfrak{g} \otimes \mathfrak{g}^*$$

This is equivalent to a map $\mathfrak{g} \otimes \mathfrak{g} \rightarrow \mathfrak{g}$, which is by defⁿ the Lie bracket of \mathfrak{g} .

one can check this gives \mathfrak{g} the structure of a Lie algebra.

Correspondence between Lie groups and Lie algebras

This construction has a (partial) inverse, giving an equivalence of categories:

Skd Reqs. L1 (4)

(1-connected Lie groups) \longleftrightarrow (finite-dimensional Lie algebras)

(where 1-connected means connected and simply-connected.)

Note: The universal cover of any connected Lie group has a natural Lie group structure. Thus there is a natural 1-connected Lie group associated to any Lie group G : simply take the universal cover of the identity component of G . Since these are locally isomorphic near I , their Lie algebras will be the same.

To get an idea of how one might construct a Lie group from a Lie algebra, consider a Lie group G .

Given a path $g(t)$ in G , we get two elements of our Lie algebra for each t :

SKI) REPS. LI (5)

$$\left. \begin{aligned} (dLg(t))^{-1}(\dot{g}(t)) &\in T_1G = \mathcal{H} \\ (dRg(t))^{-1}(\dot{g}(t)) &\in T_1G = \mathcal{H} \end{aligned} \right\}$$

After making a choice of left or right, we then see that a path $g(t)$ in G is completely determined by $g(0) \in G$ and $\xi(t)$ the path $\xi(t)$ in \mathcal{H} (given by one of the formulae above): we extend $\xi(t)$ to a time-dependent vector field (left- or right-invariant) and take the flow line passing through $g(0)$ at time 0.

We now consider paths with $g(0) = 1$. Since G is simply connected we have:

elements of $G \longleftrightarrow$ homotopy classes of paths in G with $g(0) = 1$.
 \longleftrightarrow equivalence classes of paths in \mathcal{H} .

SKD Reps. L1 ⑥

And we can describe this equivalence relation explicitly.

This gives us an invariant way to construct G from a Lie algebra.

Equations of motion for rigid bodies

Suppose we have a time-dependent family of rotations, $g_t \in \text{SO}(3)$.

Then we can interpret:

$$g^{-1}\dot{g} := (dL_g)^{-1}\dot{g} \iff \begin{array}{l} \text{angular velocity} \\ \text{in frame} \end{array}$$

$$\dot{g}g^{-1} := (dR_g)^{-1}\dot{g} \iff \begin{array}{l} \text{angular velocity} \\ \text{in space.} \end{array}$$


So the physics leads us to a study of $\text{Lie SO}(3) \cong (\mathbb{R}^3, \times)$.

SKD REPS. L1 (7)

More examples (classical groups)

We organise the classical groups as follows.

Compact:



Complex:



Most of the groups we will be interested in fall into this picture.

§2: Study of $SU(2)$

This is a very important example:

$$\begin{aligned} SU(2) &= \{A \in \mathbb{C}^{2 \times 2} : A^* A = I, \det A = 1\} \\ &= \left\{ \begin{pmatrix} z & w \\ -\bar{w} & \bar{z} \end{pmatrix} : |z|^2 + |w|^2 = 1 \right\} \cong S^3 \end{aligned}$$

(Aside: S^1 and S^3 are the only spheres which admit a Lie group structure.)

Certainly $SU(2)$ is 1-connected. The adjoint action takes the form:


$$SU(2) \longrightarrow GL(3, \mathbb{R})$$

And in fact it takes values in $SO(3)$.

This map is actually a double cover, and since $\pi_1 SO(3) = \mathbb{Z}_2$ we have:

$$SO(3) = SU(2) / \boxed{\mathbb{Z}_2} = SU(2) / \{\pm I\}$$

So that $SU(2)$ is the universal cover of $SO(3)$. Thus $\text{Lie } SU(2) = \text{Lie } SO(3) = (\mathbb{R}^3, \times)$.

SKD Reps. LI  (9)

And $\text{Lie } \text{su}(2) = \text{Lie } \text{so}(3) = \mathbb{R}^3$
equipped w/ cross product.
($\text{so } \text{su}(2)$ is universal cover of $\text{so}(3)$)

Now we also have $\text{su}(2) \subseteq \text{SL}(2, \mathbb{C})$.
Furthermore:

$$\begin{aligned}\text{Lie } \text{SL}(2, \mathbb{C}) &= \text{Lie } \text{su}(2) \otimes_{\mathbb{R}} \mathbb{C} \\ &= \text{Lie } \text{su}(2) \oplus i \text{Lie } \text{su}(2)\end{aligned}$$

we can thus consider:

$\left. \begin{array}{l} \text{Reps. of } \text{su}(2) \\ \text{Reps. of } \text{SL}(2, \mathbb{C}) \\ \text{Reps. of } \text{Lie } \text{su}(2) \\ \text{Reps. of } \text{Lie } \text{SL}(2, \mathbb{C}) \end{array} \right\}$

(All representations are complex in this course.) The point is that these four are equivalent (mostly follows from above discussion).

SKD Reps. LI (10)

Algebraic approach

We look at reps. of $\text{Lie } \mathfrak{sl}(2, \mathbb{C})$. We use the following basis:

$$H = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}, \quad X = \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix}, \quad Y = \begin{pmatrix} 0 & 0 \\ 1 & 0 \end{pmatrix}.$$

The structure constants are then:

$$[H, X] = 2X, \quad [H, Y] = -2Y, \quad [X, Y] = H$$

Let V be a f.d.V.S. acted on by $\mathfrak{sl}(2, \mathbb{C})$, irreducible as a module.

Consider $\langle iH \rangle = \text{Lie } \mathfrak{su}(2)$. This is the Lie algebra of:

$$S^1 = \left\{ \begin{pmatrix} e^{i\theta} & 0 \\ 0 & e^{-i\theta} \end{pmatrix} : \theta \in \mathbb{R} \right\} \subseteq \text{SU}(2).$$

Inters. of S^1 on $\mathbb{C}V$ given by $\lambda \mapsto \lambda^?$

So restricting action on V to $\langle iH \rangle$, can decompose V into a direct sum of $\mathbb{C}V$ reps.

Skid Reps. L1 ⑪

$V = \bigoplus_{\alpha \in S \subseteq \mathbb{Z}} V_{\alpha} \leftarrow$ as a vector space, not an $SL(2, \mathbb{C})$ -module

where $\begin{pmatrix} e^{i\alpha} & 0 \\ 0 & e^{-i\alpha} \end{pmatrix}$ acts on V_{α} as $e^{i\alpha}$, so H acts on V_{α} as α .

Let $e_{\alpha} \in V_{\alpha}$. Then $He_{\alpha} = \alpha e_{\alpha}$, and:

$$\begin{aligned} HXe_{\alpha} &= XHe_{\alpha} + 2Xe_{\alpha} \\ &= (\alpha + 2)Xe_{\alpha} \end{aligned}$$

$$\Rightarrow Xe_{\alpha} \in V_{\alpha+2}$$

By a similar argument:

$$Ye_{\alpha} \in V_{\alpha-2}$$

Note that if $g = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} \in SU(2)$ then:

$$g \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix} g^{-1} = \begin{pmatrix} -1 & 0 \\ 0 & 1 \end{pmatrix}$$

$$\Rightarrow gHg^{-1}(e_{\alpha}) = -\alpha e_{\alpha}$$

$$\Rightarrow Hg^{-1}(\boxed{e_{\alpha}}) = -\alpha g^{-1}e_{\alpha}$$

So if α is a weight, so is $-\alpha$,
(with $\boxed{\alpha} V_{-\alpha} = \langle g^{-1}\boxed{e_{\alpha}} \rangle$).

SKD Reps. LI (12)

Choose $\alpha \in S$ maximal and choose $e \in V_{\alpha_{\max}}$ (the highest weight vector).

Claim: If V is irreducible then it is generated by $\{Y^p e\}$.

Proof: we must prove $X^p e$ belongs to this span. (From this it follows that the submodule generated by e is spanned by $\{Y^p e\}$, hence is equal to V since V irreducible).

But we know how X and Y change weights. So need $X^p e$ a multiple of $Y^p e$, which we can prove easily by induction. \square

~~The claim is that~~ we conclude that V is completely determined by $\alpha_{\max} \in \mathbb{N}_{>0}$.

SKID Reps. LI (13)


Thus the irreps. of $\begin{pmatrix} \mathrm{SU}(2) \\ \mathrm{SL}(2, \mathbb{C}) \end{pmatrix}$ are the $S^d U$ (where $U \equiv \mathbb{C}^2$ w/ tautological rep.) (cf. polys. homogeneous).

Geometric approach



If $\mathrm{SL}(2, \mathbb{C}) \curvearrowright V$ then it also acts on $\mathbb{P}(V^*)$. Suppose we know that \exists an orbit isomorphic to \mathbb{CP}^1 , not contained in any proper linear subspace. Call it C . Now $\mathrm{SL}(2, \mathbb{C})$ has a standard action on $\mathbb{CP}^1 = S^2$ by Möbius transformations.

Here, $V = H^0(C, L)$ where $L \rightarrow C$ is a line bundle (see later). So the classification is that of line bundles over \mathbb{CP}^1 with a lift of the action $\mathrm{SL}(2, \mathbb{C}) \curvearrowright \mathbb{CP}^1$.

This is well-known from Cpx geometry: $L = \mathcal{O}(d)$. Then:

SKD Refs. LI  (14)

$$V = H^0(C, \mathcal{O}(d)) = S^d U \leftarrow \begin{array}{l} \text{homogeneous} \\ \text{polys.} \\ \text{of degree } d. \end{array}$$

Note: $SO(3) = SU(2)/\langle \pm 1 \rangle$. So every
rep. of $SO(3)$ lifts to a rep. of $SU(2)$.
Conversely a rep. of $SU(2)$ factors
through to one of $SO(3)$ iff -1
acts trivially. For  reps $S^d U$
this happens iff d  even.

Alternatively look at harmonic polys:
those for which $\Delta f = 0$. Then:

$$H_1 = \{\text{linear polys.}\}$$

$$H_2 = \left\{ \begin{array}{c} \text{trace-free} \\ 2 \times 2 \end{array} \text{ symmetric matrices} \right\}$$

$$H_k = S^{2k} U$$