AI Med Rec Training Dataset — Emergency Department

Medication Reconciliation Scenarios with Ranked Drug Interactions

20 Patient Scenarios | All Drug Classes | Geriatric-Inclusive

---

Dataset Purpose: Training data for AI-assisted medication reconciliation in the Emergency Department. Each scenario includes: patient demographics, multi-source medication data (verbal, written, pill bottle), presenting symptoms, and a ranked list of the top 3 clinically significant interactions (drug–drug, drug–symptom, drug–disease, drug–lab, OTC/supplement).

SCENARIO 1

Patient Demographics

- **Name:** Mary Smith
- **Age:** 62
- **Sex:** Female
- **Setting:** Emergency Department

Chief Complaint

"I've had this dry, tickling cough for about three months. My throat feels fine and I don't have a fever. I thought it was allergies but the allergy medicine didn't help."

Medication Sources

Source 1 — Handwritten List (Patient-Provided)

```
Mary Smith - Medications
- Lisinopril 10 mg once daily
- Atorvastatin 20 mg at night
- Calcium (no dose noted)
```

Source 2 — Verbal Report

"I also take metformin, one pill in the morning and one at night. I think it's 500 milligrams each."

Source 3 — Pill Bottle (Physical)

Patient places bottle on counter: "This one is my blood pressure medicine — I take one every morning."

- **Label reads:** Lisinopril 10 mg tablets — Take 1 tablet by mouth once daily

Reconciled Medication List

Medication	Dose	Frequency	Source
Lisinopril	10 mg	Once daily (AM)	Written list + pill bottle (confirmed)
Atorvastatin	20 mg	Once daily (PM)	Written list
Metformin	500 mg	Twice daily (AM/PM)	Verbal only
Calcium	Unknown dose	Unknown frequency	Written list (incomplete)

Reconciliation Flags

- Calcium: dose and frequency not documented — clarify supplement type (carbonate vs. citrate) and dose
- Metformin: verbal only, no written or bottle confirmation — verify with pharmacy or PCP

Top 3 Ranked Interactions

🥇 Rank 1 — Drug–Symptom Interaction [HIGH CLINICAL RELEVANCE]

Lisinopril → ACE Inhibitor–Induced Cough

- **Mechanism:** ACE inhibitors block degradation of bradykinin and substance P in the lungs, causing a persistent dry cough in 5–20% of patients (higher incidence in women and patients of Asian descent).
- **Clinical correlation:** Patient's cough is dry, chronic (3 months), non-productive, and did not respond to antihistamines — classic presentation. No infectious or allergic etiology identified.
- **Action:** Confirm onset relative to lisinopril initiation. Consider switching to an ARB (e.g., losartan) if ACE inhibitor is continued for the same indication.
- **Confidence:** HIGH — temporal pattern + classic symptom profile.

🥈 Rank 2 — Drug–Lab Interaction [MODERATE RELEVANCE]

Metformin → Renal Function (Creatinine / eGFR)

- **Mechanism:** Metformin is renally cleared. In renal impairment, metformin accumulates and increases risk of lactic acidosis. FDA contraindicates use if eGFR < 30 mL/min/1.73m².
- **Clinical correlation:** Patient is 62F on metformin with no documented renal function labs in this visit. Age-related GFR decline is common.
- **Action:** Order BMP/CMP; if eGFR 30–45, reduce dose and reassess; if < 30, hold metformin.
- **Confidence:** MODERATE — requires lab confirmation before clinical decision.

🥉 Rank 3 — Drug–Drug / OTC Interaction [LOW-MODERATE RELEVANCE]

Calcium Supplement → Atorvastatin Absorption

- **Mechanism:** Calcium carbonate (the most common OTC form) can reduce absorption of some statins when taken simultaneously. Evidence is modest but clinically acknowledged.
- **Clinical correlation:** Dose and timing of calcium are unknown. If taken at the same time as atorvastatin, absorption may be suboptimal.
- **Action:** Clarify calcium type and timing. Advise 2-hour separation if using carbonate form.
- **Confidence:** LOW-MODERATE — interaction magnitude depends on formulation and timing.

SCENARIO 2

Patient Demographics

- **Name:** John Rivera
- **Age:** 71
- **Sex:** Male
- **Setting:** Emergency Department

Chief Complaint

"I've been feeling lightheaded when I stand up. It started a few days ago. My legs are also a little swollen."

Medication Sources

Source 1 — Printed Clinic Summary (After-Visit Summary from PCP)

```
John Rivera – Active Medications
- Amlodipine 5 mg once daily
- Metoprolol 25 mg twice daily
- Aspirin 81 mg daily
- Omeprazole 20 mg daily
```

Source 2 — Verbal Report

"I think my doctor also told me to take a water pill but I don't see it on this list. I didn't bring it."

Reconciled Medication List

Medication	Dose	Frequency	Source
Amlodipine	5 mg	Once daily	Printed list (confirmed)
Metoprolol	25 mg	Twice daily	Printed list (confirmed)
Aspirin	81 mg	Once daily	Printed list (confirmed)
Omeprazole	20 mg	Once daily	Printed list (confirmed)
Unknown diuretic	Unknown	Unknown	Verbal only — unverified

Reconciliation Flags

- Possible unverified diuretic — contact PCP or pharmacy to confirm
- Leg edema in a patient on amlodipine warrants differentiation: drug-induced vs. new cardiac/venous pathology

---

Top 3 Ranked Interactions

🥇 Rank 1 — Drug–Symptom Interaction [HIGH CLINICAL RELEVANCE]

Amlodipine → Peripheral Edema (Dependent)

- **Mechanism:** Dihydropyridine calcium channel blockers (e.g., amlodipine) cause preferential arterial vasodilation without equivalent venous dilation, leading to capillary fluid leak and dependent edema in up to 30% of patients, more common in older adults.
- **Clinical correlation:** Patient presents with bilateral leg swelling. Orthostatic hypotension (lightheadedness on standing) may also be aggravated by vasodilatory effect. Edema is drug-induced and not necessarily indicative of heart failure.

- **Action:** Check for pitting vs. non-pitting edema; evaluate for CHF concurrently. Consider dose reduction or switching to amlodipine + ACE inhibitor combination (reduces edema via venodilation). Do not start diuretic without confirming diagnosis.
- **Confidence:** HIGH.

🥈 Rank 2 — Drug–Drug Interaction [MODERATE RELEVANCE]

Metoprolol + Amlodipine → Additive Hypotension / Bradycardia

- **Mechanism:** Beta-blockers reduce cardiac output; calcium channel blockers reduce systemic vascular resistance. Combination can produce additive hypotension and, when non-DHP CCBs are used (less relevant here with amlodipine), significant bradycardia. Orthostatic hypotension is amplified in elderly patients.
- **Clinical correlation:** Patient is 71 years old with orthostatic lightheadedness on a dual antihypertensive regimen — this combination is the most likely pharmacologic contributor.
- **Action:** Orthostatic vitals (lying/sitting/standing BP and HR). If symptomatic orthostasis confirmed, consider reducing amlodipine dose.
- **Confidence:** HIGH for symptom causation; moderate for dose adjustment decision.

🥉 Rank 3 — Drug–Drug Interaction [LOW-MODERATE RELEVANCE]

Aspirin + Omeprazole (Protective — Not Harmful)

- **Mechanism:** Aspirin (even low-dose 81 mg) causes GI mucosal irritation. Omeprazole provides gastroprotection. This is an intentional combination — likely prescribed together.
- **Clinical correlation:** Combination is appropriate. Flag: omeprazole can reduce clopidogrel efficacy (if patient is ever started on clopidogrel), but currently no issue.
- **Action:** Confirm indication for aspirin (primary vs. secondary prevention). If primary prevention only in a 71-year-old, current guidelines recommend reassessing risk-benefit.
- **Confidence:** Protective interaction — no harm expected. Reassess aspirin indication per age.

SCENARIO 3

Patient Demographics

- **Name:** Eleanor Johnson
- **Age:** 78
- **Sex:** Female

- **Setting:** Emergency Department

Chief Complaint

"I've been more tired than usual and had some mild bruising on my arms that I don't remember causing. I also had a small cut last week that took a long time to stop bleeding."

Medication Sources

Source 1 — Typed List (From Home)

```
Eleanor Johnson – Medications
- Losartan 50 mg once daily
- Furosemide 20 mg once daily
- Apixaban (Eliquis) – blood thinner [DOSE AND FREQUENCY MISSING]
- Vitamin B12 supplement
```

Source 2 — Verbal Report

"I take Eliquis, one pill in the morning and one at night. I think it's a small dose, not the strongest one. My cardiologist adjusted it because of my kidneys."

Source 3 — Pill Bottle (Physical)

Patient presents a bottle: "This is the one I really worry about getting wrong."

- **Label reads:** Apixaban (Eliquis) 2.5 mg — Take 1 tablet by mouth twice daily

Source 4 — Additional Pill Bottle

Patient produces a second bottle: "This is the heart medicine I take now."

- **Label reads:** Carvedilol 12.5 mg — Take 1 tablet by mouth twice daily

[Note: Carvedilol NOT on patient's written list — newly identified medication]

Reconciled Medication List

Medication	Dose	Frequency	Source
Losartan	50 mg	Once daily	Written list (confirmed)
Furosemide	20 mg	Once daily	Written list (confirmed)
Apixaban (Eliquis)	2.5 mg	Twice daily	Written list (incomplete) + verbal + pill bottle (confirmed)
Vitamin B12	Unknown dose	Unknown frequency	Written list only

Medication	Dose	Frequency	Source
Carvedilol	12.5 mg	Twice daily	Pill bottle ONLY — MISSING from written list ⚠️

Reconciliation Flags

- **CRITICAL:** Carvedilol identified via pill bottle — completely absent from patient's written list. High-risk discrepancy.
- Apixaban dose confirmed as 2.5 mg BID — appropriate reduced dose per renal/age criteria (Eliquis dose reduction criteria: ≥2 of: age ≥80, weight ≤60 kg, Cr ≥1.5 mg/dL)
- Vitamin B12 dose and frequency unknown

Top 3 Ranked Interactions

🥇 Rank 1 — Drug–Disease + Drug–Symptom Interaction [CRITICAL]

Apixaban (Anticoagulant) → Bleeding Symptoms (Bruising, Prolonged Wound Bleeding)

- **Mechanism:** Apixaban inhibits Factor Xa, reducing thrombin generation. At therapeutic doses in elderly patients (especially those with reduced renal clearance), excessive anticoagulation can manifest as spontaneous bruising and prolonged bleeding.
- **Clinical correlation:** Patient is 78F with easy bruising, prolonged bleeding from minor wound — these are classic signs of over-anticoagulation or drug accumulation. Risk is amplified by age-related renal decline affecting apixaban clearance.
- **Action:** Assess renal function (BMP/Cr/eGFR). Verify dose reduction criteria met (2.5 mg BID is appropriate if ≥2 criteria met). Order CBC (platelet count), consider anti-Xa level if over-anticoagulation suspected. Fall risk assessment given age.
- **Confidence:** HIGH — symptoms directly correlate with anticoagulation effect.

🥈 Rank 2 — Drug–Drug Interaction [HIGH RELEVANCE]

Furosemide + Losartan → Hypotension / Electrolyte Imbalance

- **Mechanism:** Loop diuretics (furosemide) cause volume depletion and electrolyte wasting (Na^+, K^+). ARBs (losartan) block angiotensin II-mediated aldosterone release, impairing potassium excretion and sodium retention. In combination, these effects can cause hypotension and hyperkalemia, especially with renal impairment.
- **Clinical correlation:** 78-year-old patient on dual RAAS-modifying therapy with a diuretic — volume depletion, fatigue, and electrolyte disturbance are plausible contributors to her presentation.
- **Action:** BMP (Na, K, BUN, Cr). Orthostatic vitals. Fatigue may represent hypovolemia or electrolyte disturbance rather than cardiac or hematologic cause.
- **Confidence:** HIGH.

🥉 Rank 3 — Drug–Drug Interaction [MODERATE RELEVANCE]

Carvedilol (Undocumented) + Furosemide → Masked Hypoglycemia / Bradycardia Risk

- **Mechanism:** Carvedilol (non-selective beta-blocker with alpha-blocking activity) can cause bradycardia and mask hypoglycemia symptoms (tachycardia, tremor). Combined with furosemide-induced volume depletion, orthostatic hypotension risk is amplified.
- **Clinical correlation:** Carvedilol was NOT on patient's medication list — this is a reconciliation failure. Its absence could lead to inadvertent repeat dosing or discontinuation. In a 78-year-old on furosemide and an anticoagulant, the hemodynamic effects matter.
- **Action:** Confirm indication for carvedilol with cardiologist. Add to medication list. Evaluate HR and BP for carvedilol effect. Consider whether furosemide + carvedilol combination is intentional for heart failure management.
- **Confidence:** MODERATE — drug interaction risk is real; clinical significance depends on indication and current hemodynamics.

SCENARIO 4

Patient Demographics

- **Name:** Patricia "Trish" Lee
- **Age:** 66
- **Sex:** Female
- **Setting:** Emergency Department

Chief Complaint

"My heart has been racing on and off for the past week. I've also had some muscle cramps in my legs, especially at night."

Medication Sources

Source 1 — Verbal Report Only

"I take a thyroid pill every morning… I think it's levothyroxine 75. I take a cholesterol pill at night… I think it's simvastatin, maybe 20? I take a little water pill in the morning — hydrochlorothiazide — I think 25. And I take Tylenol when my knee hurts."

Source 2 — Loose Pills in Zip-Lock Bag

"These are the ones I take every day — sorry, I don't have the bottles."

- **Pill 1:** White, round, scored tablet — visually consistent with hydrochlorothiazide 25 mg
- **Pill 2:** Peach/tan oval tablet — visually consistent with simvastatin 20 mg

Reconciled Medication List

Medication	Dose	Frequency	Source
Levothyroxine	75 mcg (unconfirmed)	Once daily (AM)	Verbal only
Simvastatin	20 mg (unconfirmed)	Once daily (PM)	Verbal + visual pill ID
Hydrochlorothiazide	25 mg (unconfirmed)	Once daily (AM)	Verbal + visual pill ID
Acetaminophen (Tylenol)	Unknown dose	PRN (knee pain)	Verbal only

Reconciliation Flags

- All doses unconfirmed — contact pharmacy for verification
- Levothyroxine: dose critical to confirm; narrow therapeutic index drug
- Acetaminophen: frequency unknown — assess for potential overuse given OTC access

---

Top 3 Ranked Interactions

🥇 Rank 1 — Drug–Symptom Interaction [HIGH CLINICAL RELEVANCE]

Levothyroxine (Potentially Supratherapeutic) → Palpitations / Tachycardia

- **Mechanism:** Levothyroxine (T4) is a narrow therapeutic index drug. Even modest over-replacement elevates free T4 and T3 levels, increasing cardiac chronotropy and causing palpitations, tachycardia, and atrial fibrillation risk, particularly in women over 60.
- **Clinical correlation:** Patient presents with episodic racing heart for one week. If levothyroxine dose is incorrect (higher than needed) or has been recently changed, thyrotoxicosis may be driving her symptoms.
- **Action:** Order TSH (with free T4 if TSH is suppressed). EKG to characterize tachycardia. Verify exact levothyroxine dose with pharmacy or PCP.
- **Confidence:** HIGH — dose uncertainty in a narrow-therapeutic-index drug + cardiac symptoms = priority interaction.

🥈 Rank 2 — Drug–Symptom + Drug–Lab Interaction [HIGH RELEVANCE]

Hydrochlorothiazide → Hypokalemia → Muscle Cramps + Cardiac Arrhythmia Risk

- **Mechanism:** Thiazide diuretics inhibit sodium-chloride cotransporter in the distal tubule, increasing urinary potassium excretion. Resulting hypokalemia causes muscle cramping, weakness, and in severe cases, cardiac arrhythmias (QT prolongation, ventricular ectopy).
- **Clinical correlation:** Patient reports leg cramps at night (hallmark hypokalemia symptom) and episodic palpitations. These two symptoms together on HCTZ are consistent with a hypokalemia presentation.
- **Action:** BMP — check K^+, Mg^{2+} (magnesium depletion amplifies hypokalemia effects). EKG for QT assessment. Potassium repletion if $K^+ < 3.5$ mEq/L.
- **Confidence:** HIGH.

🥉 Rank 3 — Drug–Drug Interaction [MODERATE RELEVANCE]

Simvastatin + Levothyroxine (Hypothyroidism Effect) → Statin Myopathy Risk

- **Mechanism:** Hypothyroidism itself increases statin-induced myopathy risk by reducing CYP3A4 activity (relevant for simvastatin) and impairing mitochondrial muscle metabolism. If patient is currently undertreated (hypothyroid), simvastatin myopathy risk is elevated.
- **Clinical correlation:** Leg cramps may not be exclusively from hypokalemia — statin myopathy (especially with untreated or undertreated hypothyroidism) should be considered.
- **Action:** TSH, CK level. If TSH is elevated (undertreated hypothyroidism), myopathy risk from simvastatin is meaningfully increased. Consider switching to a hydrophilic statin (pravastatin, rosuvastatin) which carries lower myopathy risk.
- **Confidence:** MODERATE — requires TSH confirmation to stratify risk.

SCENARIO 5

Patient Demographics

- **Name:** Robert Kim
- **Age:** 69
- **Sex:** Male
- **Setting:** Emergency Department

Chief Complaint

"I've been feeling shaky and sweaty since this morning. I checked my blood sugar and it was 54. I had my usual insulin last night."

Medication Sources

Source 1 — Verbal Report

"I take Lantus every night — 20 units. And I take Humalog before meals, sliding scale."

Source 2 — Insulin Pens (Physical)

Patient shows two insulin pens:

- **Pen 1:** Lantus SoloStar (insulin glargine) — long-acting basal insulin
- **Pen 2:** Humalog KwikPen (insulin lispro) — rapid-acting insulin

Source 3 — No Written Sliding Scale Available

"I didn't bring the paperwork with the scale."

Reconciled Medication List

Medication	Dose	Frequency	Source
Insulin glargine (Lantus)	20 units	Once daily (PM)	Verbal + pen confirmed
Insulin lispro (Humalog)	Per sliding scale (unknown)	Before meals	Verbal + pen confirmed (scale not available)

Reconciliation Flags

- Sliding scale not available — critical gap; must obtain from PCP, endocrinologist, or pharmacy
- Hypoglycemia etiology: insufficient carbohydrate intake, excessive Lantus dose, or inappropriate sliding scale dose?
- Assess for renal impairment — insulin clearance decreases with CKD, increasing hypoglycemia risk

---

Top 3 Ranked Interactions

🥇 Rank 1 — Drug–Disease Interaction [CRITICAL]

Insulin (Basal + Bolus Regimen) → Hypoglycemia in Aging Kidney (Renal Impairment)

- **Mechanism:** As eGFR declines with age, insulin clearance is reduced and hypoglycemia risk increases significantly. Insulin requirements often decrease as kidney function worsens. In a 69-year-old with possible undetected CKD, a fixed 20-unit Lantus dose may be inappropriately high.

- **Clinical correlation:** Patient presents with blood glucose of 54 after routine Lantus dose — no reported changes to diet, exercise, or dose. Renal insulin clearance impairment is a key underrecognized cause of recurrent hypoglycemia in older diabetic patients.
- **Action:** BMP (Cr/eGFR), HbA1c. If eGFR < 45, insulin dose reduction and tighter glucose monitoring are recommended. Obtain sliding scale from prescriber.
- **Confidence:** HIGH.

🥈 Rank 2 — Drug–Drug Interaction [HIGH RELEVANCE]

Insulin Glargine + Insulin Lispro → Stacking Risk (Hypoglycemia)

- **Mechanism:** Basal-bolus insulin regimens require careful coordination. If the Humalog sliding scale is applied without accounting for residual Lantus effect (which is active for ~24 hours), insulin stacking can occur — particularly with large sliding scale doses or missed meals.
- **Clinical correlation:** Patient does not have his sliding scale available. If he took a higher Humalog dose than needed at his last meal or dosed without eating sufficient carbohydrates, the combination may have pushed glucose to 54.
- **Action:** Obtain full sliding scale. Document last Humalog dose, timing, and meal intake. Education on meal-dose pairing if not already provided.
- **Confidence:** HIGH for mechanism; moderate for this specific event without sliding scale data.

🥉 Rank 3 — Drug Interaction with OTC / Other Agents [MODERATE RELEVANCE]

Insulin → Masking by Beta-Blockers (If Present) — Assess for Undisclosed Medications

- **Mechanism:** Beta-blockers mask the adrenergic symptoms of hypoglycemia (tremor, tachycardia, anxiety), leaving diaphoresis as the only reliable warning sign. Sweating without racing heart is consistent with beta-blocker–masked hypoglycemia.
- **Clinical correlation:** Patient reports shakiness and sweating — but does he take a beta-blocker (common in T2DM patients for cardiovascular protection)? This was not disclosed. Symptom profile (sweating but no reported palpitations) warrants investigation.
- **Action:** Screen for any undisclosed beta-blocker use. If present, adjust hypoglycemia education — patient must rely on blood glucose monitoring rather than symptoms.
- **Confidence:** MODERATE — conditional on undisclosed beta-blocker use; still warrants screening.

SCENARIO 6

Patient Demographics

- **Name:** Linda O'Connor
- **Age:** 74
- **Sex:** Female
- **Setting:** Emergency Department

Chief Complaint

"I've been feeling weak and confused since yesterday. My legs feel heavy and I've been peeing a lot less than normal. My family said I seemed 'off.'"

Medication Sources

Source 1 — Discharge Paperwork (Recent Hospitalization)

```
Discharge Medications
- Lasix (furosemide) 20 mg daily
- Potassium chloride 10 mEq daily
- Lisinopril 10 mg daily
- Protonix (pantoprazole) 40 mg daily
```

Source 2 — Pill Bottle (Physical)

Patient presents a bottle: bottle label reads furosemide 20 mg — consistent with discharge paperwork.

Source 3 — Verbal Report

"My primary doctor stopped one of them, but I kept taking it because I had pills left." [Patient cannot clarify which medication was stopped.]

Reconciled Medication List

Medication	Dose	Frequency	Source
Furosemide (Lasix)	20 mg	Once daily	Discharge paperwork + pill bottle (confirmed)
Potassium chloride	10 mEq	Once daily	Discharge paperwork only
Lisinopril	10 mg	Once daily	Discharge paperwork only
Pantoprazole (Protonix)	40 mg	Once daily	Discharge paperwork only

Reconciliation Flags

- **CRITICAL:** Patient reports one medication was stopped by PCP but she continued it — identity unknown. Must contact PCP urgently.

- Decreased urine output + confusion in a 74F on diuretic + ACE inhibitor = high suspicion for acute kidney injury (AKI)
- Potassium chloride on top of lisinopril — hyperkalemia risk in setting of renal impairment

---

Top 3 Ranked Interactions

🥇 Rank 1 — Drug–Drug + Drug–Disease Interaction [CRITICAL]

Furosemide + Lisinopril → AKI Risk / Volume Depletion in Elderly

- **Mechanism:** Loop diuretics (furosemide) cause volume depletion; ACE inhibitors (lisinopril) reduce efferent arteriolar tone in the kidney, maintaining GFR only when adequate perfusion pressure is present. In a volume-depleted patient, this combination can precipitate acute kidney injury by reducing glomerular filtration pressure.
- **Clinical correlation:** Patient is 74F with confusion, weakness, and decreased urine output — a classic AKI presentation. She is on both furosemide and lisinopril. This is the highest-priority interaction to evaluate.
- **Action:** STAT BMP (Cr, BUN, K^+, Na^+). Hold furosemide and lisinopril pending renal function results. IV fluid resuscitation if volume-depleted. Confusion may reflect uremia or hyponatremia.
- **Confidence:** CRITICAL — symptoms, age, and medication combination are directly consistent.

🥈 Rank 2 — Drug–Drug + Drug–Lab Interaction [HIGH RELEVANCE]

Lisinopril + Potassium Chloride → Hyperkalemia (Amplified by AKI)

- **Mechanism:** ACE inhibitors reduce aldosterone, impairing potassium excretion. Supplemental potassium chloride further raises serum K^+. If the patient is developing AKI (as suspected above), this combination can rapidly produce dangerous hyperkalemia.
- **Clinical correlation:** K^+ supplementation is appropriate when furosemide causes hypokalemia, but in the setting of ACE inhibitor use and now possible AKI, potassium can spike rapidly. Cardiac arrhythmia risk is real.
- **Action:** STAT potassium level. EKG if $K^+ > 5.5$ mEq/L. Hold potassium supplementation until renal function established. Cardiology telemetry.
- **Confidence:** HIGH — especially urgent given clinical presentation.

🥉 Rank 3 — Medication Adherence / Discontinuation Flag [HIGH RELEVANCE]

Unknown Stopped Medication — Continued Without Authorization

- **Mechanism:** Patient reports her PCP stopped a medication, but she continued taking it. Any of the four listed drugs could carry significant risk if continued inappropriately (e.g., potassium supplementation without a diuretic; lisinopril after a new renal diagnosis).
- **Clinical correlation:** The unknown stopped medication could be directly contributing to her current presentation. This is a reconciliation gap that cannot be resolved without PCP contact.
- **Action:** Contact PCP immediately for medication reconciliation. Document all four medications as "status uncertain — continued per patient." Do not assume any of the four are currently indicated without verification.
- **Confidence:** HIGH uncertainty — requires urgent pharmacist or PCP contact.

SCENARIO 7

Patient Demographics

- **Name:** Thomas Nguyen
- **Age:** 68
- **Sex:** Male
- **Setting:** Emergency Department

Chief Complaint

"I've been having muscle aches in my thighs for the past month. I don't exercise much. My urine looked a little dark this morning — I thought I was just dehydrated."

Medication Sources

Source 1 — Printed Medication List (From PCP)

```
Thomas Nguyen – Medications
- Amlodipine 10 mg daily
- Hydrochlorothiazide 25 mg daily
- Atorvastatin 40 mg nightly
- Metformin 1000 mg twice daily
- Vitamin D3 1000 IU daily
```

Reconciled Medication List

Medication	Dose	Frequency	Source
Amlodipine	10 mg	Once daily	Printed list (confirmed)
Hydrochlorothiazide	25 mg	Once daily	Printed list (confirmed)

Medication	Dose	Frequency	Source
Atorvastatin	40 mg	Once daily (PM)	Printed list (confirmed)
Metformin	1000 mg	Twice daily	Printed list (confirmed)
Vitamin D3	1000 IU	Once daily	Printed list (confirmed)

Reconciliation Flags

- Dark urine + myalgia on a statin = must rule out rhabdomyolysis immediately
- Metformin 1000 mg BID — confirm renal function given age and HCTZ-induced volume depletion

---

Top 3 Ranked Interactions

🥇 Rank 1 — Drug–Symptom Interaction [CRITICAL]

Atorvastatin → Statin-Induced Myopathy / Rhabdomyolysis

- **Mechanism:** Statins inhibit HMG-CoA reductase, which reduces mevalonate — a precursor for coenzyme Q10 and other molecules involved in mitochondrial muscle metabolism. This can lead to myopathy (muscle pain and weakness) or, in severe cases, rhabdomyolysis (muscle fiber breakdown releasing myoglobin into the urine).
- **Clinical correlation:** Patient presents with proximal muscle aches (thighs, one month duration) and dark urine — dark urine is a red flag for myoglobinuria. Atorvastatin 40 mg is a moderate-to-high intensity statin. Risk increases with age (68M), dehydration (likely given HCTZ), and possible CYP3A4 interactions.
- **Action:** STAT CK level (rhabdomyolysis if CK > 10x upper limit of normal), urinalysis (myoglobinuria), BMP (renal function). Hold atorvastatin. Aggressive IV hydration if rhabdomyolysis confirmed.
- **Confidence:** CRITICAL — dark urine + proximal myalgia on a statin is rhabdomyolysis until proven otherwise.

🥈 Rank 2 — Drug–Lab + Drug–Disease Interaction [HIGH RELEVANCE]

Metformin → Renal Function (Amplified by Volume Depletion from HCTZ)

- **Mechanism:** HCTZ causes volume depletion, which reduces renal perfusion and transiently lowers eGFR. If rhabdomyolysis is confirmed, the myoglobin-induced AKI will further impair renal function. Metformin is contraindicated in significant renal impairment (eGFR < 30) due to lactic acidosis risk.
- **Clinical correlation:** Patient is on both HCTZ (volume depletion risk) and metformin, and may be developing AKI from rhabdomyolysis. This creates a cascading renal drug interaction.

- **Action:** Hold metformin immediately. BMP for eGFR. Restart only when renal function normalized and confirmed.
- **Confidence:** HIGH — especially urgent in context of suspected rhabdomyolysis.

🥉 Rank 3 — Drug–Drug Interaction [MODERATE RELEVANCE]

Hydrochlorothiazide → Glucose Dysregulation (Metformin Offset)

- **Mechanism:** Thiazide diuretics impair insulin secretion and increase insulin resistance by promoting hypokalemia and direct pancreatic beta-cell effects. This can raise blood glucose levels in diabetic patients, partially offsetting metformin's glucose-lowering effect.
- **Clinical correlation:** Patient is on both HCTZ and metformin. If glucose is uncontrolled, the HCTZ effect may be a contributing factor.
- **Action:** Check HbA1c and fasting glucose. If glycemic control is poor, assess HCTZ contribution before escalating diabetes therapy. Alternative antihypertensive class (ACE inhibitor, ARB) may be more metabolically favorable in diabetic patients.
- **Confidence:** MODERATE — clinically relevant for longitudinal management; lower immediate priority in this acute visit.

SCENARIO 8

Patient Demographics

- **Name:** Carolyn Brooks
- **Age:** 73
- **Sex:** Female
- **Setting:** Emergency Department

Chief Complaint

"I've been having joint pain, especially in my hands, and I've been very tired. I also noticed a rash on my cheeks that's been there for a few weeks."

Medication Sources

Source 1 — Handwritten List (Brand Names Only)

```
Carolyn Brooks – Medications (as she remembers them)
- Norvasc 5 mg daily
- Lipitor 20 mg nightly
```

- Synthroid 88 mcg daily

Reconciled Medication List

Medication	Generic Name	Dose	Frequency	Source
Norvasc (amlodipine)	Amlodipine	5 mg	Once daily	Written list (brand name)
Lipitor (atorvastatin)	Atorvastatin	20 mg	Once daily (PM)	Written list (brand name)
Synthroid (levothyroxine)	Levothyroxine	88 mcg	Once daily	Written list (brand name)

Reconciliation Flags

- All three medications listed by brand name only — generic equivalents confirmed above
- Rash + joint pain + fatigue: may suggest drug-induced lupus (DIL) — screen medications for known DIL-causing agents
- Hypothyroidism (levothyroxine) must be well-controlled to assess whether symptoms are thyroid-related

---

Top 3 Ranked Interactions

🥇 Rank 1 — Drug–Disease Interaction [HIGH CLINICAL RELEVANCE]

Amlodipine (CCB) → Drug-Induced Lupus (DIL) — Rare but Documented

- **Mechanism:** While hydralazine, procainamide, and isoniazid are the classic DIL-causing agents, calcium channel blockers (particularly amlodipine) have been associated with subacute cutaneous lupus erythematosus (SCLE) and lupus-like reactions in case series and pharmacovigilance databases.
- **Clinical correlation:** Patient is 73F with malar rash, joint pain, and fatigue — the classic lupus triad. Autoimmune workup (ANA, anti-histone antibodies, anti-Ro/SSA) is needed. Age of onset in a patient on chronic amlodipine suggests drug-induced rather than idiopathic SLE.
- **Action:** Rheumatology consult. ANA, anti-dsDNA, anti-histone, anti-Ro/SSA, complement (C3/C4), CBC with differential. If DIL confirmed, trial discontinuation of amlodipine with BP management via another class.
- **Confidence:** MODERATE-HIGH — presentation is highly consistent; DIL must be excluded before other diagnoses.

🥈 Rank 2 — Drug–Symptom Interaction [MODERATE RELEVANCE]

Levothyroxine (Suboptimal Dosing) → Fatigue / Joint Pain / Myalgia

- **Mechanism:** Untreated or undertreated hypothyroidism causes fatigue, myalgia, arthralgia, and cold intolerance. Conversely, over-replacement causes palpitations, anxiety, and bone loss. Symptoms overlap significantly with autoimmune disease.
- **Clinical correlation:** Before attributing fatigue and joint pain to drug-induced lupus, confirm that thyroid function is well-controlled. Hypothyroid arthropathy and myopathy can mimic inflammatory arthritis.
- **Action:** TSH (with free T4). If TSH is elevated, this may be the primary driver of her fatigue and joint complaints, independent of the rash.
- **Confidence:** MODERATE — thyroid disease is a common mimicker; essential to rule out before committing to an autoimmune workup.

🥉 Rank 3 — Drug–Drug Interaction [LOW-MODERATE RELEVANCE]

Atorvastatin → Myalgia (Overlapping with Hypothyroid Myopathy)

- **Mechanism:** Statin-induced myopathy presents as proximal muscle aching and fatigue, which can overlap with both hypothyroid myopathy and lupus myositis. Hypothyroidism increases statin myopathy risk by reducing CYP3A4 activity.
- **Clinical correlation:** If levothyroxine is under-dosed (elevated TSH), atorvastatin myopathy risk is amplified. Joint and muscle pain may have a pharmacologic rather than autoimmune cause.
- **Action:** CK level, TSH. If both are elevated, hold atorvastatin temporarily and optimize levothyroxine before re-challenging.
- **Confidence:** LOW-MODERATE — dependent on TSH result; relevant only if hypothyroid state confirmed.

SCENARIO 9

Patient Demographics

- **Name:** Edward Collins
- **Age:** 76
- **Sex:** Male
- **Setting:** Emergency Department

Chief Complaint

"I've been feeling dizzy and unsteady on my feet. I fell this morning getting out of bed. I also feel a bit drowsy all the time."

Medication Sources

Source 1 — Verbal Report (Doses Unknown)

"I take gabapentin three times a day. I also take lisinopril once a day." (Doses unknown.)

Source 2 — Pill Bottle (Physical)

Patient hands over a bottle: "This is the nerve pain medicine."

- **Label reads:** Gabapentin 300 mg — Take 1 capsule by mouth three times daily

Reconciled Medication List

Medication	Dose	Frequency	Source
Gabapentin	300 mg	Three times daily (900 mg/day total)	Verbal + pill bottle (confirmed)
Lisinopril	Unknown dose	Once daily	Verbal only

Reconciliation Flags

- Lisinopril dose not confirmed — contact pharmacy
- Gabapentin 900 mg/day in a 76-year-old — assess for age-related clearance reduction; renal function critical
- Fall + dizziness + drowsiness = high Beers Criteria concern for CNS depressant use in elderly

---

Top 3 Ranked Interactions

🥇 Rank 1 — Drug–Disease Interaction (Geriatric) [HIGH CLINICAL RELEVANCE]

Gabapentin → CNS Depression / Fall Risk in Elderly (Beers Criteria)

- **Mechanism:** Gabapentin is a CNS depressant that causes dose-dependent sedation, dizziness, and ataxia. In older adults, these effects are amplified due to age-related changes in drug distribution, reduced renal clearance (gabapentin is renally eliminated), and increased CNS sensitivity. Gabapentin is listed in the AGS Beers Criteria as a drug to use with caution in older adults due to fall and fracture risk.
- **Clinical correlation:** Patient is 76 years old, presenting with dizziness, unsteadiness, drowsiness, and a fall — all cardinal signs of gabapentin toxicity or accumulation.
- **Action:** Renal function (BMP — if eGFR is reduced, gabapentin accumulates significantly). Consider dose reduction. Fall risk assessment. Occupational therapy / physical therapy referral. Check for other CNS-active medications not yet disclosed.

- **Confidence:** CRITICAL — Beers Criteria drug + classic presentation in high-risk age group.

🥈 Rank 2 — Drug–Drug Interaction [MODERATE RELEVANCE]

Gabapentin + Lisinopril → Additive Hypotension (Orthostatic Fall Risk)

- **Mechanism:** Gabapentin can cause orthostatic hypotension through CNS vasomotor depression. Lisinopril (ACE inhibitor) adds antihypertensive effect. In combination, postural blood pressure drops may be significant enough to cause falls in elderly patients with impaired baroreflex function.
- **Clinical correlation:** Patient fell getting out of bed — a classically orthostatic moment. The combination of gabapentin's vasodepressor effect and lisinopril's antihypertensive action may be producing unsafe postural hypotension.
- **Action:** Orthostatic vitals (lying, sitting, standing BP and HR). If significant orthostatic drop, hold or reduce gabapentin and reassess lisinopril dose. Confirm lisinopril dose with pharmacy.
- **Confidence:** MODERATE — orthostatic mechanism is plausible given presentation.

🥉 Rank 3 — Drug–Lab Interaction [MODERATE RELEVANCE]

Gabapentin → Renal Function Dependency (eGFR-Based Dosing)

- **Mechanism:** Gabapentin is almost entirely renally eliminated without hepatic metabolism. As eGFR declines, gabapentin half-life increases dramatically. A patient on 300 mg TID (900 mg/day) with eGFR of 30 may have drug exposure equivalent to 2700 mg/day in a patient with normal renal function.
- **Clinical correlation:** 76-year-old male with drowsiness and ataxia — these are hallmarks of gabapentin accumulation in the setting of occult renal impairment. Even if his creatinine appears "normal" for his age, eGFR may be substantially reduced.
- **Action:** BMP — calculate eGFR using CKD-EPI formula. Gabapentin dose-adjust per GFR: 300 mg BID for eGFR 30–59; 300 mg once daily for eGFR 15–29. Consider holding if acute presentation warrants.
- **Confidence:** HIGH — gabapentin's renal elimination makes this clinically essential to evaluate.

SCENARIO 10

Patient Demographics

- **Name:** Margaret Feldman

- **Age:** 70
- **Sex:** Female
- **Setting:** Emergency Department

Chief Complaint

"I came in because my primary care doctor sent me. My sodium was low on a blood test last week. I feel fine, but she was worried."

Medication Sources

Source 1 — Written Medication List (Patient-Provided)

```
Margaret Feldman – Medications
- Sertraline 50 mg daily
- Aspirin 81 mg daily
- Multivitamin daily
- Melatonin 5 mg at bedtime
```

Reconciled Medication List

Medication	Dose	Frequency	Source
Sertraline	50 mg	Once daily	Written list (confirmed)
Aspirin	81 mg	Once daily	Written list (confirmed)
Multivitamin	Unknown composition	Once daily	Written list (confirmed)
Melatonin	5 mg	Once daily (bedtime)	Written list (confirmed)

Reconciliation Flags

- Hyponatremia documented by PCP — medication-induced SIADH must be evaluated
- Multivitamin composition unknown — verify for vitamin E (additive antiplatelet with aspirin), iron, or other interacting components

---

Top 3 Ranked Interactions

🥇 Rank 1 — Drug–Lab Interaction [HIGH CLINICAL RELEVANCE]

Sertraline → SIADH-Induced Hyponatremia

- **Mechanism:** SSRIs (including sertraline) cause syndrome of inappropriate antidiuretic hormone secretion (SIADH) in a dose-dependent fashion. ADH retention causes

dilutional hyponatremia. Risk is highest in women, the elderly, patients on low-sodium diets, and those with low BMI.

- **Clinical correlation:** Patient is 70F with documented hyponatremia on sertraline. Even at 50 mg daily, SSRI-induced SIADH is a well-established and common cause of hyponatremia in older adults. She is asymptomatic, which is consistent with gradual/chronic hyponatremia.
- **Action:** Repeat serum Na^+, serum osmolality, urine osmolality, urine Na^+ (SIADH pattern: high urine osmolality + high urine Na^+ + low serum Na^+). If SIADH confirmed, consider dose reduction, fluid restriction, or switching to a different antidepressant class.
- **Confidence:** HIGH — sertraline is a leading cause of drug-induced SIADH in older women.

🥈 Rank 2 — Drug–Drug Interaction [MODERATE RELEVANCE]

Sertraline + Aspirin → Increased GI Bleeding Risk

- **Mechanism:** SSRIs deplete platelet serotonin, impairing platelet aggregation. Aspirin irreversibly inhibits COX-1-mediated thromboxane A2, further reducing platelet function. The combination significantly increases the risk of gastrointestinal bleeding (relative risk approximately 3–15x vs. aspirin alone, depending on study).
- **Clinical correlation:** Patient is on both. At 70 years old, GI mucosal fragility increases risk further. Ask about any GI symptoms (melena, hematemesis, heartburn).
- **Action:** Assess for GI symptoms, stool guaiac. Confirm aspirin indication (primary vs. secondary prevention). If primary prevention only, guidelines in 2022+ recommend against routine low-dose aspirin in patients over 60 due to bleeding risk. Consider adding PPI (omeprazole/pantoprazole) if aspirin is continued.
- **Confidence:** HIGH for interaction; clinical significance depends on aspirin indication and GI symptom assessment.

🥉 Rank 3 — OTC Interaction [LOW-MODERATE RELEVANCE]

Melatonin (5 mg) + Sertraline → Additive Serotonergic / Sedative Effects

- **Mechanism:** Melatonin at higher doses (5 mg, as opposed to the physiologic 0.1–0.5 mg) can interact with serotonergic pathways. Some case reports and pharmacodynamic modeling suggest additive sedation and serotonergic augmentation with SSRI co-administration, though serotonin syndrome at these doses is rare.
- **Clinical correlation:** 5 mg melatonin is a supratherapeutic dose (physiologic nighttime levels are much lower). In an elderly patient on sertraline, nighttime over-sedation and morning grogginess may result.
- **Action:** Recommend melatonin dose reduction to 0.5–1 mg IR formulation for sleep initiation. Assess for daytime sedation, morning confusion, or falls.
- **Confidence:** LOW-MODERATE — serotonin syndrome risk is low; sedation risk in elderly is moderate.

SCENARIO 11

Patient Demographics

- **Name:** Dorothy Huang
- **Age:** 84
- **Sex:** Female
- **Setting:** Emergency Department

Chief Complaint

"My daughter brought me in. I've been confused since last night, not sleeping right, and I tried to get up by myself and almost fell. I take a lot of medications and she's worried about them."

Medication Sources

Source 1 — Medication Bag (Multiple Bottles, Daughter-Assisted)

```
Dorothy Huang – Medications (daughter provided from home bottles)
- Diphenhydramine (Benadryl) 25 mg — "for sleep, takes every night"
- Oxybutynin 5 mg twice daily — "for bladder leakage"
- Donepezil 5 mg once daily — "for memory"
- Metoprolol succinate 25 mg once daily
- Lisinopril 5 mg once daily
```

Source 2 — Verbal (Daughter)

"She also takes over-the-counter Gas-X and Tums on and off. I'm not sure about anything else."

Reconciled Medication List

Medication	Dose	Frequency	Source
Diphenhydramine (Benadryl)	25 mg	Once daily (bedtime)	Bottle confirmed
Oxybutynin	5 mg	Twice daily	Bottle confirmed
Donepezil	5 mg	Once daily	Bottle confirmed
Metoprolol succinate	25 mg	Once daily	Bottle confirmed
Lisinopril	5 mg	Once daily	Bottle confirmed
Simethicone (Gas-X)	Unknown	PRN	Verbal (daughter)
Calcium carbonate (Tums)	Unknown	PRN	Verbal (daughter)

Reconciliation Flags

- CRITICAL: Diphenhydramine (strong anticholinergic) in an 84-year-old on donepezil — directly antagonistic
- CRITICAL: Oxybutynin (strong anticholinergic) in an 84-year-old — Beers Criteria highest-priority concern
- Anticholinergic burden score extremely high — major contributor to delirium

---

Top 3 Ranked Interactions

🥇 Rank 1 — Drug–Drug + Drug–Disease Interaction [CRITICAL]

Diphenhydramine + Oxybutynin → Severe Anticholinergic Burden → Delirium in Elderly

- **Mechanism:** Both diphenhydramine (H1 antihistamine with potent anticholinergic activity) and oxybutynin (muscarinic antagonist) are among the highest anticholinergic burden medications on the AGS Beers Criteria and the Anticholinergic Cognitive Burden (ACB) scale. In older adults, cumulative anticholinergic load causes confusion, urinary retention, constipation, dry mouth, blurred vision, and delirium. Risk is greatest in patients over 80.
- **Clinical correlation:** Patient is 84F presenting with acute confusion, sleep disturbance, and fall risk — a textbook anticholinergic toxidrome in the elderly. The combination of diphenhydramine nightly + oxybutynin twice daily represents an extremely high anticholinergic load.
- **Action:** Discontinue diphenhydramine and oxybutynin immediately. Substitute oxybutynin with mirabegron (beta-3 agonist for overactive bladder — no anticholinergic effect). Cognitive and delirium workup. Nursing/safety sitter.
- **Confidence:** CRITICAL.

🥈 Rank 2 — Drug–Drug Interaction [HIGH RELEVANCE]

Donepezil (Cholinesterase Inhibitor) + Oxybutynin/Diphenhydramine (Anticholinergics) → Direct Pharmacodynamic Antagonism

- **Mechanism:** Donepezil works by increasing central cholinergic tone (inhibiting acetylcholinesterase). Anticholinergic drugs (oxybutynin, diphenhydramine) directly block muscarinic acetylcholine receptors — pharmacodynamically neutralizing donepezil's cognitive benefit. This combination essentially cancels out the dementia medication's effect.
- **Clinical correlation:** Patient is on donepezil for memory impairment while also taking two drugs that actively impair cholinergic neurotransmission. The cognitive decline may be partly drug-induced rather than disease progression.
- **Action:** Discontinue anticholinergics. Reassess cognitive status after washout (at least 4–6 weeks). Communicate with her geriatric or neurology team.
- **Confidence:** HIGH — direct mechanistic antagonism with high clinical impact.

🥉 Rank 3 — Drug–Drug Interaction [MODERATE RELEVANCE]

Calcium Carbonate (Tums) → Absorption Interference with Multiple Medications

- **Mechanism:** Calcium carbonate (Tums) raises gastric pH and chelates certain medications, potentially reducing absorption of donepezil, metoprolol, and levothyroxine (if present). Timing relative to other medications matters significantly.
- **Clinical correlation:** Intermittent Tums use is low-risk, but if taken with morning medications, may reduce efficacy of critical agents.
- **Action:** Counsel family to take Tums at least 2 hours apart from scheduled medications. Low immediate priority given acute presentation, but relevant for discharge counseling.
- **Confidence:** LOW-MODERATE — depends on timing and frequency of Tums use.

SCENARIO 12

Patient Demographics

- **Name:** James Whitfield
- **Age:** 58
- **Sex:** Male
- **Setting:** Emergency Department

Chief Complaint

"I came in because I've been having chest pain that comes and goes over the past two days. I also take a medication for depression and I started a new antibiotic last week for a sinus infection."

Medication Sources

Source 1 — Written List (From Wallet Card)

```
James Whitfield – Medications
- Citalopram 40 mg daily
- Atorvastatin 40 mg nightly
- Metformin 500 mg twice daily
- Azithromycin 500 mg daily (recently started — 5-day course)
```

Reconciled Medication List

Medication	Dose	Frequency	Source
Citalopram	40 mg	Once daily	Wallet card confirmed

Medication	Dose	Frequency	Source
Atorvastatin	40 mg	Once daily (PM)	Wallet card confirmed
Metformin	500 mg	Twice daily	Wallet card confirmed
Azithromycin	500 mg	Once daily	Wallet card (recent addition)

Reconciliation Flags

- CRITICAL: Citalopram 40 mg + Azithromycin — both prolong QTc interval; combination is high-risk for Torsades de Pointes
- Chest pain in this context must prompt immediate EKG

Top 3 Ranked Interactions

🥇 Rank 1 — Drug–Drug Interaction [CRITICAL]

Citalopram + Azithromycin → QTc Prolongation → Risk of Torsades de Pointes

- **Mechanism:** Citalopram dose-dependently prolongs the cardiac QT interval by blocking the hERG potassium channel (IKr), with 40 mg being the FDA maximum approved dose due to QT concerns. Azithromycin (a macrolide antibiotic) also blocks IKr and independently prolongs QTc. In combination, additive QT prolongation raises risk for Torsades de Pointes (TdP), a potentially fatal polymorphic ventricular tachycardia.
- **Clinical correlation:** Patient is 58M with chest pain starting two days ago — after beginning azithromycin. The temporal relationship is highly suspicious. QTc > 500 ms is associated with substantially elevated TdP risk.
- **Action:** STAT EKG — measure QTc. If QTc > 500 ms or > 60 ms increase from baseline, discontinue azithromycin immediately and consider telemetry. Switch to amoxicillin or doxycycline for sinusitis if antibiotic still needed. Electrolytes (K^+, Mg^{2+}) — hypokalemia/hypomagnesemia amplify QT risk.
- **Confidence:** CRITICAL — this is a well-documented, high-risk drug interaction.

🥈 Rank 2 — Drug–Drug Interaction [MODERATE RELEVANCE]

Azithromycin → CYP3A4 Inhibition → Atorvastatin Exposure Increase

- **Mechanism:** Azithromycin is a mild CYP3A4 inhibitor. Atorvastatin is metabolized by CYP3A4. Inhibition increases atorvastatin plasma levels, theoretically increasing myopathy risk. Effect is mild and short-term (5-day course), but worth flagging.
- **Clinical correlation:** Not the primary concern in this visit, but worth monitoring for myalgia if the antibiotic course was longer.
- **Action:** Monitor for muscle symptoms. Given the 5-day course, risk is low but should be documented.

- **Confidence:** LOW-MODERATE — short course limits clinical impact.

🥉 Rank 3 — Drug–Symptom Interaction [MODERATE RELEVANCE]

Citalopram (SSRI) → Chest Pain / Cardiac Awareness

- **Mechanism:** SSRIs can cause chest tightness, palpitations, and heightened somatic awareness as side effects, particularly early in treatment or with dose changes. These symptoms can mimic or coexist with cardiac pathology.
- **Clinical correlation:** If citalopram was recently started or the dose increased, SSRI-related chest symptoms are possible. However, this is a diagnosis of exclusion — QT-related arrhythmia must be ruled out first.
- **Action:** Rule out QT-related arrhythmia (EKG, troponin) before attributing symptoms to SSRI effect. Document any recent dose changes.
- **Confidence:** LOW — exclusion diagnosis only.

SCENARIO 13

Patient Demographics

- **Name:** Gloria Martinez
- **Age:** 67
- **Sex:** Female
- **Setting:** Emergency Department

Chief Complaint

"I've had a bad headache for two days, my blood pressure at home was 190/105, and I've been taking a lot of ibuprofen for the headache."

Medication Sources

Source 1 — Verbal Report

"I take amlodipine 10 mg every day and I think I take another blood pressure pill — ramipril, 5 mg. I've been taking ibuprofen 400 mg about three times a day for this headache."

Source 2 — Pill Bottles (Two Brought)

- **Bottle 1:** Amlodipine 10 mg — confirmed verbal report

- **Bottle 2:** Ramipril 5 mg — confirmed verbal report

Reconciled Medication List

Medication	Dose	Frequency	Source
Amlodipine	10 mg	Once daily	Verbal + bottle confirmed
Ramipril	5 mg	Once daily (assumed)	Verbal + bottle confirmed
Ibuprofen (OTC)	400 mg	Three times daily (self-reported, recent)	Verbal only

Reconciliation Flags

- NSAID use actively counteracting antihypertensive therapy — likely contributor to BP crisis
- Ramipril + ibuprofen = nephrotoxic combination in the setting of hypertensive urgency

---

Top 3 Ranked Interactions

🥇 Rank 1 — Drug–Drug Interaction [CRITICAL]

Ibuprofen (NSAID) → Antihypertensive Resistance (Amlodipine + Ramipril Blunting)

- **Mechanism:** NSAIDs inhibit prostaglandin synthesis (COX-1 and COX-2), causing sodium and water retention and vasoconstriction. This directly counteracts the mechanisms of both CCBs (amlodipine) and ACE inhibitors (ramipril). Even short-term NSAID use (3–5 days) can raise systolic BP by 3–5 mmHg on average, with higher surges in patients on antihypertensives.
- **Clinical correlation:** Patient's BP is 190/105 despite being on two antihypertensives. She has been taking ibuprofen TID for two days — this is very likely the pharmacologic cause of her hypertensive urgency.
- **Action:** Discontinue ibuprofen immediately. Acetaminophen is the safe alternative for pain in hypertensive patients. Reassess BP after NSAID washout (12–24 hours). If headache is the result of elevated BP, treating the BP may resolve it.
- **Confidence:** CRITICAL — temporal relationship is direct; NSAIDs are a classic cause of antihypertensive failure.

🥈 Rank 2 — Drug–Drug + Drug–Disease Interaction [HIGH RELEVANCE]

Ibuprofen + Ramipril (ACE Inhibitor) → Acute Kidney Injury Risk

- **Mechanism:** ACE inhibitors dilate the efferent glomerular arteriole to maintain filtration. NSAIDs vasoconstrict the afferent arteriole by reducing prostaglandin-mediated dilation. In combination (the "triple whammy" is ACE inhibitor + diuretic + NSAID, but the dual

combination alone is nephrotoxic), glomerular filtration pressure drops precipitously. This causes AKI, particularly in volume-depleted or elderly patients.
- **Clinical correlation:** Patient is on ramipril + self-administering ibuprofen TID during a hypertensive urgency — she may already have reduced renal perfusion.
- **Action:** BMP (Cr, BUN, K^+). Hold ibuprofen. Monitor urine output. If eGFR is declining, hold ramipril temporarily during the acute period.
- **Confidence:** HIGH.

🥉 Rank 3 — Drug–Symptom Interaction [MODERATE RELEVANCE]

Hypertensive Headache vs. Analgesic Rebound Headache

- **Mechanism:** Ibuprofen taken daily for more than 10–15 days can cause medication-overuse headache (MOH), also called rebound headache. This creates a self-perpetuating cycle where the patient takes more analgesic to treat the headache caused by the analgesic.
- **Clinical correlation:** Patient has been taking ibuprofen TID for at least two days for a headache — if this has been ongoing longer, MOH must be considered as a contributor to the unremitting headache.
- **Action:** Detailed headache history (duration, prior episodes, analgesic use patterns). If MOH suspected, education on rebound headache and transition to scheduled acetaminophen with gradual analgesic weaning.
- **Confidence:** MODERATE — requires further history to confirm.

SCENARIO 14

Patient Demographics

- **Name:** Harold Jensen
- **Age:** 80
- **Sex:** Male
- **Setting:** Emergency Department

Chief Complaint

"I've been really constipated for a week and my belly is bloated. I also started taking something for bone strength my doctor suggested."

Medication Sources

Source 1 — Written List (From Spouse)

```
Harold Jensen – Medications
- Amlodipine 5 mg daily
- Atorvastatin 20 mg nightly
- Tamsulosin 0.4 mg daily (for prostate)
- Alendronate 70 mg once weekly (recently started)
- Calcium carbonate 500 mg twice daily (recently started with alendronate)
- Vitamin D 800 IU daily
```

Source 2 — Verbal (Spouse)

"His doctor started the bone medicine and calcium a few weeks ago. He also takes Milk of Magnesia when he's backed up — he's used it a few times this week."

Reconciled Medication List

Medication	Dose	Frequency	Source
Amlodipine	5 mg	Once daily	Written list confirmed
Atorvastatin	20 mg	Once daily (PM)	Written list confirmed
Tamsulosin	0.4 mg	Once daily	Written list confirmed
Alendronate	70 mg	Once weekly	Written list confirmed
Calcium carbonate	500 mg	Twice daily	Written list confirmed
Vitamin D	800 IU	Once daily	Written list confirmed
Milk of Magnesia (Mg hydroxide)	Unknown dose	PRN (recent use)	Verbal (spouse)

Reconciliation Flags

- Alendronate administration instructions critical — must be taken correctly or causes esophageal injury
- Calcium carbonate + alendronate timing conflict — must be separated
- Tamsulosin: orthostatic hypotension risk especially combined with amlodipine in an 80-year-old

---

Top 3 Ranked Interactions

🥇 Rank 1 — Drug–Drug Interaction [HIGH RELEVANCE]

Calcium Carbonate → Alendronate Absorption (Near-Complete Blockade)

- **Mechanism:** Alendronate (bisphosphonate) has extremely poor oral bioavailability (~1%) that is critically dependent on fasting conditions. Calcium, antacids, food, and other minerals bind alendronate in the GI tract, reducing absorption by up to 60–90%.

The drug must be taken on an empty stomach with a full glass of water, sitting upright, 30–60 minutes before any other food, drink, or medication.

- **Clinical correlation:** Patient is on both alendronate and calcium carbonate twice daily — if calcium is taken too close to alendronate (even within 2 hours), the bisphosphonate may be nearly completely unabsorbed, rendering therapy ineffective.
- **Action:** Educate patient on alendronate administration: AM on empty stomach, full glass of water, upright position for 30 minutes. Calcium and all other medications should be taken at least 30–60 minutes later. Separate calcium doses from alendronate administration window.
- **Confidence:** HIGH — this is a well-documented, clinically significant absorption interaction.

🥈 Rank 2 — Drug–Symptom Interaction [MODERATE-HIGH RELEVANCE]

Calcium Carbonate (High Dose) → Constipation + Milk-Alkali Syndrome Risk

- **Mechanism:** Calcium carbonate is a known cause of constipation through slowing colonic transit. At doses ≥1000 mg/day (patient is on 500 mg BID = 1000 mg/day), constipation is common. Additionally, high-dose calcium carbonate combined with Milk of Magnesia and vitamin D can cause milk-alkali syndrome (hypercalcemia, metabolic alkalosis, renal impairment).
- **Clinical correlation:** Patient has severe constipation and bloating starting after initiating calcium + alendronate. Calcium carbonate-induced constipation is the most likely cause.
- **Action:** Switch to calcium citrate (less constipating, better absorbed, doesn't require stomach acid). Stop Milk of Magnesia as a constipation treatment while assessing for milk-alkali syndrome. Check serum Ca^{2+}, BMP.
- **Confidence:** HIGH for constipation causation.

🥉 Rank 3 — Drug–Drug Interaction (Geriatric) [MODERATE RELEVANCE]

Tamsulosin + Amlodipine → First-Dose and Orthostatic Hypotension in Elderly

- **Mechanism:** Alpha-1 blockers (tamsulosin) and calcium channel blockers (amlodipine) both cause vasodilation and reduction in blood pressure. In older adults, baroreceptor sensitivity is blunted, making postural hypotension more likely. Risk is highest immediately after a dose or upon standing.
- **Clinical correlation:** 80-year-old male on dual vasodilatory therapy — fall and orthostatic hypotension risk is elevated. This is a Beers Criteria–relevant combination.
- **Action:** Orthostatic vitals. Fall risk education. Confirm whether antihypertensive effect of amlodipine is appropriate given current BP.
- **Confidence:** MODERATE — important for fall prevention in discharge planning.

SCENARIO 15

Patient Demographics

- **Name:** Sylvia Okafor
- **Age:** 55
- **Sex:** Female
- **Setting:** Emergency Department

Chief Complaint

"I have HIV and I'm on my medications. I started a new medication for a fungal infection two days ago and since then I've been having severe nausea, vomiting, and I feel drunk — really dizzy."

Medication Sources

Source 1 — Pharmacy Printout (Brought to ED)

```
Sylvia Okafor – Current Medications
- Bictegravir/tenofovir alafenamide/emtricitabine (Biktarvy) 1 tablet daily
- Fluconazole 200 mg daily (new — started 2 days ago, prescribed at urgent
care)
```

Source 2 — Verbal Report

"My HIV doctor gave me Biktarvy. The urgent care gave me the antifungal. They didn't talk to each other."

Reconciled Medication List

Medication	Dose	Frequency	Source
Bictegravir/TAF/FTC (Biktarvy)	1 tablet	Once daily	Pharmacy printout confirmed
Fluconazole	200 mg	Once daily	Pharmacy printout confirmed

Reconciliation Flags

- Fluconazole is a potent CYP3A4 and CYP2C19 inhibitor — interacts with many antiretrovirals
- Symptoms began with fluconazole initiation — pharmacokinetic interaction likely

---

Top 3 Ranked Interactions

🥇 Rank 1 — Drug–Drug Interaction [HIGH CLINICAL RELEVANCE]

Fluconazole → CYP3A4 Inhibition → Bictegravir Plasma Level Increase

- **Mechanism:** Bictegravir is metabolized by CYP3A4 and UGT1A1. Fluconazole is a potent inhibitor of CYP3A4 (and CYP2C19), reducing bictegravir metabolism and increasing plasma drug concentrations. Elevated bictegravir levels cause nausea, vomiting, dizziness, and CNS effects — exactly matching patient's symptoms.
- **Clinical correlation:** Symptoms began 2 days after fluconazole was added — a textbook pharmacokinetic interaction with temporal precision. Urgent care prescribed fluconazole without checking antiretroviral interactions.
- **Action:** Contact HIV specialist immediately. Hold fluconazole pending specialist guidance. Consider alternative antifungals with fewer CYP interactions (e.g., topical agents for candidiasis; IV micafungin or caspofungin for systemic infection). Supportive care for nausea/vomiting. Do not adjust Biktarvy dose without HIV specialist.
- **Confidence:** HIGH — temporal relationship and mechanism are unambiguous.

🥈 Rank 2 — Drug–Drug Interaction [MODERATE RELEVANCE]

Fluconazole → QTc Prolongation (Additive with TAF Component)

- **Mechanism:** Fluconazole prolongs the QT interval by blocking the hERG potassium channel. Although Biktarvy components are not the most potent QT prolongers, combined QTc effects in the setting of drug-induced nausea (with potential electrolyte loss from vomiting) create arrhythmia risk.
- **Clinical correlation:** Patient is vomiting — electrolyte depletion (K^+, Mg^{2+}) amplifies fluconazole's QTc effect.
- **Action:** EKG (QTc measurement). Electrolytes (K^+, Mg^{2+}). Replete if depleted.
- **Confidence:** MODERATE.

🥉 Rank 3 — System/Process Interaction [HIGH RELEVANCE — Non-Pharmacologic]

Siloed Prescribing (Urgent Care + Specialist) → Dangerous Drug Interaction Gap

- **Mechanism:** This is a care coordination failure. The urgent care provider did not review the patient's antiretroviral regimen before prescribing fluconazole — a high-risk antifungal with extensive drug interactions.
- **Clinical correlation:** Patient explicitly states "they didn't talk to each other." This is a sentinel reconciliation event.
- **Action:** Document care coordination failure. Ensure HIV specialist is notified. Provide patient with a comprehensive medication card listing antiretrovirals and key interaction

warnings (e.g., "No azole antifungals without HIV specialist approval"). Flag for pharmacist review on discharge.
- **Confidence:** HIGH — process failure directly caused patient harm.

SCENARIO 16

Patient Demographics

- **Name:** Frank Deluca
- **Age:** 72
- **Sex:** Male
- **Setting:** Emergency Department

Chief Complaint

"I've been having black, tarry stools for the past two days. I feel a little weak and dizzy."

Medication Sources

Source 1 — Written List (Printed, Brought by Son)

```
Frank Deluca – Medications
- Warfarin 5 mg daily (for atrial fibrillation)
- Clopidogrel 75 mg daily (after coronary stent, 1 year ago)
- Aspirin 81 mg daily
- Pantoprazole 40 mg daily
- Atorvastatin 40 mg nightly
```

Source 2 — Verbal (Son)

"He also takes fish oil — two capsules a day. He started that on his own from the health food store."

Reconciled Medication List

Medication	Dose	Frequency	Source
Warfarin	5 mg	Once daily	Written list confirmed
Clopidogrel	75 mg	Once daily	Written list confirmed
Aspirin	81 mg	Once daily	Written list confirmed
Pantoprazole	40 mg	Once daily	Written list confirmed

Medication	Dose	Frequency	Source
Atorvastatin	40 mg	Once daily (PM)	Written list confirmed
Fish oil (omega-3)	Unknown dose	Twice daily	Verbal (son)

Reconciliation Flags

- CRITICAL: Triple antithrombotic therapy (warfarin + clopidogrel + aspirin) — GI bleed is extremely high risk
- Black tarry stools = melena = presumed upper GI bleed until proven otherwise
- Fish oil adds additional antiplatelet effect — undisclosed OTC supplement

---

Top 3 Ranked Interactions

🥇 Rank 1 — Drug–Drug Interaction [CRITICAL]

Warfarin + Clopidogrel + Aspirin → Triple Antithrombotic Therapy → GI Hemorrhage

- **Mechanism:** This combination (anticoagulant + two antiplatelets) is sometimes indicated post-stent with atrial fibrillation, but carries dramatically increased major bleeding risk — up to 3–4x vs. dual therapy. Warfarin anticoagulates via clotting factor inhibition; aspirin irreversibly inhibits COX-1 platelet aggregation; clopidogrel inhibits P2Y12 ADP receptor. Each mechanism is additive.
- **Clinical correlation:** Melena (black tarry stool) in a 72-year-old on triple antithrombotic therapy = upper GI hemorrhage until proven otherwise. This is a medical emergency.
- **Action:** STAT CBC (hemoglobin), INR/PT, type and screen. IV access x2. GI consult for urgent endoscopy. Hold all antithrombotics pending hemostasis. Correct coagulopathy (vitamin K, FFP if INR markedly elevated, 4-factor PCC if severe). Confirm whether triple therapy is still indicated (may be appropriate to de-escalate after 1 year post-stent to DAPT or anticoagulant alone).
- **Confidence:** CRITICAL — highest-risk drug interaction in dataset.

🥈 Rank 2 — OTC Supplement + Drug Interaction [HIGH RELEVANCE]

Fish Oil (High-Dose Omega-3) + Warfarin + Antiplatelet Agents → Additional Bleeding Risk

- **Mechanism:** High-dose omega-3 fatty acids (≥1–3 g/day) have antiplatelet and mild anticoagulant properties, potentially enhancing warfarin effect and adding to platelet inhibition from aspirin and clopidogrel. The patient takes "two capsules daily" — dose unknown but common doses are 1–2 g each.
- **Clinical correlation:** Fish oil was not on the written medication list — this is an OTC supplement that the son revealed. It adds yet another antithrombotic agent to an already critically high-risk regimen.

- **Action:** Discontinue fish oil immediately. Document as "undisclosed OTC supplement with antithrombotic effect." Reinforce comprehensive medication reconciliation including all supplements.
- **Confidence:** HIGH — additive to an already critical interaction.

🥉 Rank 3 — Drug Assessment — Protective Interaction (PPI Coverage)

Pantoprazole → Partial GI Protection (Possibly Insufficient for Triple Antithrombotic Regimen)

- **Mechanism:** Proton pump inhibitors reduce gastric acid secretion, protecting the GI mucosa from NSAID/aspirin-induced injury. However, in triple antithrombotic therapy, PPI protection may be insufficient to prevent all bleeding events — it reduces, but does not eliminate, GI hemorrhage risk.
- **Clinical correlation:** Patient is appropriately on pantoprazole with his triple antithrombotic regimen. However, he is still bleeding — confirming that PPI does not fully mitigate risk at this level of antithrombotic therapy.
- **Action:** Continue pantoprazole. Do not interpret PPI use as eliminating bleeding risk assessment in discharge planning. After bleed is controlled, cardiology/hematology review of antithrombotic regimen is essential.
- **Confidence:** HIGH for assessment — reinforces that de-escalation of antithrombotic therapy should be explored.

SCENARIO 17

Patient Demographics

- **Name:** Barbara Kowalski
- **Age:** 77
- **Sex:** Female
- **Setting:** Emergency Department

Chief Complaint

"I've been having tremors in my hands, I've felt stiff, and I've been a bit nauseous. My family said I've been shuffling when I walk, worse than usual."

Medication Sources

Source 1 — Medication List (Printed, From Assisted Living Facility)

```
Barbara Kowalski – Active Medications
- Metoclopramide 10 mg three times daily (for gastroparesis, 8 months)
- Levodopa/carbidopa 25/100 mg three times daily (Parkinson's disease)
- Donepezil 10 mg once daily
- Atorvastatin 40 mg nightly
- Lisinopril 10 mg daily
```

Reconciled Medication List

Medication	Dose	Frequency	Source
Metoclopramide	10 mg	Three times daily	Facility list confirmed
Levodopa/carbidopa	25/100 mg	Three times daily	Facility list confirmed
Donepezil	10 mg	Once daily	Facility list confirmed
Atorvastatin	40 mg	Once daily (PM)	Facility list confirmed
Lisinopril	10 mg	Once daily	Facility list confirmed

Reconciliation Flags

- CRITICAL: Metoclopramide (dopamine antagonist) in a patient on levodopa for Parkinson's disease — direct pharmacodynamic antagonism
- Metoclopramide has been prescribed for 8 months — FDA black box warning for tardive dyskinesia with use > 12 weeks
- Tremor + shuffling = extrapyramidal symptoms (EPS) — likely metoclopramide-induced

---

Top 3 Ranked Interactions

🥇 Rank 1 — Drug–Drug Interaction [CRITICAL]

Metoclopramide (Dopamine Antagonist) + Levodopa (Dopamine Precursor) → Direct Pharmacodynamic Antagonism + Worsening Parkinsonism

- **Mechanism:** Metoclopramide blocks central and peripheral D2 dopamine receptors. This directly counteracts levodopa's mechanism of action (restoring dopaminergic neurotransmission in the basal ganglia). In patients with Parkinson's disease, metoclopramide can precipitate or dramatically worsen parkinsonian symptoms (tremor, rigidity, bradykinesia, shuffling gait).
- **Clinical correlation:** Patient on long-term metoclopramide (8 months) for gastroparesis is developing worsening tremor and shuffling. This is a critical prescribing error — metoclopramide is contraindicated in Parkinson's disease.
- **Action:** Discontinue metoclopramide immediately. Substitute domperidone (preferred in Parkinson's — peripherally acting D2 antagonist with minimal CNS penetration) or low-dose erythromycin for gastroparesis. Neurology/movement disorder specialist consultation. Symptoms may improve gradually over weeks.

- **Confidence:** CRITICAL — textbook contraindicated drug-disease interaction.

🥈 Rank 2 — Drug–Disease Interaction [HIGH RELEVANCE]

Metoclopramide → Tardive Dyskinesia / EPS Risk with Prolonged Use (FDA Black Box)

- **Mechanism:** Metoclopramide carries an FDA black box warning for tardive dyskinesia (TD) — involuntary, repetitive movements — with use beyond 12 weeks. Risk increases with cumulative exposure, older age, and female sex. TD can be irreversible even after discontinuation.
- **Clinical correlation:** This patient has been on metoclopramide for 8 months — nearly 35 weeks past the 12-week safety threshold. She is a 77-year-old female — the highest-risk demographic for TD.
- **Action:** Assess for tardive dyskinesia (orofacial movements, lip smacking, tongue protrusion) in addition to Parkinson's-related symptoms. Report to prescribing provider. Document duration of use and FDA warning exceedance.
- **Confidence:** HIGH — black box violation with high-risk patient profile.

🥉 Rank 3 — Drug–Drug Interaction [MODERATE RELEVANCE]

Donepezil (Cholinesterase Inhibitor) + Metoclopramide → Additive GI Motility Effects / Nausea

- **Mechanism:** Donepezil increases cholinergic activity throughout the body, including in the GI tract (increasing gut motility and secretions). Metoclopramide also enhances GI motility. Together, they can cause additive nausea, cramping, and diarrhea.
- **Clinical correlation:** Patient reports nausea — this may be partly a donepezil + metoclopramide GI effect, though the more urgent concern is the parkinsonian interaction above.
- **Action:** After discontinuing metoclopramide, reassess whether GI symptoms were primarily drug-induced. Donepezil at 10 mg is the higher dose — consider whether 5 mg is sufficient for cognitive benefit.
- **Confidence:** MODERATE — secondary to the higher-priority parkinsonian interaction.

SCENARIO 18

Patient Demographics

- **Name:** Samuel Tran
- **Age:** 63
- **Sex:** Male

- **Setting:** Emergency Department

Chief Complaint

"I've been having episodes of flushing and sweating that come on suddenly. My heart races for a few minutes and then it goes back to normal. I recently started a new supplement for bodybuilding."

Medication Sources

Source 1 — Verbal Report

"I take metoprolol for my heart — I think it's 50 mg once a day. I also take a statin, I can never remember the name. The new thing I started is pre-workout powder — I take it before I go to the gym."

Source 2 — Supplement Container (Physical)

Patient shows container:

- **Label shows:** Pre-workout powder containing caffeine 250 mg/serving, niacin (nicotinic acid) 50 mg/serving, synephrine 50 mg/serving, beta-alanine 3.2 g/serving

Reconciled Medication List

Medication	Dose	Frequency	Source
Metoprolol	50 mg (unconfirmed)	Once daily	Verbal only
Unknown statin	Unknown	Unknown	Verbal only — must verify
Pre-workout supplement	Per serving	PRN before gym	Physical container confirmed

Reconciliation Flags

- Synephrine (sympathomimetic) + metoprolol (beta-blocker) — opposing cardiovascular effects
- Caffeine (high dose) + metoprolol — chronotropic interaction
- Niacin in supplement: flushing, sweating — patient's primary presenting symptom
- Statin type unknown — niacin interaction risk depends on statin identity

---

Top 3 Ranked Interactions

🥇 Rank 1 — Drug–Symptom / OTC Supplement Interaction [HIGH CLINICAL RELEVANCE]

Niacin (50 mg/serving) in Pre-Workout → Flushing / Sweating / Tachycardia (Prostaglandin-Mediated)

- **Mechanism:** Niacin (nicotinic acid) causes prostaglandin D2-mediated vasodilation, producing intense skin flushing, warmth, sweating, and a sensation of racing heart, typically 15–30 minutes after ingestion. This is a pharmacologic effect of niacin, not an allergic reaction — it is dose-dependent and very common.
- **Clinical correlation:** Patient's episodic flushing, sweating, and palpitations that resolve after a few minutes are a textbook description of niacin-induced flushing. The episodes correlate with pre-workout use.
- **Action:** Educate patient on niacin flushing — this is the primary cause of his symptoms. If niacin flushing is intolerable, recommend switching to a niacin-free pre-workout. Aspirin 325 mg taken 30 minutes before niacin can mitigate flushing (prostaglandin inhibition).
- **Confidence:** HIGH — classic presentation with direct supplement correlation.

🥈 Rank 2 — Drug–Supplement Interaction [HIGH RELEVANCE]

Synephrine + Caffeine (High Dose) → Sympathomimetic Surge → Metoprolol Antagonism

- **Mechanism:** Synephrine (a sympathomimetic amine similar to ephedrine) and caffeine (250 mg = ~2.5 cups of coffee equivalent) stimulate adrenergic receptors, increasing heart rate and blood pressure. Metoprolol (a beta-1 selective blocker) blunts the HR response but cannot fully counteract the vasoconstrictive alpha-adrenergic effects of synephrine. Combined, the interaction creates unpredictable cardiovascular stress.
- **Clinical correlation:** Patient reports episodic tachycardia and flushing while taking this supplement despite being on a beta-blocker. The beta-blocker may be partially masking the full sympathomimetic response while allowing unopposed alpha-adrenergic vasoconstriction (reflex hypertension).
- **Action:** Strongly advise discontinuation of synephrine-containing supplements in patients on any cardiac medication. EKG. BP assessment. Verify metoprolol dose with pharmacy.
- **Confidence:** HIGH — synephrine in cardiac patients is a documented risk.

🥉 Rank 3 — Drug–Drug (Statin + Niacin) Interaction [MODERATE RELEVANCE]

Unknown Statin + Niacin (Pre-Workout) → Myopathy Risk

- **Mechanism:** Niacin combined with statins (particularly simvastatin and lovastatin) increases myopathy and rhabdomyolysis risk. At 50 mg/serving (and possibly multiple servings per week), cumulative niacin exposure may be significant. Risk is statin-specific — highest with simvastatin, lower with rosuvastatin.

- **Clinical correlation:** Patient is on an unknown statin. Until the statin is identified, myopathy risk cannot be fully assessed. Given his gym use and the potential for high-exertion exercise, muscle injury risk is additive.
- **Action:** Identify statin immediately (pharmacy contact). If simvastatin or lovastatin, strong recommendation to switch to rosuvastatin or pravastatin and discontinue niacin-containing supplements. CK level.
- **Confidence:** MODERATE — statin identity determines risk level.

SCENARIO 19

Patient Demographics

- **Name:** Alice Pemberton
- **Age:** 89
- **Sex:** Female
- **Setting:** Emergency Department

Chief Complaint

"I've been falling asleep at the dinner table, my speech seems slurred to my daughter, and I've had two falls this week. I live alone. My daughter found me on the floor this morning."

Medication Sources

Source 1 — Medication Bag (Daughter Brought All Bottles)

```
Alice Pemberton - Medications (bottle review)
- Alprazolam (Xanax) 0.5 mg three times daily — prescribed by psychiatrist
- Zolpidem (Ambien) 10 mg at bedtime — prescribed by PCP
- Oxycodone 5 mg every 6 hours PRN (for spinal stenosis pain)
- Metoprolol 25 mg twice daily
- Losartan 25 mg daily
```

Source 2 — Verbal (Daughter)

"She takes all of them. I didn't know she was on so many. She also drinks a small glass of wine at night — has for years."

Reconciled Medication List

Medication	Dose	Frequency	Source
Alprazolam (Xanax)	0.5 mg	Three times daily	Bottle confirmed
Zolpidem (Ambien)	10 mg	Once daily (bedtime)	Bottle confirmed
Oxycodone	5 mg	Every 6 hours PRN	Bottle confirmed
Metoprolol	25 mg	Twice daily	Bottle confirmed
Losartan	25 mg	Once daily	Bottle confirmed
Alcohol (wine)	~1 glass	Daily (evening)	Verbal (daughter)

Reconciliation Flags

- CRITICAL: Triple CNS depression (benzodiazepine + Z-drug + opioid) in 89-year-old = life-threatening respiratory depression and fall risk
- Alcohol + CNS depressant polypharmacy = compound sedation
- Alprazolam is on the Beers Criteria — all three CNS agents are inappropriate in this patient
- Zolpidem 10 mg in an elderly female — FDA recommends 5 mg maximum in women

---

Top 3 Ranked Interactions

🥇 Rank 1 — Drug–Drug Interaction [CRITICAL / LIFE-THREATENING]

Alprazolam + Zolpidem + Oxycodone + Alcohol → CNS/Respiratory Depression in Elderly

- **Mechanism:** Benzodiazepines and Z-drugs (zolpidem) both act on GABA-A receptors, causing sedation and respiratory depression. Opioids (oxycodone) cause respiratory depression via mu-opioid receptors. Alcohol potentiates all CNS depressants. In combination (four CNS depressants simultaneously), the risk of fatal respiratory depression is dramatically elevated. The FDA has issued a Black Box Warning for BZD + opioid co-prescribing due to overdose fatalities.
- **Clinical correlation:** 89-year-old with daytime hypersomnia, slurred speech, and repeated falls — she is experiencing CNS depressant toxidrome. Found on the floor by her daughter suggests she may have had a syncopal/sedation episode.
- **Action:** Respiratory assessment immediately (SpO_2, respiratory rate). If RR < 10 or SpO_2 < 90%, consider naloxone (reverses opioid component). Supportive care. Discontinue zolpidem immediately (5 mg maximum for elderly women). Taper alprazolam (do not abruptly stop — seizure risk). Oxycodone reassessment: is PRN use being used regularly? Telemetry. Social work / safety evaluation for living alone.
- **Confidence:** CRITICAL.

🥈 Rank 2 — Drug–Disease Interaction (Geriatric) [CRITICAL]

Alprazolam + Zolpidem → Beers Criteria: Fall Risk, Cognitive Impairment, Hip Fracture in Elderly

- **Mechanism:** AGS Beers Criteria strongly recommends AVOIDING all benzodiazepines (including short-acting alprazolam) and all Z-drugs (zolpidem, eszopiclone) in adults 65+. These medications impair psychomotor function, gait, balance, and reaction time, tripling fall and hip fracture risk. Cognitive effects (confusion, anterograde amnesia) are also pronounced.
- **Clinical correlation:** Patient is 89, living alone, with two falls this week. Both her benzodiazepine and Z-drug are explicitly contraindicated by geriatric prescribing guidelines.
- **Action:** Geriatrics or palliative care consultation. Deprescribing plan for alprazolam (slow taper) and zolpidem. Non-pharmacologic sleep hygiene. CBT-I (Cognitive Behavioral Therapy for Insomnia) if appropriate. Home safety evaluation.
- **Confidence:** CRITICAL — Beers Criteria highest priority recommendation violated.

🥉 Rank 3 — Drug + Substance Interaction [HIGH RELEVANCE]

Alcohol → Potentiation of All CNS Depressants + Warfarin/Anticoagulant Interaction (If Present)

- **Mechanism:** Alcohol (even one glass daily) significantly potentiates all CNS depressants through GABA-ergic and NMDA mechanisms, amplifying sedation, respiratory depression, and fall risk. In elderly patients, age-related decreases in alcohol dehydrogenase mean blood alcohol levels are higher per gram of alcohol consumed than in younger adults.
- **Clinical correlation:** Nightly wine is being consumed alongside alprazolam, zolpidem, and PRN oxycodone. This is a high-risk substance interaction that was undisclosed until the daughter reported it.
- **Action:** Non-judgmental counseling on alcohol-drug interactions. Discuss elimination or timing adjustment. Document as CNS depressant in medication reconciliation. Screen for alcohol use disorder using AUDIT-C.
- **Confidence:** HIGH.

SCENARIO 20

Patient Demographics

- **Name:** Calvin Reeves
- **Age:** 61
- **Sex:** Male
- **Setting:** Emergency Department

Chief Complaint

"I was just diagnosed with a blood clot in my leg. The ER doc wants to start me on a blood thinner. I take St. John's Wort and I also take a medication for seizures. I wanted someone to check before they give me anything new."

Medication Sources

Source 1 — Written List (Phone Notes App, Patient Shown)

```
Calvin Reeves – Medications
- Carbamazepine 400 mg twice daily (seizure disorder)
- Lisinopril 10 mg daily
- Atorvastatin 20 mg nightly
- St. John's Wort 300 mg three times daily (self-prescribed)
```

Source 2 — Verbal Report

"I've been taking St. John's Wort for depression for about a year. My neurologist doesn't know about it."

Reconciled Medication List

Medication	Dose	Frequency	Source
Carbamazepine	400 mg	Twice daily	Written list confirmed
Lisinopril	10 mg	Once daily	Written list confirmed
Atorvastatin	20 mg	Once daily (PM)	Written list confirmed
St. John's Wort	300 mg	Three times daily	Written list confirmed — undisclosed to specialist

Reconciliation Flags

- CRITICAL: Carbamazepine is a potent CYP3A4 inducer — will massively reduce levels of many anticoagulants
- CRITICAL: St. John's Wort is also a potent CYP3A4 inducer — combined induction effect is extreme
- DVT requires anticoagulation — but anticoagulant choice is critically dependent on drug interaction profile
- Neurologist unaware of St. John's Wort — adds CYP3A4 induction to seizure drug regimen (may reduce carbamazepine efficacy through autoinduction + SJW interaction)

Top 3 Ranked Interactions

🥇 Rank 1 — Drug–Drug + OTC Supplement Interaction [CRITICAL]

Carbamazepine + St. John's Wort → Potent CYP3A4 Induction → Anticoagulant Failure Risk

- **Mechanism:** Carbamazepine is among the most potent inducers of CYP3A4, CYP2C9, and P-glycoprotein — the primary enzymes metabolizing most direct oral anticoagulants (DOACs: rivaroxaban, apixaban, edoxaban) and warfarin. St. John's Wort (hypericum) is an independently potent CYP3A4 and P-gp inducer. Together, induction is additive or synergistic — plasma levels of co-administered drugs can be reduced by 50–90%.
- **Clinical correlation:** This patient needs anticoagulation for DVT. Standard DOAC doses will likely be sub-therapeutic due to extreme enzyme induction from two concurrent inducers. Under-anticoagulation increases pulmonary embolism risk, which can be fatal.
- **Action:** Do NOT start a standard DOAC regimen without pharmacist and hematology/neurology consultation. Warfarin (with careful INR monitoring) is a possible option (INR titration accounts for induction), but SJW and carbamazepine also affect warfarin metabolism. Parenteral anticoagulation (LMWH/enoxaparin) is less affected by CYP induction and may be the safest bridge. Discontinue St. John's Wort — it also reduces carbamazepine levels and may be destabilizing seizure control. Notify neurologist immediately.
- **Confidence:** CRITICAL — this interaction is documented and potentially life-threatening.

🥈 Rank 2 — Drug + OTC Supplement Interaction [HIGH RELEVANCE]

St. John's Wort → Carbamazepine Level Reduction → Seizure Risk

- **Mechanism:** St. John's Wort induces CYP3A4, which is also involved in carbamazepine metabolism (carbamazepine is a substrate of CYP3A4 as well as an auto-inducer). SJW may paradoxically reduce carbamazepine plasma levels by increasing its metabolism. This is underrecognized because patients self-prescribe SJW without telling their neurologist.
- **Clinical correlation:** Patient has been on St. John's Wort for one year without his neurologist's knowledge. If SJW is reducing carbamazepine levels, seizure control may be suboptimal or unstable.
- **Action:** Carbamazepine drug level (therapeutic range 4–12 mcg/mL). Notify neurologist of SJW use immediately. Discontinue SJW — this also allows accurate reassessment of carbamazepine levels after induction effect resolves (2–4 weeks).
- **Confidence:** HIGH.

🥉 Rank 3 — Drug–Drug Interaction [MODERATE RELEVANCE]

Carbamazepine → Atorvastatin Level Reduction (CYP3A4 Induction)

- **Mechanism:** Atorvastatin is extensively metabolized by CYP3A4. Carbamazepine induction can reduce atorvastatin plasma concentrations by 60–80%, potentially rendering the statin ineffective for cholesterol management.
- **Clinical correlation:** Patient may have inadequate cardiovascular protection from atorvastatin despite being on a standard dose. This is a chronic, ongoing interaction that should be addressed in outpatient follow-up.
- **Action:** Fasting lipid panel to assess current control. Switch to a non-CYP3A4-dependent statin (rosuvastatin or pravastatin) — these are minimally affected by CYP3A4 induction and will provide reliable cardiovascular protection.
- **Confidence:** HIGH for mechanism; MODERATE for immediate urgency (not life-threatening but clinically important for long-term management).

APPENDIX: AI MODEL TRAINING NOTES

Interaction Classification Schema Used in This Dataset

Category	Code	Definition
Drug–Drug	DDI	Two or more medications interact via pharmacokinetic or pharmacodynamic mechanisms
Drug–Symptom	DSI	A medication causes or mimics a clinical symptom (e.g., ACE inhibitor cough)
Drug–Disease	DDSI	A medication is contraindicated or high-risk given a patient's specific diagnosis
Drug–Lab	DLI	A medication affects laboratory values or requires lab monitoring
OTC/Supplement	OTCI	Over-the-counter drugs or supplements interact with prescription medications
Process	PI	Care coordination failures that create medication reconciliation risk

Confidence Level Definitions

Level	Definition
CRITICAL	Immediate patient harm possible; stop/start medication action required now
HIGH	Strong evidence-based interaction; action required this visit
MODERATE	Clinically relevant; requires monitoring and follow-up
LOW-MODERATE	Possible interaction; context-dependent; document and monitor

Geriatric-Specific Flags Used

- **Beers Criteria:** American Geriatrics Society list of medications potentially inappropriate in adults ≥65
- **Anticholinergic Burden:** Cumulative anticholinergic drug load — increases delirium, falls, cognitive impairment
- **Renal dose adjustment:** Most medications require eGFR-based dosing in elderly patients
- **CNS sensitivity:** Elderly patients have increased blood-brain barrier permeability and reduced CNS reserve

Key Reconciliation Data Sources (By Scenario)

Source Type	Scenarios
Handwritten list only	1, 8, 10
Printed PCP/clinic list	2, 7, 17
Typed patient list	3
Verbal only	4 (primary), 5, 9
Discharge paperwork	6
Pill bottles (critical finding)	3 (carvedilol), 6 (furosemide)
OTC/supplement undisclosed	3 (B12), 4 (Tylenol), 5 (scope), 16 (fish oil), 18 (pre-workout), 20 (SJW)
Multi-provider prescribing gap	15, 17, 20
