1

Assignment-1 (12.13.5.5)

Uttam Paharia CS22BTECH11060

12.)Question: Assume that each born child is equally likely to be a boy or a girl. If a family has two children, what is the conditional probability that both are girls given that

- 1) the youngest is a girl
- 2) at least one is a girl?

Solution: Let random variable $X_i = 0$, if it is a girl child and $X_i = 1$ if it is a boy.

Where i = 1 for first child and i = 2 for second child.

It is given that:

$$\Pr(X_i = 1) = \Pr(X_i = 0)$$
 (1)

$$Pr(X_i = 1) + Pr(X_i = 0) = 1$$
 (2)

$$\implies \Pr(X_i = 0) = \Pr(X_i = 1) = 0.5$$
 (3)

We are supposed to find Probability of both girl child(for some given conditions)

$$\implies \Pr(X_1 = 0) \times \Pr(X_2 = 0) \tag{4}$$

This is because gender of each child is independent of another

Variables	Corresponding Child	Probability
$X_1 = 0$	Younger girl child	$\frac{1}{2}$
$X_1 = 1$	Younger boy child	$\frac{1}{2}$
$X_2 = 0$	Elder girl child	$\frac{1}{2}$
$X_2 = 1$	Elder boy child	1/2

TABLE II Variables Declaration

Probabilities	Values
$\Pr\left(X_1 + X_2 = 2\right)$	$\frac{1}{4}$
$\Pr\left(X_1 + X_2 = 1\right)$	$\frac{1}{2}$
$\Pr\left(X_1 + X_2 = 0\right)$	$\frac{1}{4}$

TABLE IV Some useful Probabilities

If the youngest child is girl:
 By conditional Probability we have

$$\Pr((X_1 + X_2 = 0) | X_1 = 0) = \frac{\Pr(X_1 + X_2 = 0)}{\Pr(X_1 = 0)}$$
(5)

$$Pr(X_1 + X_2 = 0) = Pr(X_1 = 0) \times Pr(X_2 = 0)$$
(6)

$$\implies \Pr(X_2 = 0) = 0.5 \tag{7}$$

2) If at least one of the child is girl: By conditional Probability we have

$$\Pr(X_1 + X_2 = 0 | (X_1 + X_2 \le 1)) = \frac{\Pr(X_1 + X_2 = 0)}{\Pr(X_1 + X_2 \le 1)}$$
(8)

$$Pr(X_1 + X_2 \le 1) = 1 - Pr(X_1 + X_2 = 2)$$
 (9)

$$=1-\frac{1}{4}=\frac{3}{4} \qquad (10)$$

Probability of both are girl given that at least one is girl is

From (8)

$$\frac{\frac{1}{4}}{\frac{3}{4}} = \frac{1}{3} \tag{11}$$