

DROWNING OR SUBMERSION

ALL PROVIDERS

- ☐ Focused history and physical exam
 - Blood glucose, core body temperature and oxygen saturation assessment.
 - Assess the scene for other environmental issues or possible toxins.
- ☐ Cardiac monitor, ETCO₂, pulse oximetry monitoring, blood pressure when available.
- ☐ **Treatment Plan**
 - Safely remove patient from the water
 - Place patient supine
 - Remove wet clothing and wrap in blankets
 - Ensure patient warmth
 - If concern for spinal injury refer to *Spinal Motion Restriction Guideline*.
 - Scuba divers “Dive Computer” or Dive Log Book should be transported with the patient.
- ☐ **Key Considerations**
 - Airway maintenance is the primary consideration.
 - Unlike the “CAB” strategy used in standard cardiac arrest, patients suffering cardiac arrest from drowning require an “ABC” approach with emphasis prompt airway management and supplemental ventilations.
 - There can be co-existing conditions depending on the type of submersion injury including trauma, hypothermia, and intoxication.
 - Hypotension is associated with a worse outcome, monitor closely and treat per the *Shock, Sepsis and Fluid Therapy Guideline*, as needed.
 - Initiation of in-water ventilations may increase survival; however, in-water chest compressions are futile.
 - Submersion in cold water will often cause severe hypothermia, notify receiving hospital so that appropriate resources can be mobilized.
 - Pediatric cardiac arrest due to drowning and hypothermia (temperature <30 C/86 F): consider direct transport to Primary Children’s Medical Center for ECMO and do NOT rewarm this patient.
 - Adult cardiac arrest due to drowning and hypothermia (temperature <30 C/86 F): consider direct transport to University of Utah Medical Center or Intermountain Medication Center for ECMO and do NOT rewarm this patient.

ADULT

PEDIATRIC (<15 years of Age)

NOTE: Pediatric weight based dosing should not exceed Adult dosing.

EMT

EMT

- | | |
|--|--|
| <ul style="list-style-type: none"> <input type="checkbox"/> If breathing spontaneously apply oxygen at 15 LPM via non-rebreather mask to maintain oxygen saturations >95% <input type="checkbox"/> Ventilate with BVM when apneic or exhibiting respiratory distress. Consider a nasal or oral airway <input type="checkbox"/> Initiate 5 rescue breaths followed by 30 chest compressions, then use a 30:2 compression: ventilation ratio | <ul style="list-style-type: none"> <input type="checkbox"/> If breathing spontaneously apply oxygen at 15 LPM via non-rebreather mask to maintain oxygen saturations >95% <input type="checkbox"/> Ventilate with BVM when apneic or exhibiting respiratory distress. Consider a nasal or oral airway <input type="checkbox"/> Initiate 5 rescue breaths followed by 30 chest compressions, then use a 15:2 compression: ventilation ratio |
|--|--|

AEMT

- ☐ Advanced airway, vascular access and fluid therapy.
 - Albuterol 2.5 every 10 minutes via nebulization for bronchospasm/wheezing until symptoms subside
 - Reassess patient after each dose to determine need for additional dosing
- ☐ Consider CPAP in awake patients with respiratory distress

PARAMEDIC

- ☐ **Epinephrine 2–10 mcg/min IV/IO** infusion for hypoperfusion. Titrate to maintain a SBP >100 mmHg
- ☐ **Push Dose Epinephrine 10mcg** as needed to maintain a SBP >100 mmHg after fluid bolus
- ☐ **Norepinephrine** initial dose: **0.05 – 1 mcg/kg/min IV/IO** for hypoperfusion. Titrate to maintain a SBP > 100 mmHg. For patients in refractory shock: 8-30 mcg/minute

AEMT

- ☐ Advanced airway, vascular access and fluid therapy.
 - Albuterol 2.5 every 10 minutes via nebulization for bronchospasm/wheezing until symptoms subside. Start with 1.25 mg if age <1yr
 - Reassess patient after each dose to determine need for additional dosing
- ☐ Consider CPAP in awake patients with respiratory distress

PARAMEDIC

- ⌚ **Epinephrine 0.1–1 mcg/kg/min IV/IO** infusion for hypoperfusion. Titrate to maintain a SBP >70 + (age in years x 2) mmHg
- ⌚ **Push Dose Epinephrine (dose per appendix)** as needed to maintain a SBP >70 + (age in years x 2) mmHg after fluid bolus
- ⌚ **Norepinephrine** initial dose: **0.05 - 0.1 mcg/kg/min**, titrate to max of 2 mcg/kg/min to maintain SBP >70 + (age in years x 2) mmHg