

GLUCOSE EMERGENCIES

HYPOGLYCEMIA / HYPERGLYCEMIA

ALL PROVIDERS

- ☐ Focused history and physical exam
 - Blood glucose assessment (heel stick is preferred in newborns or infants).
 - Hypoglycemia is defined as blood glucose level <50 mg/dl for adults, <60 mg/dl for children, and <40 mg/dl for the term neonate (<30days of age) with any degree of altered mentation.
- ☐ **Treatment Plan**
 - Insulin pump in place: Hypoglycemic patient with altered mentation -
 - Care is directed at treating hypoglycemia first, then stopping administration of insulin.
 - Turn off insulin pump, if able.
 - If no one familiar with the device is available to assist, disconnect pump from patient by either:
 - Using quick-release where the tubing enters the dressing on patient's skin.
 - Completely remove the dressing, thereby removing the subcutaneous needle and catheter from under patient's skin.
 - When mental status returns to normal, patient should be strongly encouraged to eat.
 - Criteria for scene release of hypoglycemic patient:
 - Patient does not want to be transported.
 - Return to apparent normal mental capacity following treatment.
 - Insulin only. The patient does not have access to oral medications for diabetes.
 - No suicidal ideations or recent suicide attempt.
 - There is at least one responsible party that can assist them in their recovery and is comfortable in their care.
 - Children should be considered for transport for evaluation regardless of improvement in the field due to other possible etiologies for the episode.
- ☐ **Key Considerations**
 - Do NOT attempt to give oral glucose to those who are unconscious, cannot swallow or whose gag reflex is diminished.
 - Transport any patient who is at risk for prolonged or recurrent hypoglycemia such as long acting insulin or oral hypoglycemic overdose.
 - If the patient is hypoglycemic and has a seizure, recheck blood glucose every 15 minutes to check for recurrent low blood sugar that may need treatment.

ADULT

PEDIATRIC (<15 years of Age)

NOTE: Pediatric weight based dosing should not exceed Adult dosing.

EMT

- ☐ **Dextrose Oral glucose 15 grams** if patient is able to protect airway
 - Repeat in 15 minutes as needed

EMT

- ☐ **Dextrose Oral glucose 7.5 grams** if patient is able to protect airway
 - Repeat in 15 minutes as needed

AEMT

- ☐ Vascular access and fluid therapy

AEMT

- ☐ Vascular access and fluid therapy

HYPOGLYCEMIA

- ☐ **Dextrose 50% 12.5 grams** (25mL) IV/IO. May repeat as necessary
- ☐ **Dextrose 10%:** Infuse **125 mL** (12.5 grams), then recheck blood sugar. If still low, may repeat
- ☐ **Glucagon 1 mg** IM if no IV/IO access available

HYPOGLYCEMIA

- ☐ Infants up to 1 year
 - **Dextrose 10% 5 mL/kg** (0.5 grams/kg) IV/IO. May repeat as necessary up to a MAX of 125 mL (12.5 grams).
- ☐ Children greater than 1 year

- HYPERGLYCEMIA (BS >300 mg/dL)**
- ☐ **Normal Saline 1000 mL IV/IO** over 30–60 minutes

- **Dextrose 25% 2 mL/kg IV/IO:** repeat as necessary (max 12.5G/ 50mL)
- **Dextrose 10% 5 mL/kg (0.5 grams/kg)** IV/IO. May repeat as necessary up to a MAX of 125 mL (12.5 grams).
- **Glucagon 0.01 mg/kg (max dose of 1 mg)** IM if no IV/IO access available

- HYPERGLYCEMIA (BS >300 mg/dL)**
- ☐ **Normal Saline 20 mL/kg IV/IO** over 30–60 minutes for hyperglycemic patient

PARAMEDIC

PARAMEDIC