Compiled using LATEX 1

STAT 430-1, Fall 2024 HW2

Tuorui Peng¹

Contents

Atoms

Fatou's Lemma	1
Dominated convergence theorem	2
On interchanging limits and integration	2
	Dominated convergence theorem

1

 $\mathbf{2}$

Exercise 1 Atoms

On interchanging integrals

Denote by K_n the set of atoms with mass greater than 1/n, then we have

$$\mu(E) \ge \mu(K_n) \ge \frac{|K_n|}{n} \Rightarrow |K_n| \le n\mu(E)$$

which means that K_n is finite for any $n \in \mathbb{N}^+$. And we also note that $\{K_n\}_{n \in \mathbb{N}^+}$ is an increasing sequence of sets. By definition of stones, we have the set of all stones being $S = \bigcup_{n=1}^{\infty} K_n = \bigoplus_{n=1}^{\infty} K_n \setminus K_{n-1}$. i.e. $\{K_n \setminus K_{n-1}\}_{n \in \mathbb{N}^+}$ is a partition of S.

If S is uncountable, then there exists $n_0 \in \mathbb{N}^+$ such that $K_{n_0} \setminus K_{n_0-1}$ is uncountable, an thus $K_{n_0} = \bigcup_{k=1}^{n_0} K_{n_0} \setminus K_{n_0-1}$ is also uncountable, which contradicts the fact that $|K_{n_0}| \leq n\mu(E) < \infty$.

Thus S is countable.

Exercise 2 Fatou's Lemma

Note that $\{\inf_{m\geq n} f_m\}_{n\in\mathbb{N}}$ is a monotone increasing sequence of functions. Using monotone convergence theorem, we have

$$\mu\left(\liminf_{n\to\infty} f_n\right) = \mu\left(\liminf_{n\to\infty} \inf_{m\geq n} f_m\right)$$

$$= \lim_{n\to\infty} \mu\left(\inf_{m\geq n} f_m\right)$$

$$\leq \lim_{n\to\infty} \inf_{m\geq n} \mu(f_m)$$

$$= \lim_{n\to\infty} \inf \mu(f_n)$$

 $^{^1}$ TuoruiPeng2028@u.northwestern.edu

Tuorui Peng 2

Exercise 3 Dominated convergence theorem

From above, we already have $\mu(f) \leq \liminf_{n \to \infty} \mu(f_n)$. Now it suffice to show the other direction.

Note that we have dominance condition $-g \leq f_n \leq g$, $\forall n \in \mathbb{N}^+$, thus we can consider the following sequence of functions: $\{g - f_n\}_{n \in \mathbb{N}^+}$, which has

$$0 \le g - f_n \le 2g$$

thus is still integrable. By Fatou's lemma, we have

$$\mu(g-f) \le \liminf_{n \to \infty} \mu(g-f_n) \Rightarrow \mu(f) \ge \limsup_{n \to \infty} \mu(f_n)$$

which concludes the proof.

Exercise 4 On interchanging limits and integration

• Left hand side: we have

$$\lim_{\lambda \to 0} \int_0^\infty f_{\lambda}(x) \, \mathrm{d}x = \lim_{\lambda \to 0} \int_0^\infty \lambda e^{-\lambda x} \, \mathrm{d}x = \lim_{\lambda \to 0} 1 = 1$$

• Right hand side: we have

$$\int_0^\infty \lim_{\lambda \to 0} f_{\lambda}(x) \, \mathrm{d}x = \int_0^\infty \lim_{\lambda \to 0} \lambda e^{-\lambda x} \, \mathrm{d}x = \int_0^\infty 0 \, \mathrm{d}x = 0$$

Thus the two sides are not equal, and the limit cannot be interchanged with the integral.

To explain why this happens, we illustrate the following that there is no dominating function for the sequence of functions $\{f_{\lambda}(x)\}_{\lambda>0}$: for any $x \in \mathbb{R}^+$, and any $\varepsilon \in (0,1)$, we can find λ_{ε} s.t. $\lambda_{\varepsilon}e^{-\lambda_{\varepsilon}x} = \varepsilon$, thus to dominate the sequence, we need to have $g(x) \geq \varepsilon$, $\forall x \in \mathbb{R}^+$, which makes g(x) un-integrable. Thus there is no dominating function for the sequence of functions, and the condition of Lebesgue's dominated convergence theorem is not satisfied.

Exercise 5 On interchanging integrals

• We have

$$I = \int_0^\infty dx \int_0^\infty dy f(x, y) = \int_0^\infty dx \begin{cases} x - 1, & x \in [0, 1] \\ 0, & x \in (1, \infty) \end{cases}$$
$$= \int_0^1 dx (x - 1) = -\frac{1}{2}$$

• We have

$$J = \int_0^\infty dy \int_0^\infty dx f(x, y) = \int_0^\infty dy \begin{cases} 1 - y, & y \in [0, 1] \\ 0, & y \in (1, \infty) \end{cases}$$
$$= \int_0^1 dy (1 - y) = \frac{1}{2}$$

Tuorui Peng 3

i.e. we got $I \neq J$, and the integrals cannot be interchanged. We can explain this by noticing that the condition of Fubini's theorem is not satisfied: the function f(x,y) is not integrable on \mathbb{R}^2 :

$$\int_0^\infty dx \int_0^\infty dy |f(x,y)| = \infty.$$