

A Brief Summary of Statistics Course

统计学课程知识总结

Vincent

2021 年 1 月 13 日

目录

1 概率论部分	3
1.1 Some Important Distributions	3
1.2 Probability and Probability Model	3
1.2.1 Sample and σ -Field	3
1.2.2 Axioms of Probability	4
1.2.3 Conditional Probability	5
1.3 Properties of Random Variable and Vector	5
1.3.1 Random Variable	5
1.3.2 Random Vector	6
1.4 Properties of E , σ^2 and cov	7
1.4.1 Expection	7
1.4.2 Variance	8
1.4.3 Covariance and Correlation	8
1.5 PGF, MGF and C.F	9
1.5.1 Probability Generating Function	9
1.5.2 Moment Generating Function	10
1.5.3 Characteristic Function	10
1.6 Convergence and Limit Distribution	10
1.6.1 Convergence Mode	10
1.6.2 Law of Large Number & Central Limit Theorem	11
1.7 Inequalities	12
1.8 Multivariate Normal Distribution	12
1.8.1 Linear Transform	12

1.8.2	Distributions of Function of Normal Variable: χ^2 , t & F	13
2	统计推断部分	16
2.1	Statistical Model and Statistics	16
2.1.1	Statistics	16
2.1.2	Exponential Family	17
2.1.3	Sufficient and Complete Statistics	18
2.2	Point Estimation	19
2.2.1	Optimal Criterion	19
2.2.2	Method of Moments	20
2.2.3	Maximum Likelihood Estimation	21
2.2.4	Uniformly Minimum Variance Unbiased Estimate	22
2.2.5	MoM and MLE in Linear Regression	24
2.2.6	Kernel Density Estimate	26
2.3	Interval Estimation	27
2.3.1	Confidence Interval	27
2.3.2	Pivot Variable Method	28
2.3.3	Confidence Interval for Common Distributions	28
2.3.4	Fisher Fiducial Argument*	30
2.4	Hypothesis Testing	30
2.4.1	Basic Concepts	31
2.4.2	Hypothesis Testing of Common Distributions	32
2.4.3	Likelihood Ratio Test	34
2.4.4	Uniformly Most Powerful Test	34
2.4.5	Duality of Hypothesis Testing and Interval Estimation	35
2.4.6	Introduction to Non-Parametric Hypothesis Testing	36

1 概率论部分

Cover: Basic axioms, random events, σ -field; random variable/vector and their properties, some special distributions; E & σ^2 & cov and their properties; probability-generating/moment-generating/characteristic function; weak/strong law of large number, central limit thm.; intro. to multivariate normal distribution.

1.1 Some Important Distributions

X	$p_X(k)/f_X(x)$	E	σ^2	PGF	MGF
$B(p)$		p	pq		$q + pe^s$
$B(n, p)$	$C_n^k p^k (1-p)^{n-k}$	np	npq	$(q + ps)^n$	$(q + pe^s)^n$
$G(p)$	$(1-p)^{k-1} p$	$\frac{1}{p}$	$\frac{q}{p^2}$	$\frac{ps}{1-qs}$	$\frac{pe^s}{1-qe^s}$
$H(n, M, N)$	$\frac{C_M^k C_{N-M}^{n-k}}{C_N^n}$	$n \frac{M}{N}$	$\frac{nM(N-n)(N-M)}{N^2(n-1)}$		
$P(\lambda)$	$\frac{\lambda^k}{k!} e^{-\lambda}$	λ	λ	$e^{\lambda(s-1)}$	$e^{\lambda(e^s-1)}$
$U(a, b)$	$\frac{1}{b-a}$	$\frac{a+b}{2}$	$\frac{(b-a)^2}{12}$		$\frac{e^{sb} - e^{sa}}{(b-a)s}$
$N(\mu, \sigma^2)$	$\frac{1}{\sigma\sqrt{2\pi}} e^{-\frac{(x-\mu)^2}{2\sigma^2}}$	μ	σ^2		$e^{\frac{\sigma^2 s^2}{2} + \mu s}$
$\epsilon(\lambda)$	$\lambda e^{-\lambda x}$	$\frac{1}{\lambda}$	$\frac{1}{\lambda^2}$		$\frac{\lambda}{\lambda-s}$
$\Gamma(\alpha, \lambda)$	$\frac{\lambda^\alpha}{\Gamma(\alpha)} x^{\alpha-1} e^{-\lambda x}$	$\frac{\alpha}{\lambda}$	$\frac{\alpha}{\lambda^2}$		
$B(\alpha, \beta)$	$\frac{1}{B(\alpha, \beta)} x^{\alpha-1} (1-x)^{\beta-1}$	$\frac{\alpha}{\alpha+\beta}$	$\frac{\alpha\beta}{(\alpha+\beta)^2(\alpha+\beta+1)}$		
χ_n^2	$\frac{1}{2^{\frac{n}{2}} \Gamma(\frac{n}{2})} x^{\frac{n}{2}-1} e^{-\frac{x}{2}}$	n	$2n$		
t_ν	$\frac{\Gamma(\frac{\nu+1}{2})}{\sqrt{\nu\pi} \Gamma(\frac{\nu}{2})} (1 + \frac{x^2}{\nu})^{-\frac{\nu+1}{2}}$	0	$\frac{\nu}{\nu-2}$		
$F(m, n)$	$\frac{\Gamma(\frac{m+n}{2})}{\Gamma(\frac{m}{2}) \Gamma(\frac{n}{2})} \frac{m^{\frac{m}{2}} x^{\frac{m}{2}-1}}{(mx+n)^{\frac{m+n}{2}}}$	$\frac{n}{n-2}$	$\frac{2n^2(m+n-2)}{m(n-2)^2(n-4)}$		

More Properties of χ^2, t, F see section 1.8.2.

Definition of PGF, MGF, CF see section 1.5.

1.2 Probability and Probability Model

What is **Probability**? A 'belief' in the chance of an event occurring.

1.2.1 Sample and σ -Field

Def. sample space Ω : The set of all possible outcomes of one particular experiment.

Def. \mathcal{F} a σ -field(or a σ -algebra) as a collection of some subsets of Ω if

- $\Omega \in \mathcal{F}$

- if $A \in \mathcal{F}$, then $A^C \in \mathcal{F}$
- if $A_n \in \mathcal{F}$, then $\bigcup_{n=1}^{\infty} A_n \in \mathcal{F}$

And (Ω, \mathcal{F}) is a measurable space.

1.2.2 Axioms of Probability

P is probability measure (or probability function) defined on (Ω, \mathcal{F}) , satisfying

- Nonnegativity

$$P(A) \geq 0 \quad \forall A \in \Omega$$

- Normalization

$$P(\Omega) = 1$$

- Countable Additivity

$$P(A_1 \cup A_2 \cup \cdots) = P(A_1) + P(A_2) + \cdots \quad (A_i \parallel A_j \quad \forall i \neq j)$$

Then (Ω, \mathcal{F}, P) is probability space.

Properties of Probability:

- Monotonicity

$$P(A) \leq P(B) \quad \text{for } A \subset B$$

- Finite Subadditivity (Boole Inequality)

$$P\left(\bigcup_{i=1}^n A_i\right) \leq \sum_{i=1}^n P(A_i)$$

- Inclusion-Exclusion Formula

$$\begin{aligned} P(\mathbb{U}_{i=1}^n A_i) &= \sum_{1 \leq i \leq n} P(A_i) - \sum_{1 \leq i < j \leq n} P(A_i \cap A_j) \\ &+ \sum_{1 \leq i < j < k \leq n} P(A_i \cap A_j \cap A_k) - \cdots \\ &+ (-1)^{n-1} P(A_1 \cap A_2 \cap \cdots \cap A_n) \end{aligned}$$

- Borel-Cantelli Lemma

$$\begin{aligned} \sum_{n=1}^{\infty} P(A_n) < \infty &\Rightarrow P(\limsup_{n \rightarrow \infty} A_n) = 0 \\ \sum_{n=1}^{\infty} P(A_n) = \infty &\Rightarrow P(\limsup_{n \rightarrow \infty} A_n) = 1 \quad \text{if } A_i \text{ independent} \end{aligned}$$

1.2.3 Conditional Probability

Def. **Conditional Probability** of B given A :

$$P(B|A) = \frac{P(A \cap B)}{P(A)}$$

(Actually a change of σ -field from Ω to B)

Application of conditional probability:

- Multiplication Formula

$$P(\cap_{i=1}^n A_i) = P(A_1) \prod_{i=2}^n P(A_i | A_1 \cap A_2 \cap \cdots \cap A_{i-1})$$

- Total Probability Thm

$$P(B) = \sum_{i=1}^n P(A_i)P(B|A_i)$$

where $\{A_i\}$ is a partition of Ω .

- Bayes's Rule

$$P(A_i|B) = \frac{P(A_i)P(B|A_i)}{\sum_{j=1}^n P(A_j)P(B|A_j)}$$

where $\{A_i\}$ is a partition of Ω .

- Statistically Independence

$$P(A \cap B) = P(A)P(B), \text{ for } A \parallel B$$

1.3 Properties of Random Variable and Vector

1.3.1 Random Variable

Def. Random Variable: a **function** X defined on sample space Ω , mapping from Ω to some $\mathcal{X} \in \mathbb{R}$.

Then def. Cumulative Distribution Function (CDF).

$$F_X(x) = P(X \leq x)$$

For Discrete case, consider CDF as right-continuity.

- PMF:

$$p_X(x) = F_X(x^+) - F_X(x^-)$$

PDF

$$f_X(x) = \frac{dF_X(x)}{dx}$$

- Indicator function:

$$I_{x \in A}(x) = \begin{cases} 1 & x \in A \\ 0 & x \notin A \end{cases}$$

- Convolution

$$- W = X + Y$$

$$f_W(w) = \int_{-\infty}^{\infty} f_X(x)f_Y(w-x)dx$$

$$- V = X - Y$$

$$f_V(v) = \int_{-\infty}^{\infty} f_X(x)f_Y(x-v)dx$$

$$- Z = XY$$

$$f_Z(z) = \int_{-\infty}^{\infty} \frac{1}{|x|} f_X(x)f_Y\left(\frac{z}{x}\right)dx$$

- Order Statistics

Def $X_{(1)}, X_{(2)}, \dots, X_{(n)}$ as order statistics of \vec{X}

$$g_{X_{(i)}} = n! \prod f(x_i) \quad \text{for } x_1 < x_2 < \dots < x_n$$

PDF of $X_{(k)}$

$$g_k(x_k) = nC_{n-1}^{k-1}[F(x_k)]^{k-1}[1-F(x_k)]^{n-k}f(x_k)$$

- p -fractile

$$\xi_p = F^{-1}(p) = \inf\{x | F(x) \geq p\}$$

1.3.2 Random Vector

A general case of random variable.

n -dimension Random Vector $\vec{X} = (X_1, X_2, \dots, X_n)$ defined on (Ω, \mathcal{F}, P) .

CDF $F(x_1, \dots, x_n)$ defined on \mathbb{R}^n :

$$F(x_1, \dots, x_n) = P(X_1 \leq x_1, \dots, X_n \leq x_n)$$

Joint PDF of random vector:

$$f(x_1, \dots, x_n) = \frac{\partial^n F(x_1, \dots, x_n)}{\partial x_1 \dots \partial x_n}$$

k -dimensional Marginal Distribution: For $1 \leq k < n$ and index set $S_k = \{i_1, \dots, i_k\}$, distribution of $\vec{X} = (X_{i_1}, X_{i_2}, \dots, X_{i_k})$

$$F_{S_k}(x_{i_1}, X_{i_2} \leq x_{i_2}, \dots, x_{i_k}) = P(X_{i_1} \leq x_{i_1}, \dots, X_{i_k} \leq x_{i_k}; X_{i_{k+1}}, \dots, X_{i_n} \leq \infty)$$

Marginal distribution:

$$g_{S_k}(x_{i_1}, \dots, x_{i_k}) = \int_{\mathbb{R}^{n-k}} f(x_1, \dots, x_n) dx_{i_{k+1}} \dots dx_{i_n} = \frac{\partial^{n-k} F(x_1, \dots, x_n)}{\partial x_{i_{k+1}} \dots \partial x_{i_n}}$$

Δ Function of r.v.

For $\vec{X} = (X_1, X_2, \dots, X_n)$ with PDF $f(\vec{X})$ and define

$$\vec{Y} = (Y_1, Y_2, \dots, Y_n) = (y_1(\vec{X}), y_2(\vec{X}), \dots, y_n(\vec{X}))$$

with inverse mapping

$$\vec{X} = (X_1, X_2, \dots, X_n) = (x_1(\vec{Y}), x_2(\vec{Y}), \dots, x_n(\vec{Y}))$$

then

$$g(\vec{Y}) = f(x_1(\vec{Y}), x_2(\vec{Y}), \dots, x_n(\vec{Y})) \left| \frac{\partial \vec{X}}{\partial \vec{Y}} \right| I_{D_Y}$$

(Intuitively: $g(\vec{Y})d\vec{Y} = dP = f(\vec{X})d\vec{X}$)

1.4 Properties of E , σ^2 and cov

Expectation and Variance of common distributions see sec.1.1.

1.4.1 Expection

Expectation of r.v. $g(X)$ def.:

$$E[g(X)] = \begin{cases} \int_{\Omega} g(x)f_X(x)dx = \int_{\Omega} g(x)dF(x) \\ \sum_{\Omega} g(X)f_X(x) \end{cases}$$

Properties of expectation $E(\cdot)$:

- Linearity of Expectation

$$E(aX + bY) = aE(X) + bE(Y)$$

- Conditional Expectation

$$E(X|A) = \frac{E(XI_A)}{P(A)}$$

Note: if take A as Y is also a r.v. then

$$m(Y) = E(X|Y) = \int x f_{X|Y}(x)dx$$

is actually a function of Y

- Law of Total Expectation

$$E\{E[g(X)|Y]\} = E[g(X)]$$

- r.v.& Event

$$P(A|X) = E(I_A|X) \Rightarrow E[P(A|X)] = E(I_A) = P(A)$$

-

$$E[h(Y)g(X)|Y] = h(Y)E[g(X)|Y]$$

1.4.2 Variance

Variance of r.v. X :

$$\text{var}(X) = E[(X - E(X))^2] = E(X^2) - (E(X))^2$$

(sometimes denoted as σ_X^2 .)

Properties:

- Linear combination of Variance

$$\text{var}(aX + b) = a^2 \text{var}(X)$$

- Conditional Variance

$$\text{var}(X|Y) = E[X - E(X|Y)]^2|Y$$

- Law of Total Variance

$$\text{var}(X) = E[\text{var}(X|Y)] + \text{var}[E(X|Y)]$$

Standard Deviation def. as :

$$\sigma_X = \sqrt{\text{var}(X)}$$

Then can construct **Standardization** of r.v.

$$Y + \frac{X - E(X)}{\sqrt{\text{var}(X)}}$$

1.4.3 Covariance and Correlation

Covariance of r.v. X and Y :

$$\text{cov}(X, Y) = E[(X - \mu_X)(Y - \mu_Y)] = E(XY) - E(X)E(Y)$$

And (Pearson's) Correlation Coefficient

$$\rho_{X,Y} = \text{corr}(X, Y) = \frac{\text{cov}(X, Y)}{\sqrt{\text{var}(X)\text{var}(Y)}}$$

Remark: correlation \nRightarrow cause and effect.

Properties:

- Bilinear of Covariance

$$\text{cov}(X + Y, Z) = \text{cov}(X, Z) + \text{cov}(Y, Z)$$

$$\text{cov}(X, Y + Z) = \text{cov}(X, Y) + \text{cov}(X, Z)$$

- Variance and Covariance

$$\text{var}(X + Y) = \text{var}(X) + \text{var}(Y) + 2\text{cov}(X, Y)$$

- Covariance Matrix

Def $\Sigma = E[(X - \mu)^T(X - \mu)] = \{\sigma_{ij}\}$

$$\Sigma = \begin{pmatrix} \text{var}(X_1) & \text{cov}(X_1, X_2) & \dots & \text{cov}(X_1, X_n) \\ \text{cov}(X_2, X_1) & \text{var}(X_2) & \dots & \text{cov}(X_2, X_n) \\ \vdots & \vdots & \ddots & \vdots \\ \text{cov}(X_n, X_1) & \text{cov}(X_n, X_2) & \dots & \text{var}(X_n) \end{pmatrix} \quad (1.1)$$

Attachment: Independence:

$$X_i || X_j \Rightarrow \begin{cases} f(x_1, x_2, \dots, x_n) = \prod f(x_i) \\ F(x_1, x_2, \dots, x_n) = \prod F(x_i) \\ E(\prod X_i) = \prod E(X_i) \\ \text{var}(\sum X_i) = \sum \text{var}(X_i) \end{cases}$$

1.5 PGF, MGF and C.F

Generating Function: Representation of P in function space. $P \Leftrightarrow$ Generating Function.

1.5.1 Probability Generating Function

PGF: used for non-negative, integer X

$$g(s) = E(s^X) = \sum_{j=0}^{\infty} s^j P(X = j), s \in [-1, 1]$$

Properties

- $P(X = k) = \frac{g^{(k)}(0)}{k!}$
- $E(X) = g^{(1)}(1)$
- $\text{var}(X) = g^{(2)}(1) + g^{(1)}(1) - [g^{(1)}(1)]^2$
- For X_1, X_2, \dots, X_n independent with $g_i(s) = E(s^{X_i})$, $Y = \sum_{i=1}^n X_i$, then

$$g_Y(s) = \prod_{i=1}^n g_i(s), s \in [-1, 1]$$

- For X_i i.i.d with $\psi(s) = E(s^{X_i})$, Y with $G(s) = E(s^Y)$, $W = X_1 + X_2 + \dots + X_Y$, then

$$g_W(s) = G[\psi(s)]$$

- 2-Dimensional PGF of (X, Y)

$$g(s, t) = E(s^X t^Y) = \sum_{i=0}^{\infty} \sum_{j=0}^{\infty} P_{(X,Y)}(X = i, Y = j) s^i t^j, s, t \in [-1, 1]$$

1.5.2 Moment Generating Function

MGF:

$$M_X(s) = E(e^{sX}) = \begin{cases} \sum_j e^{sx} P(X = x_j) \\ \int_{-\infty}^{\infty} e^{sx} f_X(x) dx \end{cases}$$

Properties

- MGF of $Y = aX + b$: $M_Y(s) = e^{sb} M(sa)$
- $E(X^k) = M^{(k)}(0)$
- $P(X = 0) = \lim_{s \rightarrow -\infty} M(s)$
- For X_1, X_2, \dots, X_n independent with $M_{X_i}(s) = E(e^{sX_i})$, $Y = \sum_{i=1}^n X_i$, then

$$M_Y(s) = \prod_{i=1}^n M_{X_i}(s)$$

1.5.3 Characteristic Function

C.F is actually the Fourier Transform of f .

$$\phi(t) = E(e^{itX}) = \int_{-\infty}^{\infty} e^{itx} f_X(x) dx$$

Properties

- if $E(|X|^k) < \infty$, then
- $$\phi^{(k)}(t) = i^k E(X^k e^{itX}) \quad \phi^{(k)}(0) = i^k E(X^k)$$
- For X_1, X_2, \dots, X_n independent with $\phi_{X_i}(t) = E(e^{itX_i})$, $Y = \sum_{i=1}^n X_i$, then

$$\phi_Y(t) = \prod_{i=1}^n \phi_{X_i}(t)$$

- Inverse (Fourier) Transform


$$f(x) = \frac{1}{2\pi} \int_{-\infty}^{\infty} e^{-itx} \phi(t) dt$$

1.6 Convergence and Limit Distribution

1.6.1 Convergence Mode

$$\left\{ \begin{array}{ll} \text{Convergence in Distribution} & X_n \xrightarrow{\mathcal{L}} X : \lim_{n \rightarrow \infty} F_n(x) = F(x) \\ \text{Convergence in Probability} & X_n \xrightarrow{p} X : \lim_{n \rightarrow \infty} P(|X_n - X| \geq \varepsilon) = 0, \forall \varepsilon > 0 \\ \text{Almost Sure Convergence} & X_n \xrightarrow{\text{a.s.}} X : P(\lim_{n \rightarrow \infty} X_n = X) = 1 \\ L_p \text{ Convergence} & X_n \xrightarrow{L_p} X : \lim_{n \rightarrow \infty} E(|X_n - X|^p) = 0 \end{array} \right.$$

Relations between convergence:



Useful Thm.:

- Continuous Mapping Thm.: For continuous function $g(\cdot)$

1. $X_n \xrightarrow{a.s.} X \Rightarrow g(X_n) \xrightarrow{a.s.} g(X)$
2. $X_n \xrightarrow{p} X \Rightarrow g(X_n) \xrightarrow{p} g(X)$
3. $X_n \xrightarrow{\mathcal{L}} X \Rightarrow g(X_n) \xrightarrow{\mathcal{L}} g(X)$

- Slutsky's Thm.: For $X_n \xrightarrow{\mathcal{L}} X, Y_n \xrightarrow{p} c$

1. $X_n + Y_n \xrightarrow{\mathcal{L}} X + c$
2. $X_n Y_n \xrightarrow{\mathcal{L}} cX$
3. $X_n / Y_n \xrightarrow{\mathcal{L}} X / c$

- Continuity Thm.

$$\lim_{n \rightarrow \infty} \phi_n(t) = \varphi(t) \Leftrightarrow X_n \xrightarrow{\mathcal{L}} X$$

1.6.2 Law of Large Number & Central Limit Theorem

- WLLN

$$\frac{1}{n} \sum X_i \xrightarrow{p} E(X_1)$$

- SLLN

$$\frac{1}{n} \sum X_i \xrightarrow{a.s.} C$$

- CLT

$$\frac{1}{\sigma\sqrt{n}} \sum (X_k - \mu) \xrightarrow{\mathcal{L}} N(0, 1)$$

- de Moivre-Laplace Thm.

$$P(k \leq S_n \leq m) \approx \Phi\left(\frac{m + 0.5 - np}{\sqrt{npq}}\right) - \Phi\left(\frac{k - 0.5 - np}{\sqrt{npq}}\right)$$

- Stirling Eqa

$$\frac{\lambda^k}{k!} e^{-\lambda} \approx \frac{1}{\sqrt{\lambda} \sqrt{2\pi}} e^{-\frac{(k-\lambda)^2}{2\lambda}} \xrightarrow[k=\lambda]{k=n} n! \approx \sqrt{2\pi n} \left(\frac{n}{e}\right)^n$$

1.7 Inequalities

- Cauchy-Schwarz Inequality

$$|E(XY)| \leq \sqrt{E(X^2)E(Y^2)}$$

- Bonferroni Inequality

$$P\left(\bigcup_{i=1}^n A_i\right) \geq \sum_{1 \leq i \leq n} P(A_i) + \sum_{1 \leq i < j \leq n} P(A_i \cap A_j)$$

- Markov Inequality

$$P(|X| \geq \epsilon) \leq \frac{E(|X|^\alpha)}{\epsilon^\alpha}$$

- Chebyshev Inequality

$$P(|X - E(X)| \geq \epsilon) \leq \frac{\text{var}(X)}{\epsilon^2}$$

- Jensen Inequality: For convex function $g(x)$:

$$E[g(X)] \geq g(E(X))$$

1.8 Multivariate Normal Distribution

For X_1, X_2, \dots, X_n independent and $X_k \sim N(\mu_k, \sigma_k^2)$, $k = 1, \dots, n$, $T = \sum_{k=1}^n c_k X_k$, (c_k const), then

$$T \sim N\left(\sum_{k=1}^n c_k \mu_k, \sum_{k=1}^n c_k^2 \sigma_k^2\right)$$

Deduction in some special cases:

- Given $\mu_1 = \mu_2 = \dots = \mu_n = \mu$, $\sigma_1^2 = \sigma_2^2 = \dots = \sigma_n^2 = \sigma^2$, i.e. X_k i.i.d., then

$$T \sim N\left(\mu \sum_{k=1}^n c_k, \sigma^2 \sum_{k=1}^n c_k^2\right)$$

- Further take $c_1 = c_2 = \dots = c_n = \frac{1}{n}$, i.e. $T = \sum_{k=1}^n X_k/n = \bar{X}$, then

$$T = \bar{X} \sim N\left(\mu, \frac{\sigma^2}{n}\right)$$

1.8.1 Linear Transform

First consider $\epsilon_1, \epsilon_2, \dots, \epsilon_m$ i.i.d. $\sim N(0, 1)$, $n \times 1$ const column vector $\vec{\mu}$, $n \times m$ const matrix $\mathbf{B} = \{b_{ij}\}$,
 def. $X_i = \sum_{j=1}^m b_{ij} \epsilon_j$, i.e.

$$\vec{X} = \begin{pmatrix} X_1 \\ X_2 \\ \vdots \\ X_n \end{pmatrix} = \begin{pmatrix} b_{11} & b_{12} & \dots & b_{1m} \\ b_{21} & b_{22} & \dots & b_{2m} \\ \vdots & \vdots & \ddots & \vdots \\ b_{n1} & b_{n2} & \dots & b_{nm} \end{pmatrix} \begin{pmatrix} \epsilon_1 \\ \epsilon_2 \\ \vdots \\ \epsilon_m \end{pmatrix}$$

We have: $\vec{X} \sim N(\vec{\mu}, \Sigma)$, where Σ , as defined in eqa.1.1 is

$$\Sigma = \mathbf{B}\mathbf{B}^T = \begin{pmatrix} \text{var}(X_1) & \text{cov}(X_1, X_2) & \dots & \text{cov}(X_1, X_n) \\ \text{cov}(X_2, X_1) & \text{var}(X_2) & \dots & \text{cov}(X_2, X_n) \\ \vdots & \vdots & \ddots & \vdots \\ \text{cov}(X_n, X_1) & \text{cov}(X_n, X_2) & \dots & \text{var}(X_n) \end{pmatrix} = \{\sigma_{ij}\}$$

Furthur Consider $\vec{Y} = (Y_1, \dots, Y_n)^T$, $n \times n$ const square matrix $\mathbf{A} = \{a_{ij}\}$ and def. $\vec{Y} = \mathbf{A}\vec{X}$ i.e.

$$\begin{pmatrix} Y_1 \\ Y_2 \\ \vdots \\ Y_n \end{pmatrix} = \begin{pmatrix} a_{11} & a_{12} & \dots & a_{1n} \\ a_{21} & a_{22} & \dots & a_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ a_{n1} & a_{n2} & \dots & a_{nn} \end{pmatrix} \begin{pmatrix} X_1 \\ X_2 \\ \vdots \\ X_n \end{pmatrix}$$

Then $\vec{Y} \sim N(\mathbf{A}\vec{\mu}, \mathbf{A}\Sigma\mathbf{A}^T)$

Special case: X_1, \dots, X_n i.i.d. $\sim N(\mu, \sigma^2)$, $\vec{X} = (X_1, \dots, X_n)^T$,

$$\begin{aligned} E(Y_i) &= \mu \sum_{k=1}^n a_{ik} \\ \text{var}(Y_i) &= \sigma^2 \sum_{k=1}^n a_{ik}^2 \\ \text{cov}(Y_i, Y_j) &= \sigma^2 \sum_{k=1}^n a_{ik} a_{jk} \end{aligned}$$

Specially when $\mathbf{A} = \{a_{ij}\}$ orthonormal, we have Y_1, \dots, Y_n independent

$$Y_i \sim N\left(\mu \sum_{k=1}^n a_{ik}, \sigma^2\right)$$

1.8.2 Distributions of Function of Normal Variable: χ^2 , t & F

Consider X_1, X_2, \dots, X_n i.i.d. $\sim N(0, 1)$; Y, Y_1, Y_2, \dots, Y_m i.i.d. $\sim N(0, 1)$

- χ^2 Distribution: Def. χ^2 distribution with degree of freedom n :

$$\xi = \sum_{i=1}^n X_i^2 \sim \chi_n^2$$

PDF of χ_n^2 :

$$g_n(x) = \frac{1}{2^{n/2}\Gamma(n/2)} x^{n/2-1} e^{-x/2} I_{x>0}$$

Properties

– E and var of $\xi \sim \chi_n^2$

$$E(\xi) = n \quad \text{var}(\xi) = 2n$$

- For independent $\xi_i \sim \chi_{n_i}^2$, $i = 1, 2, \dots, k$:

$$\xi_0 = \sum_{i=1}^k \xi_i \sim \chi_{n_1+\dots+n_k}^2$$

- Denoted as $\Gamma(\alpha, \lambda)$:

$$\xi = \sum_{i=1}^n X_i \sim \Gamma\left(\frac{n}{2}, \frac{1}{2}\right) = \chi_n^2$$

- t Distribution: Def. t distribution with degree of freedom n :

$$T = \frac{Y}{\sqrt{\frac{\sum_{i=1}^n X_i^2}{n}}} = \frac{Y}{\sqrt{\frac{\xi}{n}}} \sim t_n$$

(Usually take ν instead of n)

PDF of t_ν :

$$t_\nu(x) = \frac{\Gamma(\frac{\nu+1}{2})}{\Gamma(\frac{\nu}{2})\sqrt{\nu\pi}} \left(1 + \frac{x^2}{\nu}\right)^{-\frac{\nu+1}{2}}$$

Denote: Upper α -fractile of t_ν , satisfies $P(T \geq c) = \alpha$:

$$c = t_{\nu, \alpha}$$

(Similar for χ_n^2 and $F_{m,n}$ etc.)

- F Distribution: Def. F distribution with degree of freedom m and n :

$$F = \frac{\sum_{i=1}^m Y_i}{\sum_{i=1}^n X_i} \sim F_{m,n}$$

PDF of $F_{m,n}$:

$$f_{m,n}(x) = \frac{\Gamma(\frac{m+n}{2})}{\Gamma(\frac{m}{2})\Gamma(\frac{n}{2})} m^{\frac{m}{2}} n^{\frac{n}{2}} x^{\frac{m}{2}-1} (n+mx)^{-\frac{m+n}{2}} I_{x>0}$$

Properties

- If $Z \sim F_{m,n}$, then $\frac{1}{Z} \sim F_{n,m}$.
- If $T \sim t_n$, then $T^2 \sim F_{1,n}$
- $F_{m,n,1-\alpha} = \frac{1}{F_{n,m,\alpha}}$

► Some useful Lemma (used in statistic inference):

- For X_1, X_2, \dots, X_n independent with $X_i \sim N(\mu_i, \sigma_i^2)$, then

$$\sum_{i=1}^n \left(\frac{X_i - \mu_i}{\sigma_i} \right)^2 \sim \chi_n^2$$

- For X_1, X_2, \dots, X_n i.i.d. $\sim N(\mu, \sigma^2)$, then

$$T = \frac{\sqrt{n}(\bar{X} - \mu)}{S} \sim t_{n-1}$$

For X_1, X_2, \dots, X_m i.i.d. $\sim N(\mu_1, \sigma^2)$, Y_1, Y_2, \dots, Y_n i.i.d. $\sim N(\mu_2, \sigma^2)$,
denote $S_\omega^2 = \frac{(m-1)S_1^2 + (n-1)S_2^2}{m+n-2}$, then

$$T = \frac{(\bar{X} - \bar{Y}) - (\mu_1 - \mu_2)}{S_\omega} \cdot \sqrt{\frac{mn}{m+n}} \sim t_{m+n-2}$$

- For X_1, X_2, \dots, X_m i.i.d. $\sim N(\mu, \sigma^2)$, Y_1, Y_2, \dots, Y_n i.i.d. $\sim N(\mu_2, \sigma^2)$, then

$$T = \frac{S_1^2 \sigma_2^2}{S_2^2 \sigma_1^2} \sim F_{m-1, n-1}$$

- For X_1, X_2, \dots, X_n i.i.d. $\sim \epsilon(\lambda)$, then

$$2\lambda n \bar{X} = 2\lambda \sum_{i=1}^n X_i \sim \chi_{2n}^2$$

Remark: for $X_i \sim \epsilon(\lambda) = \Gamma(1, \lambda) \Rightarrow 2\lambda \sum_{i=1}^n X_i \sim \Gamma(n, 1/2) = \chi_{2n}^2$.

2 统计推断部分

Statistical Inference: use sample to estimate population.

Two main tasks of Statistical Inference:

- Parameter Estimation
 - Point Estimation: 2.2
 - Interval Estimation: 2.3
- Hypothesis Testing: 2.4

2.1 Statistical Model and Statistics

Random sample comes from population X . In parametric model case, we have population distribution family:

$$\mathcal{F} = \{f(x; \vec{\theta}) | \vec{\theta} \in \Theta\}$$

where parameter $\vec{\theta}$ reflect some quantities of population (e.g. mean, variance, etc.), each $\vec{\theta}$ corresponds to a distribution of population X .

Sample space: Def. as $\mathcal{X} = \{\{x_1, x_2, \dots, x_n\}, \forall x_i\}$, then $\{X_i\} \in \mathcal{X}$ is random sample from population $X \sim f(x; \vec{\theta})$.

2.1.1 Statistics

Statistic(s): function of random sample $\vec{T}(X_1, X_2, \dots, X_n)$, **but not a function of parameter**.

Some useful statistics, e.g.

- Sample mean (Consider X_i i.i.d.)

$$\bar{X} = \frac{1}{n} \sum_{i=1}^n X_i$$

- Sample variance

$$S^2 = \frac{1}{n-1} \sum_{i=1}^n (X_i - \bar{X})^2$$

- Sample moments

- Origin moment

$$a_{n,k} = \frac{1}{n} \sum_{i=1}^n X_i^k \quad k = 1, 2, 3, \dots$$

- Center moment

$$m_{n,k} = \frac{1}{n} \sum_{i=1}^n (X_i - \bar{X})^k \quad k = 2, 3, 4, \dots$$

- Order statistics

$$(X_{(1)}, X_{(2)}, \dots, X_{(n)}), \text{ for } X_{(1)} \leq X_{(2)} \leq \dots \leq X_{(n)}$$

- Sample p -fractile

$$m_p = X_{(m)}, \quad m = [(n+1)p]$$

- Sample coefficient of variation

$$\hat{\nu} = \frac{S}{\bar{X}}$$

- Skewness and Kurtosis

$$\hat{\beta}_1 = \frac{m_{n,3}}{m_{n,2}^{3/2}} \quad \hat{\beta}_2 = \frac{m_{n,4}}{m_{n,2}^2}$$

► Properties

Statistic T is a function of random sample $\{X_i\}$, thus has distribution (say $g_T(t)$) called **Sampling Distribution**.

For X_i i.i.d. from $X \sim f(x)$ with population mean μ and variance σ^2

- Calculation of S^2

$$(n-1)S^2 = \sum_{i=1}^n x_i^2 - n\bar{x}^2$$

- E and var of \bar{X} and S^2

$$E(\bar{X}) = \mu \quad var(\bar{X}) = \frac{\sigma^2}{n} \quad E(S^2) = \sigma^2$$

Further if X_i i.i.d. from $X \sim N(\mu, \sigma^2)$ where μ and σ^2 unknown.

- Independence of \bar{X} and S^2

\bar{X} and S^2 independent

- Distribution of $\bar{X} = \frac{1}{n} \sum_{i=1}^n X_i$

$$\bar{X} \sim N\left(\mu, \frac{\sigma^2}{n}\right)$$

- Distribution of $S^2 = \frac{1}{n-1} \sum_{i=1}^n (X_i - \bar{X})^2$

$$\frac{(n-1)S^2}{\sigma^2} \sim \chi_{n-1}^2$$

2.1.2 Exponential Family

Def. $\mathcal{F} = \{f(x; \vec{\theta}) | \vec{\theta} \in \Theta\}$ is **Exponential Family** if $f(x; \vec{\theta})$ has the form as

$$f(x; \vec{\theta}) = C(\theta)h(x) \exp \left[\sum_{i=1}^k Q_i(\theta)T_i(x) \right] \quad \vec{\theta} \in \Theta$$

Canonical Form: Take $Q_i(\theta) = \varphi_i$, then $\vec{\varphi} = (\varphi_1, \varphi_2, \dots, \varphi_k) = (Q_1(\theta), Q_2(\theta), \dots, Q_k(\theta))$ is a transform from Θ to Θ^* , s.t. \mathcal{F} has canonical form, i.e.

$$f(x; \vec{\varphi}) = C^*(\vec{\varphi})h(x) \exp \left[\sum_{i=1}^k \varphi_i T_i(x) \right] \quad \vec{\varphi} \in \Theta^*$$

Θ^* is canonical parameter space.

- Why we need exponential family? Have some nice properties.

2.1.3 Sufficient and Complete Statistics

- A **Sufficient Statistic** $T(\vec{X})$ for $\vec{\theta}$ contains all the information of sample when infer $\vec{\theta}$, i.e.

$$f(\vec{X}; T(\vec{X})) = f(\vec{X}; T(\vec{X}), \vec{\theta})$$

Properties

- **Factorization Thm.** $T(\vec{X})$ is sufficient **if and only if** $f_{\vec{X}}(\vec{x}; \vec{\theta}) = f(\vec{x}; \vec{\theta})$ can be written as

$$f(\vec{x}; \vec{\theta}) = g[t(\vec{x}); \vec{\theta}]h(\vec{x})$$

- If $T(\vec{X})$ sufficient, then $T'(\vec{X}) = g[T(\vec{X})]$ also. (require g single-valued and invertible)
- If $T(\vec{X})$ sufficient, then (T, T_1) also.
- Minimal sufficient statistic $T_{\theta}(\vec{X})$ satisfies

$$\forall \text{ sufficient statistic } S, \exists q_S(\cdot), \text{ s.t. } T_{\theta} = q_S(S)$$

A minimal sufficient statistic not always exists.

Sufficient & Complete \Rightarrow Minimal sufficient.

- Usually dimension of \vec{T}_{θ} and $\vec{\theta}$ equals.

Sufficient statistic is not unique.

- A **Complete Statistic** $T(\vec{X})$ for $\vec{\theta}$ satisfies

$$\forall \vec{\theta} \in \Theta; \forall \varphi \text{ satisfies } E[\varphi(T(\vec{X}))] = 0, \text{ we have } P[\varphi(T) = 0; \vec{\theta}] = 1$$

Explanation: $T \sim g_T(t)$. Rewrite as

$$\int \varphi(t) g_T(t) dt = 0 \quad \forall \vec{\theta} \Rightarrow \varphi(T) = 0 \text{ a.s.}$$

i.e. $\text{span}\{g_T(t); \forall \vec{\theta}\}$ is a complete space. Or to say that \nexists none-zero $\varphi(t)$ so that $E(\varphi(T)) = 0$ (unbiased estimation)

$$\varphi(T) \neq 0 \quad \forall \vec{\theta} \Rightarrow E[\varphi(T(\vec{X}))] \neq 0$$

So make sure the uniqueness of unbiased estimation of $\hat{\theta}$ using T .

Properties

- If $T(\vec{X})$ complete, then $T'(\vec{X}) = g[T(\vec{X})]$ also. (require g measurable)
- A complete statistic not always exists.

- An **Ancillary Statistic** $S(\vec{X})$ is a statistic whose distribution does not depend on $\vec{\theta}$

Basu Thm: $\vec{X} = (X_1, X_2, \dots, X_n)$ is sample from $\mathcal{F} = \{f(x; \theta), \theta \in \Theta\}$. $T(\vec{X})$ is a complete and minimal sufficient statistic, $S(\vec{X})$ is ancillary statistic, then $S(\vec{X}) \perp\!\!\!\perp T(\vec{X})$.

- Exponential family: For $\vec{X} = (X_1, X_2, \dots, X_n)$ from exponential family with canonical form, i.e.

$$f(\vec{x}; \vec{\theta}) = C(\vec{\theta})h(\vec{x}) \exp \left[\sum_{i=1}^k \theta_i T_i(\vec{x}) \right], \quad \vec{\theta} = (\theta_1, \theta_2, \dots, \theta_k) \in \Theta$$

Then if $\Theta \in \mathbb{R}^k$ interior point exists, then $T(\vec{X}) = (T_1(\vec{X}), T_2(\vec{X}), \dots, T_k(\vec{X}))$ is sufficient & complete statistic.

2.2 Point Estimation

For parametric distribution family $\mathcal{F} = \{f(x, \vec{\theta}), \vec{\theta} \in \Theta\}$, random sample $\vec{X} = (X_1, X_2, \dots, X_n)$ from \mathcal{F} . $g(\vec{\theta})$ is a function defined on Θ .

Mission: use sample $\{X_i\}$ to estimate $g(\vec{\theta})$, called **Parameter Estimation**.

$$\text{Parameter Estimation} \begin{cases} \text{Point Estimation} & \checkmark \\ \text{Interval Estimation} \end{cases}$$

Point estimation: when estimating θ or $g(\theta)$, denote the estimator (defined on sample space \mathcal{X}) as

$$\hat{\theta}(\vec{X}) \quad \hat{g}(\vec{X})$$

Estimator is a statistic, with sampling distribution.

2.2.1 Optimal Criterion

Some nice properties of estimators (that we expect)

- Unbiasedness

$$E(\hat{\theta}) = \theta \quad \text{or} \quad E(\hat{g}(\vec{X})) = g(\theta)$$

Otherwise, say $\hat{\theta}$ or \hat{g} biased. Def. **Bias:** $E(\hat{\theta}) - \theta$

Asymptotically unbiasedness

$$\lim_{n \rightarrow \infty} E(\hat{g}(\vec{X})) = g(\theta)$$

- Efficiency: say $\hat{g}_1(\vec{X})$ is more efficient than $\hat{g}_2(\vec{X})$, if

$$\text{var}(\hat{g}_1) \leq \text{var}(\hat{g}_2) \quad \forall \theta \in \Theta$$

- Mean Squared Error (MSE)

$$\text{MSE} = E[(\hat{\theta} - \theta)^2] = \text{var}(\hat{\theta}) + [\text{Bias}(\hat{\theta})]^2$$

For unbiased estimator, i.e. $Bias(\hat{\theta}) = 0$, we have

$$MSE = E[(\hat{\theta} - \theta)^2] = var(\hat{\theta})$$

- Consistency

$$\lim_{n \rightarrow \infty} P(|\hat{g}_n(\vec{X}) - g(\theta)| \geq \varepsilon) = 0 \quad \forall \varepsilon > 0$$

- Asymptotic Normality

2.2.2 Method of Moments

Review: Population moments & Sample moments

$$\begin{aligned} \alpha_k &= E(X^k) & \mu_k &= E[(X - E(X))^k] \\ a_{n,k} &= \frac{1}{n} \sum_{i=1}^n X_i^k & m_{n,k} &= \frac{1}{n} \sum_{i=1}^n (X_i - \bar{X})^k \end{aligned}$$

Property: $a_{n,k}$ is the unbiased estimator of α_k (while $m_{n,k}$ usually biased for μ_k)

For sample $\vec{X} = (X_1, X_2, \dots, X_n)$ from $\mathcal{F} = \{f(x; \theta, \theta \in \Theta)\}$, unknown parameter (or its function) $g(\theta)$ can be written as

$$g(\theta) = G(\alpha_1, \alpha_2, \dots, \alpha_k; \mu_2, \mu_3, \dots, \mu_l)$$

Then its **Moment Estimate** $\hat{g}(\vec{X})$ is

$$\hat{g}(\vec{X}) = G(a_{n,1}, a_{n,2}, \dots, a_{n,k}; m_{n,2}, m_{n,3}, \dots, m_{n,l})$$

Example: coefficient of variance & skewness

$$\hat{\nu} = \frac{S}{\bar{X}} \quad \hat{\beta}_1 = \frac{m_{n,3}}{m_{n,2}^{3/2}} = \sqrt{n} \frac{\sum_{i=1}^n (X_i - \bar{X})^3}{[\sum_{i=1}^n (X_i - \bar{X})^2]^{\frac{3}{2}}}$$

► Note:

- G may not have explicit expression.
- Moment estimate may not be unique.
- If $G = \sum_{i=1}^k c_i \alpha_i$ (linear combination of α , without μ), then $\hat{g}(\vec{X}) = \sum_{i=1}^k c_i a_{n,i}$ unbiased.

Usually $\hat{g}(\vec{X})$ is asymptotically unbiased.

- For small sample, not so accurate.
- May not contain all the information about $\vec{\theta}$, i.e. may not be sufficient statistic.
- Do not require a statistic model.

2.2.3 Maximum Likelihood Estimation

For sample $\vec{X} = (X_1, X_2, \dots, X_n)$ with distribution $f(\vec{x}; \vec{\theta})$ from $\mathcal{F} = \{f(x; \vec{\theta}), \vec{\theta} \in \Theta\}$, def. **Likelihood Function** $L(\vec{\theta}; \vec{x})$, defined on Θ (as a function of $\vec{\theta}$)

$$L(\vec{\theta}; \vec{x}) = f(\vec{x}; \vec{\theta}) \quad \vec{\theta} \in \Theta, \vec{x} \in \mathcal{X}$$

Also def. log-likelihood function $l(\vec{\theta}; \vec{x}) = \ln L(\vec{\theta}; \vec{x})$.

If estimator $\hat{\theta} = \hat{\theta}(\vec{X})$ satisfies

$$L(\hat{\theta}; \vec{x}) = \sup_{\vec{\theta} \in \Theta} L(\vec{\theta}; \vec{x}), \quad \vec{x} \in \mathcal{X}$$

Or equivalently take $l(\vec{\theta}; \vec{x})$ instead of $L(\vec{\theta}; \vec{x})$.

Then $\hat{\theta} = \hat{\theta}(\vec{X})$ is a **Maximum Likelihood Estimate**(MLE) of $\vec{\theta} = (\theta_1, \theta_2, \dots, \theta_k)$

How to identify MLE?

- Differentiation: Fermat Lemma

$$\left. \frac{\partial L}{\partial \theta_i} \right|_{\vec{\theta}=\hat{\theta}} = 0 \quad \left. \frac{\partial^2 L}{\partial \theta_i \partial \theta_j} \right|_{\vec{\theta}=\hat{\theta}} \text{ negative definite} \quad \forall i, j = 1, 2, \dots, k$$

- Graphing method.
- Numerically compute maximum.

► Some properties of MLE

- (Depend on the case, not always) unbiased.
- Invariance of MLE: If $\hat{\theta}$ is MLE of $\vec{\theta}$, invertible function $g(\vec{\theta})$, then $g(\hat{\theta})$ is MLE of $g(\vec{\theta})$.
- MLE and Sufficiency: $T = T(X_1, X_2, \dots, X_n)$ is a sufficient statistic of $\vec{\theta}$, if MLE of $\vec{\theta}$ exists, say $\hat{\theta}$, then $\hat{\theta}$ is a function of T , i.e.

$$\hat{\theta} = \hat{\theta}(\vec{X}) = \hat{\theta}^*(T(\vec{X}))$$

- Asymptotic Normality:

$$\sqrt{n}(\hat{\theta}_n - \theta) \xrightarrow{d} N(0, \sigma_\theta^2), \quad \sigma_\theta^2 = \frac{1}{E_\theta[\frac{\partial}{\partial \theta} \ln f(\vec{X}; \theta)]^2}$$

i.e.

$$\hat{\theta}_n \xrightarrow{d} N(\theta, \frac{\sigma_\theta^2}{n})$$

► Comparison: MoM and MLE

- MoM do not require statistic model; MLE need to know PDF.
- MoM is more robust than MLE.

MLE in Exponential Family:

For sample $\vec{X} = (X_1, X_2, \dots, X_n)$ from canonical exponential family $\mathcal{F} = \{f(x; \vec{\theta}), \vec{\theta} \in \Theta\}$

$$f(x; \vec{\theta}) = C(\vec{\theta})h(x) \exp \left[\sum_{i=1}^k \theta_i T_i(x) \right] \quad \vec{\theta} = (\theta_1, \dots, \theta_k) \in \Theta$$

Likelihood function $L(\vec{\theta}, \vec{x}) = \prod_{j=1}^n f(x_j; \vec{\theta})$ and log-likelihood function $l(\vec{\theta}, \vec{x})$

$$L(\vec{\theta}, \vec{x}) = C^n(\vec{\theta}) \prod_{j=1}^n h(x_j) \exp \left[\sum_{i=1}^k \theta_i \sum_{j=1}^n T_i(x_j) \right]$$

$$l(\vec{\theta}, \vec{x}) = n \ln C(\vec{\theta}) + \sum_{j=1}^n \ln h(x_j) + \sum_{i=1}^k \theta_i \sum_{j=1}^n T_i(x_j)$$

Solution of MLE: (Require $\hat{\theta} \in \Theta$)

$$\frac{n}{C(\vec{\theta})} \frac{\partial C(\vec{\theta})}{\partial \theta_i} \bigg|_{\vec{\theta}=\hat{\theta}} = - \sum_{j=1}^n T_i(x_j), \quad i = 1, 2, \dots, k$$

2.2.4 Uniformly Minimum Variance Unbiased Estimate

MSE: For $\hat{g}(\vec{X})$ is estimate of $g(\vec{\theta})$, then MSE

$$\text{MSE}(\hat{g}(\vec{X})) = E[(\hat{g}(\vec{X}) - g(\vec{\theta}))^2] = \text{var}(\hat{g}) + [\text{Bias}(\hat{g})]^2$$

► Unbiased estimator (i.e. $\text{Bias}(\hat{g}) = 0$) not unique; not always exist.

Now only consider unbiased estimators of $g(\vec{\theta})$ exists, say $\hat{g}(\vec{X})$, then

$$\text{MSE}(\hat{g}(\vec{X})) = \text{var}(\hat{g}(\vec{X}))$$

If \forall unbiased estimate $\hat{g}'(\vec{X})$, \hat{g} satisfies

$$\text{var}[\hat{g}(\vec{X})] \leq \text{var}[\hat{g}'(\vec{X})]$$

Then $\hat{g}(\vec{X})$ is **Uniformly Minimum Variance Unbiased Estimate (UMVUE)** of $g(\vec{\theta})$

How to determine UMVUE? (Not an easy task)

- Zero Unbiased Estimate Method
- Sufficient and Complete Statistic Method
- Cramer-Rao Inequality

1. Zero Unbiased Estimate Method

Let $\hat{g}(\vec{X})$ be an unbiased estimate with $\text{var}(\hat{g}) < \infty$. If $\forall E(\hat{l}(\vec{X})) = 0$, \hat{g} holds that

$$\text{cov}(\hat{g}, \hat{l}) = E(\hat{g} \cdot \hat{l}) = 0, \quad \forall \theta \in \Theta$$

Then \hat{g} is a UMVUE of $g(\vec{\theta})$ (sufficient & necessary).

2. Sufficient and Complete Statistic Method

For $T(\vec{X})$ sufficient statistic, $\hat{g}(\vec{X})$ unbiased estimate of $g(\vec{\theta})$, then

$$h(T) = E(\hat{g}(\vec{X})|T)$$

is an unbiased estimate of $g(\vec{\theta})$ and $\text{var}(h(T)) \leq \text{var}(\hat{g})$.

Remark:

- A method to improve estimator.
- A UMVUE has to be a function of sufficient statistic.

Lehmann-Scheffé Thm.: For $\vec{X} = (X_1, X_2, \dots, X_n)$ from population $X \sim \mathcal{F} = \{f(x, \vec{\theta}, \theta \in \Theta)\}$. $T(\vec{X})$ sufficient and complete, and $\hat{g}(T(\vec{X}))$ be an unbiased estimator, then $\hat{g}(T(\vec{X}))$ is the unique UMVUE.

Can be used to construct UMVUE: given $T(\vec{X})$ sufficient and complete and some unbiased estimator $\hat{g}'(\vec{\theta})$ then

$$\hat{g}(T) = E(\hat{g}'|T)$$

is the unique UMVUE.

3. Cramer-Rao Inequality

Core idea: determine a lower bound of $\text{var}(\hat{g})$.

Consider $\vec{\theta} = \theta$ (One dimension parameter); For $\{X_i\}$ i.i.d. $f(x, \theta)$: def.

- **Score function:** Reflects the steepness of likelihood function f .

$$S(\vec{x}; \theta) = \frac{\partial \ln f(\vec{x}; \theta)}{\partial \theta} = \sum_{i=1}^n \frac{\partial \ln f(x_i; \theta)}{\partial \theta}$$

$$E[S(\vec{X}; \theta)] = 0$$

- **Fisher Information:** Variance of $S(\vec{x}; \theta)$, reflects the accuracy to conduct estimation, i.e. reflects information of statistic model.

$$I(\theta) = E \left[\left(\frac{\partial \ln f(\vec{x}; \theta)}{\partial \theta} \right)^2 \right] = -E \left[\frac{\partial^2 \ln f(\vec{x}; \theta)}{\partial \theta^2} \right]$$

Consider \mathcal{F} satisfies some regularity conditions (in most cases, regularity conditions do hold), then the lower bound of $\text{var}(\hat{g})$ satisfies **Cramer-Rao Inequality**:

$$\text{var}(\hat{g}(\vec{X})) \geq \frac{[g'(\theta)]^2}{nI(\theta)}$$

Special case: $g(\theta) = \theta$ then

$$\text{var}(\hat{\theta}) \geq \frac{1}{nI(\theta)}$$

note:

- C-R Inequality determine a lower bound, not the infimum(i.e. UMVUE $\nRightarrow \text{var}(\hat{g}(\vec{X})) = \frac{[g'(\theta)]^2}{nI(\theta)}$).
- Take '=': Only some cases in Exponential family.
- **Efficiency**: How good the estimator is.

$$e_{\hat{g}(\vec{X})}(\theta) = \frac{[g'(\theta)]^2 / (nI(\theta))}{\text{var}(\hat{g}(\vec{X}))}$$

4. Multi-Dimensional Cramer-Rao Inequality

ReDef. Fisher Information:

$$\mathbf{I}(\vec{\theta}) = \{I_{ij}(\vec{\theta})\} = \{E \left[\left(\frac{\partial \ln f(\vec{x}; \theta)}{\partial \theta_i} \right) \left(\frac{\partial \ln f(\vec{x}; \theta)}{\partial \theta_j} \right) \right]\}$$

Then covariance matrix $\Sigma(\vec{\theta})$ satisfies **Cramer-Rao Inequality**

$$\Sigma(\vec{\theta}) \geq (n\mathbf{I}(\vec{\theta}))^{-1}$$

Note: ' \geq ' holds for all diagonal elements, i.e.

$$\text{var}(\hat{\theta}_i) \geq \frac{I_{ii}^*(\vec{\theta})}{n}, \quad \forall i = 1, 2, \dots, k$$

2.2.5 MoM and MLE in Linear Regression

- Linear Regression Model(1-dimension case):

$$y_i = \beta_0 + \beta_1 x_i + \epsilon_i$$

where β_0, β_1 are regression coefficient, and ϵ_i are unknown random **error**. Assume:

ϵ_i are i.i.d.

$$E(\epsilon_i | x_i) = 0$$

$$\text{var}(\epsilon_i) = \sigma^2$$

Mission: use data $\{(x_i, y_i)\}$ to estimate β_0, β_1 (i.e. regression line), and error ϵ_i .

1. OLS: Take β_0, β_1 so that MSE min, i.e.

$$(\hat{\beta}_0, \hat{\beta}_1) = \arg \min \sum_{i=1}^n (y_i - \beta_0 - \beta_1 x_i)^2$$

Solution:

$$\begin{cases} \hat{\beta}_0 &= \bar{y} - \beta_1 \bar{x} \\ \hat{\beta}_1 &= \frac{\sum_{i=1}^n (x_i - \bar{x})(y_i - \bar{y})}{\sum_{i=1}^n (x_i - \bar{x})^2} = \{\sum_{i=1}^n (x_i - \bar{x})^2\}^{-1} \{\sum_{i=1}^n (x_i - \bar{x})(y_i - \bar{y})\} \end{cases}$$

So get regression line: $y = \hat{\beta}_0 + \hat{\beta}_1 x$

Def. Residuals

$$\hat{\epsilon}_i = y_i - \hat{y}_i = y_i - (\hat{\beta}_0 + \hat{\beta}_1 x_i)$$

Residuals can be used to estimate ϵ_i : $E[(\epsilon_i)^2] = \sigma^2$

$$\hat{\sigma}^2 = \frac{1}{n-2} \sum_{i=1}^n (y_i - \hat{\beta}_0 - \hat{\beta}_1 x_i)^2$$

2. MoM: Consider r.v. $\epsilon \sim f(\epsilon; x, y, \beta_0, \beta_1)$, sample $\{\epsilon_i | \epsilon_i = y_i - \beta_0 - \beta_1 x_i\}$, then obviously

$$\bar{\epsilon} = \bar{y} - \beta_0 - \beta_1 \bar{x}$$

Take moment estimate of ϵ , we have

$$E(\epsilon_i) = 0 \quad E(\epsilon_i x_i) = 0 \text{ (note that)} E(\epsilon | x) = 0$$

$$\text{i.e.} \begin{cases} \frac{1}{n} \sum_{i=1}^n (y_i - \beta_0 - \beta_1 x_i) = 0 \\ \frac{1}{n} \sum_{i=1}^n x_i (y_i - \beta_0 - \beta_1 x_i) = 0 \end{cases}$$

Solution:

$$\begin{cases} \hat{\beta}_0 = \bar{y} - \beta_1 \bar{x} \\ \hat{\beta}_1 = \frac{\sum_{i=1}^n (x_i - \bar{x})(y_i - \bar{y})}{\sum_{i=1}^n (x_i - \bar{x})^2} \end{cases}$$

(Same as OLS)

Moment estimate of σ^2

$$\hat{\sigma}_n^2 = \frac{1}{n} \sum_{i=1}^n (y_i - \hat{\beta}_0 - \hat{\beta}_1 x_i)^2$$

3. MLE: Assume $\epsilon_i \sim N(0, \sigma^2)$, then $y_i | x_i \sim N(\beta_0 + \beta_1 x_i, \sigma^2)$. Get likelihood function:

$$L(\beta_0, \beta_1, \sigma^2; x_1, \dots, x_n, y_1, \dots, y_n) = (2\pi\sigma^2)^{-\frac{n}{2}} \exp \left[-\frac{\sum_{i=1}^n (y_i - \beta_0 - \beta_1 x_i)^2}{2\sigma^2} \right]$$

Take differentiation, also get the same result.

► Linear Regression Model(Multi-dimension case):

$$y_i = \beta_0 + \beta_1 x_{i1} + \beta_2 x_{i2} + \dots + \beta_p x_{ip} + \epsilon_i$$

Denote: $\vec{\beta} = (\beta_0, \beta_1, \dots, \beta_p)$, $\vec{x}_i = (1, x_{i1}, x_{i2}, \dots, x_{ip})$, then for each i : $y_i = \vec{x}_i^T \vec{\beta} + \epsilon_i$

Further denote: Matrix form:

$$Y = \begin{pmatrix} y_1 \\ y_2 \\ \vdots \\ y_n \end{pmatrix} = \begin{pmatrix} 1 & x_{11} & \dots & x_{1p} \\ 1 & x_{21} & \dots & x_{2p} \\ \vdots & \vdots & \ddots & \vdots \\ 1 & x_{n1} & \dots & x_{np} \end{pmatrix} \begin{pmatrix} \beta_0 \\ \beta_1 \\ \vdots \\ \beta_p \end{pmatrix} + \begin{pmatrix} \epsilon_1 \\ \epsilon_2 \\ \vdots \\ \epsilon_n \end{pmatrix} = X\vec{\beta} + \vec{\epsilon}$$

Basic Assumptions: Gauss-Markov Assumptions

- OLS unbiased

$$E(\epsilon_i|x_i) = 0 \quad E(y_i|x_i) = x_i^T \beta$$

- Homogeneity of ϵ_i

$$\text{var}(\epsilon_i) = \sigma^2$$

- Independent of ϵ
- (For MLE) ϵ_i i.i.d. $\sim N(0, \sigma^2)$

Residuals:

$$\hat{\epsilon}_i = y_i - \hat{y}_i = y_i - x_i^T \hat{\beta}$$

Def. Residual Sum of Squares (RSS)

$$\text{RSS} = \sum_{i=1}^n \hat{\epsilon}_i^2 = \sum_{i=1}^n (y_i - x_i^T \hat{\beta})^2$$

Estimator exists and unique: ($\hat{\sigma}^2$ after bias correction)

$$\begin{aligned} \hat{\beta} &= (X^T X)^{-1} (X^T Y) \\ \hat{\sigma}_n^2 &= \frac{1}{n} \sum_{i=1}^n (y_i - x_i^T \hat{\beta})^2 \\ \hat{\sigma}^2 &= \frac{1}{n - p - 1} \sum_{i=1}^n (y_i - x_i^T \hat{\beta})^2 \end{aligned}$$

2.2.6 Kernel Density Estimate

Given random sample $\{X_i\}$. Def. Empirical Distribution Function

$$\hat{F}_n(x) = \frac{1}{n} \sum_{i=1}^n I_{(-\infty, x]}(X_i) \quad (2.1)$$

Problem: Overfitting when getting \hat{f} . Solution: Using **Kernel Estimate**, replace $I_{(-\infty, x]}(\cdot)$ with Kernel function $K(\cdot)$, then

$$\hat{f}_n(x) = \frac{F_n(x + h_n) - F_n(x - h_n)}{2h_n} = \frac{1}{nh_n} \sum_{i=1}^n K\left(\frac{x - X_i}{h_n}\right)$$

where h_n is **bandwidth**. Take proper kernel function K to get estimate of f .

Can be considered as a convolution of sample $\{X_i\}$ and kernel function K .

Useful Kernel Functions:

- $K(x) = \frac{1}{2} I_{[-\frac{1}{2}, \frac{1}{2}]}$
- $K(x) = (1 - |x|) I_{[-1, 1]}$
- $K(x) = \frac{1}{2\pi} e^{-\frac{x^2}{2}}$

- $K(x) = \frac{1}{\pi(1+x^2)}$
- $K(x) = \frac{1}{2\pi} \text{sinc}^2\left(\frac{x}{2}\right)$

2.3 Interval Estimation

$$\text{Parameter Estimation} \begin{cases} \text{Point Estimation} \\ \text{Interval Estimation} \end{cases} \quad \checkmark$$

Interval Estimation: to estimate $g(\vec{\theta})$, give **two** estimators $\hat{g}_1(\vec{X})$, $\hat{g}_2(\vec{X})$ defined on \mathcal{X} as the two ends of interval (i.e. give an interval $[\hat{g}_1(\vec{X}), \hat{g}_2(\vec{X})]$), then random interval $[\hat{g}_1(\vec{X}), \hat{g}_2(\vec{X})]$ is an **Interval Estimation** of $g(\theta)$.

2.3.1 Confidence Interval

How to judge an interval estimation?

- Reliability

$$P(g(\theta) \in [\hat{g}_1, \hat{g}_2])$$

- Precision

$$E(\hat{g}_2 - \hat{g}_1)$$

Trade off: (in most cases)

Given a level of reliability, find an interval with the highest precision above the level

► For a given $0 < \alpha < 1$, if

$$P(\hat{g}_1 \leq g(\vec{\theta}) \leq \hat{g}_2) \geq 1 - \alpha$$

then $[\hat{g}_1, \hat{g}_2]$ is a **Confidence Interval** for $g(\vec{\theta})$, with **Confidence Level** $1 - \alpha$.

Confidence Coefficient:

$$\inf_{\forall \vec{\theta} \in \Theta} P(\vec{\theta} \in \Theta)$$

Other cases:

- **Confidence Limit:** Upper/Lower Confidence Limit

$$P(g \leq \hat{g}_U) \geq 1 - \alpha$$

$$P(\hat{g}_L \leq \theta) \geq 1 - \alpha$$

- **Confidence Region:** For high dimensional parameters $\vec{g} = (g_1, g_2, \dots, g_k)$

$$P(\vec{g} \in S(\vec{X})) \geq 1 - \alpha \quad \forall \vec{\theta} \in \Theta$$

Mission: Determine \hat{g}_1, \hat{g}_2 .

2.3.2 Pivot Variable Method

Idea: Based on point estimation, construct a new variable and thus find the interval estimation.

Def. **Pivot Variable** T , satisfies:

- Expression of T contains θ (thus T is not a statistic).
- Distribution of T independent of θ .

In different cases, construct different pivot variable, usually base on sufficient statistics and transform.

Knowing a proper pivot variable $T = T(\hat{\varphi}, g(\theta)) \sim f$, (f is some distribution independent of $\vec{\theta}$), $\hat{\varphi}$ is a sufficient statistic), then we can take T satisfies:

$$P(f_{1-\frac{\alpha}{2}} \leq T \leq f_{\frac{\alpha}{2}}) = 1 - \alpha$$

Construct the inverse mapping of $T = T(\hat{\varphi}, g(\theta)) \Leftrightarrow g(\theta) = T^{-1}(T, \hat{\varphi})$, we get

$$P[T^{-1}(f_{1-\frac{\alpha}{2}}, \hat{\varphi}) \leq \hat{g} \leq T^{-1}(f_{\frac{\alpha}{2}}, \hat{\varphi})] = 1 - \alpha$$

Thus get a confidence interval for θ with confidence coefficient $1 - \alpha$.

2.3.3 Confidence Interval for Common Distributions

Some important properties of χ^2 , t and F see section 1.8.2.

1. Single normal population: $\vec{X} = \{X_1, X_2, \dots, X_n\} \in \mathcal{X}$ i.i.d from Normal Distribution population $N(\mu, \sigma^2)$. Denote sample mean and sample variance:

$$\bar{X} = \frac{1}{n} \sum_{i=1}^n X_i \quad S^2 = \frac{1}{n-1} \sum_{i=1}^n (X_i - \bar{X})^2 \quad S_\mu = \frac{1}{n} \sum_{i=1}^n (X_i - \mu)^2, (\text{for } \mu \text{ known})$$

Estimating μ & σ^2 : construction of pivot variable under different circumstances:

Estimation	Pivot Variable	Confidence Interval
σ^2 known, estimate μ	$T = \frac{\sqrt{n}(\bar{X} - \mu)}{\sigma} \sim N(0, 1)$	$\left[\bar{X} - \frac{\sigma}{\sqrt{n}} N_{\frac{\alpha}{2}}, \bar{X} + \frac{\sigma}{\sqrt{n}} N_{\frac{\alpha}{2}} \right]$
σ^2 unknown, estimate μ	$T = \frac{\sqrt{n}(\bar{X} - \mu)}{S} \sim t_{n-1}$	$\left[\bar{X} - \frac{S}{\sqrt{n}} t_{n-1, \frac{\alpha}{2}}, \bar{X} + \frac{S}{\sqrt{n}} t_{n-1, \frac{\alpha}{2}} \right]$
μ known, estimate σ^2	$T = \frac{nS_\mu^2}{\sigma^2} \sim \chi_n^2$	$\left[\frac{nS_\mu^2}{\chi_{n, \frac{\alpha}{2}}^2}, \frac{nS_\mu^2}{\chi_{n, 1-\frac{\alpha}{2}}^2} \right]$
μ unknown, estimate σ^2	$T = \frac{(n-1)S^2}{\sigma^2} \sim \chi_{n-1}^2$	$\left[\frac{(n-1)S^2}{\chi_{n-1, \frac{\alpha}{2}}^2}, \frac{(n-1)S^2}{\chi_{n-1, 1-\frac{\alpha}{2}}^2} \right]$

2. Double normal population: $\vec{X} = \{X_1, X_2, \dots, X_m\}$ i.i.d. from $N(\mu_1, \sigma_1^2)$; $\vec{Y} = \{Y_1, Y_2, \dots, Y_n\}$ i.i.d. from $N(\mu_2, \sigma_2^2)$

Denote sample mean, sample variance and pooled sample variance:

$$\begin{aligned}\bar{X} &= \frac{1}{m} \sum_{i=1}^m X_i & S_X^2 &= \frac{1}{m-1} \sum_{i=1}^m (X_i - \bar{X})^2 & S_{\mu_1}^2 &= \frac{1}{m} \sum_{i=1}^m (X_i - \mu_1)^2, (\mu_1 \text{ known}) \\ \bar{Y} &= \frac{1}{n} \sum_{i=1}^n Y_i & S_Y^2 &= \frac{1}{n-1} \sum_{i=1}^n (Y_i - \bar{Y})^2 & S_{\mu_2}^2 &= \frac{1}{n} \sum_{i=1}^n (Y_i - \mu_2)^2, (\mu_2 \text{ known}) \\ S_\omega^2 &= \frac{(m-1)S_X^2 + (n-1)S_Y^2}{m+n-2}\end{aligned}$$

Estimating $\mu_1 - \mu_2$:

When $\sigma_1^2 \neq \sigma_2^2$ unknown, estimate $\mu_1 - \mu_2$: Behrens-Fisher Problem, remain unsolved, but can deal with simplified cases.

Estimation	Pivot Variable	Confidence Interval
$\sigma_1^2 \& \sigma_2^2$ known, estimate $\mu_1 - \mu_2$	$T = \frac{\bar{X} - \bar{Y} - (\mu_1 - \mu_2)}{\sqrt{\frac{\sigma_1^2}{m} + \frac{\sigma_2^2}{n}}} \sim N(0, 1)$	$\left[\bar{X} - \bar{Y} - N_{\frac{\alpha}{2}} \sqrt{\frac{\sigma_1^2}{m} + \frac{\sigma_2^2}{n}}, \right. \\ \left. \bar{X} - \bar{Y} + N_{\frac{\alpha}{2}} \sqrt{\frac{\sigma_1^2}{m} + \frac{\sigma_2^2}{n}} \right]$
$\sigma_1^2 = \sigma_2^2$ unknown, estimate $\mu_1 - \mu_2$	$T = \frac{\bar{X} - \bar{Y} - (\mu_1 - \mu_2)}{S_\omega \sqrt{\frac{1}{m} + \frac{1}{n}}} \sim t_{m+n-2}$	$\left[\bar{X} - \bar{Y} - S_\omega t_{m+n-2, \frac{\alpha}{2}} \sqrt{\frac{1}{m} + \frac{1}{n}}, \right. \\ \left. \bar{X} - \bar{Y} + S_\omega t_{m+n-2, \frac{\alpha}{2}} \sqrt{\frac{1}{m} + \frac{1}{n}} \right]$
Welch's t -Interval (when m, n large enough)	$T = \frac{\bar{X} - \bar{Y} - (\mu_1 - \mu_2)}{\sqrt{\frac{S_X^2}{m} + \frac{S_Y^2}{n}}} \xrightarrow{\mathcal{L}} N(0, 1)$	$\left[\bar{X} - \bar{Y} - N_{\frac{\alpha}{2}} \sqrt{\frac{S_1^2}{m} + \frac{S_2^2}{n}}, \right. \\ \left. \bar{X} - \bar{Y} + N_{\frac{\alpha}{2}} \sqrt{\frac{S_1^2}{m} + \frac{S_2^2}{n}} \right]$

Estimating $\frac{\sigma_1^2}{\sigma_2^2}$:

Estimation	Pivot Variable	Confidence Interval
μ_1, μ_2 known, estimate $\frac{\sigma_1^2}{\sigma_2^2}$	$T = \frac{S_{\mu_2}^2 \sigma_1^2}{S_{\mu_1}^2 \sigma_2^2} \sim F_{n,m}$	$\left[\frac{S_{\mu_1}^2}{S_{\mu_2}^2} \frac{1}{F_{m,n, \frac{\alpha}{2}}}, \frac{S_{\mu_1}^2}{S_{\mu_2}^2} \frac{1}{F_{m,n, 1-\frac{\alpha}{2}}} \right]$ or $\left[\frac{S_{\mu_1}^2}{S_{\mu_2}^2} F_{m,n, \frac{\alpha}{2}}, \frac{S_{\mu_1}^2}{S_{\mu_2}^2} F_{m,n, \frac{\alpha}{2}} \right]$
μ_1, μ_2 unknown, estimate $\frac{\sigma_1^2}{\sigma_2^2}$	$T = \frac{S_Y^2 \sigma_1^2}{S_X^2 \sigma_2^2} \sim F_{n-1, m-1}$	$\left[\frac{S_X^2}{S_Y^2} \frac{1}{F_{m-1, n-1, \frac{\alpha}{2}}}, \frac{S_X^2}{S_Y^2} \frac{1}{F_{m-1, n-1, 1-\frac{\alpha}{2}}} \right]$ or $\left[\frac{S_X^2}{S_Y^2} \frac{1}{F_{m-1, n-1, \frac{\alpha}{2}}}, \frac{S_X^2}{S_Y^2} F_{n-1, m-1, \frac{\alpha}{2}} \right]$

3. Non-normal population:

Estimation	Pivot Variable	Confidence Interval
Uniform Distribution: \vec{X} i.i.d. from $U(0, \theta)$	$T = \frac{X_{(n)}}{\theta} \sim U(0, 1)$	$\left[X_{(n)}, \frac{X_{(n)}}{\sqrt[n]{\alpha}} \right]$
Exponential Distribution: \vec{X} i.i.d. from $e(\lambda)$	$T = 2n\lambda\bar{X} \sim \chi_{2n}^2$	$\left[\frac{\chi_{2n, 1-\frac{\alpha}{2}}^2}{2n\bar{X}}, \frac{\chi_{2n, \frac{\alpha}{2}}^2}{2n\bar{X}} \right]$
Bernoulli Distribution: \vec{X} i.i.d. from $B(1, \theta)$	$T = \frac{\sqrt{n}(\bar{X} - \theta)}{\sqrt{\bar{X}(1 - \bar{X})}} \xrightarrow{\mathcal{L}} N(0, 1)$	$\left[\bar{X} - N_{\frac{\alpha}{2}} \sqrt{\frac{\bar{X}(1 - \bar{X})}{n}}, \bar{X} + N_{\frac{\alpha}{2}} \sqrt{\frac{\bar{X}(1 - \bar{X})}{n}} \right]$
Poisson Distribution: \vec{X} i.i.d. from $P(\lambda)$	$T = \frac{\sqrt{n}(\bar{X} - \lambda)}{\sqrt{\bar{X}}} \xrightarrow{\mathcal{L}} N(0, 1)$	$\left[\bar{X} - N_{\frac{\alpha}{2}} \sqrt{\frac{\bar{X}}{n}}, \bar{X} + N_{\frac{\alpha}{2}} \sqrt{\frac{\bar{X}}{n}} \right]$

4. General Case: Use asymptotic normality of MLE to construct CLT for large sample. MLE of θ satisfies:

$$\sqrt{n}(\hat{\theta}^* - \theta) \xrightarrow{\mathcal{L}} N(0, \frac{1}{I(\theta)})$$

where $\hat{\theta}^*$ is MLE of θ . Replace $\frac{1}{I(\theta)}$ by $\sigma^2(\hat{\theta}^*)$, then

$$T = \frac{\sqrt{n}(\hat{\theta}^* - \theta)}{\sigma(\hat{\theta}^*)} \xrightarrow{\mathcal{L}} N(0, 1)$$

confidence interval:

$$\left[\hat{\theta}^* - \frac{N_{\frac{\alpha}{2}}}{\sqrt{n}} \sigma(\hat{\theta}^*), \hat{\theta}^* + \frac{N_{\frac{\alpha}{2}}}{\sqrt{n}} \sigma(\hat{\theta}^*) \right]$$

2.3.4 Fisher Fiducial Argument*

Idea: When sample is known, we can get '**Fiducial Probability**' of θ , thus can find an interval estimation based on fiducial distribution. (Similar to the idea of MLE)

Remark: Fiducial probability (denoted as $\tilde{P}(\theta)$) is 'probability of parameter', in the case that sample is known. **Fiducial probability is different from Probability.**

Thus get

$$\tilde{P}(\hat{g}_1 \leq g(\theta) \leq \hat{g}_2) = 1 - \alpha$$

2.4 Hypothesis Testing

Hypothesis is a statement about the characteristic of population, e.g. distribution form, parameters, etc.

Mission: Use sample to test the hypothesis, i.e. judge whether population has some characteristic.

2.4.1 Basic Concepts

Parametric hypothesis testing.

For random sample $\vec{X} = (X_1, X_2, \dots, X_n) \in \mathcal{X}$ i.i.d. from $\mathcal{F} = \{f(x; \theta); \theta \in \Theta\}$

- Null Hypothesis H_0 & Alternative Hypothesis H_1 : Wonder whether a statement is true. Def. **Null Hypothesis**: $H_0 : \theta \in \Theta_0 \subset \Theta$, **a statement that we try to reject based on sample**; $H_1 : \theta \in \Theta_1 = \Theta - \Theta_0$ is **Alternative Hypothesis**.

Thus Hypothesis Testing:

$$H_0 : \theta \in \Theta_0 \longleftrightarrow H_1 : \theta \in \Theta_1$$

- Rejection Region R & Acceptance Region R^C : Judge whether to reject H_0 from sample, Def. **Rejection Region**:

$$R \subset \mathcal{X}: \text{reject } H_0 \text{ if } \vec{X} \in R$$

Acceptance Region: accept H_0 if $\vec{X} \in R^C$

- Test Function: Describe how to make a decision.

– Continuous Case:

$$\varphi(\vec{X}) = \begin{cases} 1, & \vec{X} \in R \\ 0, & \vec{X} \in R^C \end{cases}$$

i.e. $R = \{\vec{X} : \varphi(\vec{X}) = 1\}$. Where R to be determined.

– Discrete Case: Randomized Test Function

$$\varphi(\vec{X}) = \begin{cases} 1, & \vec{X} \in R - \partial R \\ r, & \vec{X} \in \partial R \\ 0, & \vec{X} \in R^C \end{cases}$$

Where R and r to be determined.

- Type I Error & Type II Error: Sample is random, possible to make a wrong judge.

– Type I Error (弃真): H_0 is true but sample falls in R , thus H_0 is rejected.

$$P(\text{type I error}) = P(\vec{X} \in R | H_0) = \alpha(\theta)$$

– Type II Error (取伪): H_0 is wrong but sample falls in R^C , thus H_0 is accepted.

$$P(\text{type II error}) = P(\vec{X} \notin R | H_1) = \beta(\theta)$$

Impossible to make probability of Type I & II Error small simultaneously, how to pick a proper test $\varphi(\vec{x})$?

Neyman-Pearson Principle: First control $\alpha \leq \alpha_0$, then take $\min \beta$.

How to determine α_0 ? Depend on specific problem.¹

¹In most cases, take $\alpha_0 = 0.05$.

	Judgement		
	Accept H_0		Reject H_0
Real Case	H_0	✓	Type I Error
	H_1	Type II Error	✓

- p -value: probability to get larger bias than observed \vec{x}_0 under H_0 .

For reject region $R = \{\vec{X} | T(\vec{X}) \geq C\}$, p -value:

$$p(\vec{x}) = P[T(\vec{X}) \geq t(\vec{x}_0) | H_0]$$

Remark: Under H_0 , the probability to get a worse result than \vec{x}_0 .

Rule: Reject H_0 if $p(\vec{x}_0) \leq \alpha_0$

- Power Function: (when H_0 is given), probability to reject H_0 by sampling.

$$\pi(\theta) = \begin{cases} P(\text{type I error}), & \theta \in \Theta_0 \\ 1 - P(\text{type II error}), & \theta \in \Theta_1 \end{cases} = \begin{cases} \alpha(\theta), & \theta \in \Theta_0 \\ 1 - \beta(\theta), & \theta \in \Theta_1 \end{cases}$$

Express as test function:

$$\pi(\theta) = E[\varphi(\vec{X}) | \theta]$$

A nice test: $\pi(\theta)$ small under H_0 , large under H_1 .

► General Steps of Hypothesis Testing:

1. Propose H_0 & H_1 .
2. Determine R (usually in the form of a statistic, e.g. $R = \{\vec{X} : T(\vec{X}) \geq c\}$).
3. Select a proper α (to determine c).
4. Sampling, get sample (as well as $t(\vec{x})$), compare with R and determine whether to reject/accept H_0

2.4.2 Hypothesis Testing of Common Distributions

For some common distribution populations, determine rejection region R under certain H_0 with confidence coefficient α .

Definition of necessary statistics see section 2.3.3.

1. Single normal population:

Condition	H_0	H_1	Testing Statistic T	Rejection Region R
σ^2 known, test μ	$\mu = \mu_0$ $\mu \leq \mu_0$ $\mu \geq \mu_0$	$\mu \neq \mu_0$ $\mu > \mu_0$ $\mu < \mu_0$	$T = \frac{\sqrt{n}(\bar{X} - \mu_0)}{\sigma} \sim N(0, 1)$	$ T > N_{\frac{\alpha}{2}}$ $T > N_{\alpha}$ $T < -N_{\alpha}$
σ^2 unknown, test μ	$\mu = \mu_0$ $\mu \leq \mu_0$ $\mu \geq \mu_0$	$\mu \neq \mu_0$ $\mu > \mu_0$ $\mu < \mu_0$	$T = \frac{\sqrt{n}(\bar{X} - \mu_0)}{S} \sim t_{n-1}$	$ T > t_{n-1, \frac{\alpha}{2}}$ $T > t_{n-1, \alpha}$ $T < -t_{n-1, \alpha}$
μ known, test σ^2	$\sigma^2 = \sigma_0^2$ $\sigma^2 \leq \sigma_0^2$ $\sigma^2 \geq \sigma_0^2$	$\sigma^2 \neq \sigma_0^2$ $\sigma^2 > \sigma_0^2$ $\sigma^2 < \sigma_0^2$	$T = \frac{nS_{\mu}^2}{\sigma_0^2} \sim \chi_n^2$	$T < \chi_{n, 1-\frac{\alpha}{2}}^2 \cup T > \chi_{n, \frac{\alpha}{2}}^2$ $T > \chi_{n, \alpha}^2$ $T < \chi_{n, 1-\alpha}^2$
μ unknown, test σ^2	$\sigma^2 = \sigma_0^2$ $\sigma^2 \leq \sigma_0^2$ $\sigma^2 \geq \sigma_0^2$	$\sigma^2 \neq \sigma_0^2$ $\sigma^2 > \sigma_0^2$ $\sigma^2 < \sigma_0^2$	$T = \frac{(n-1)S^2}{\sigma_0^2} \sim \chi_{n-1}^2$	$T < \chi_{n-1, 1-\frac{\alpha}{2}}^2 \cup T > \chi_{n-1, \frac{\alpha}{2}}^2$ $T > \chi_{n-1, \alpha}^2$ $T < \chi_{n-1, 1-\alpha}^2$

2. Double normal population:

Condition	H_0	H_1	Testing Statistic T	Rejection Region R
σ_1^2, σ_2^2 known, test $\mu_1 - \mu_2$	$\mu_1 - \mu_2 = \mu_0$ $\mu_1 - \mu_2 \leq \mu_0$ $\mu_1 - \mu_2 \geq \mu_0$	$\mu_1 - \mu_2 \neq \mu_0$ $\mu_1 - \mu_2 > \mu_0$ $\mu_1 - \mu_2 < \mu_0$	$T = \frac{\bar{X} - \bar{Y} - \mu_0}{\sqrt{\frac{\sigma_1^2}{m} + \frac{\sigma_2^2}{n}}} \sim N(0, 1)$	$ T > N_{\frac{\alpha}{2}}$ $T > N_{\alpha}$ $T < -N_{\alpha}$
σ_1^2, σ_2^2 unknown, test $\mu_1 - \mu_2$	$\mu_1 - \mu_2 = \mu_0$ $\mu_1 - \mu_2 \leq \mu_0$ $\mu_1 - \mu_2 \geq \mu_0$	$\mu_1 - \mu_2 \neq \mu_0$ $\mu_1 - \mu_2 > \mu_0$ $\mu_1 - \mu_2 < \mu_0$	$T = \frac{\bar{X} - \bar{Y} - \mu_0}{S_w} \sqrt{\frac{mn}{m+n}} \sim t_{m+n-2}$	$ T > t_{m+n-2, \frac{\alpha}{2}}$ $T > t_{m+n-2, \alpha}$ $T < -t_{m+n-2, \alpha}$
μ_1, μ_2 known, test $\frac{\sigma_1^2}{\sigma_2^2}$	$\sigma_1^2 = \sigma_2^2$ $\sigma_1^2 \geq \sigma_2^2$ $\sigma_1^2 \leq \sigma_2^2$	$\sigma_1^2 \neq \sigma_2^2$ $\sigma_1^2 < \sigma_2^2$ $\sigma_1^2 > \sigma_2^2$	$T = \frac{S_{\mu_2}^2}{S_{\mu_1}^2} \sim F_{n,m}$	$T < F_{n,m, 1-\frac{\alpha}{2}}$ $\cup T > F_{n,m, \frac{\alpha}{2}}$ $T > F_{n,m, \alpha}$ $T < F_{n,m, 1-\alpha}$
μ_1, μ_2 unknown, test $\frac{\sigma_1^2}{\sigma_2^2}$	$\sigma_1^2 = \sigma_2^2$ $\sigma_1^2 \geq \sigma_2^2$ $\sigma_1^2 \leq \sigma_2^2$	$\sigma_1^2 \neq \sigma_2^2$ $\sigma_1^2 < \sigma_2^2$ $\sigma_1^2 > \sigma_2^2$	$T = \frac{S_2^2}{S_1^2} \sim F_{n-1, m-1}$	$T < F_{n-1, m-1, 1-\frac{\alpha}{2}}$ $\cup T > F_{n-1, m-1, \frac{\alpha}{2}}$ $T > F_{n-1, m-1, \alpha}$ $T < F_{n-1, m-1, 1-\alpha}$

3. None normal population:

Condition	H_0	H_1	Testing Statistic T	Rejection Region R
\vec{X} from $B(1, p)$, test p	$p = p_0$	$p \neq p_0$	$T = \frac{\sqrt{n}(\bar{X} - p_0)}{\sqrt{p_0(1 - p_0)}} \xrightarrow{\mathcal{L}} N(0, 1)$	$ T > N_{\frac{\alpha}{2}}$
\vec{X} from $P(\lambda)$, test λ	$\lambda = \lambda_0$	$\lambda \neq \lambda_0$	$T = \frac{\sqrt{n}(\bar{X} - \lambda_0)}{\sqrt{\lambda_0}} \xrightarrow{\mathcal{L}} N(0, 1)$	$ T > N_{\frac{\alpha}{2}}$

2.4.3 Likelihood Ratio Test

Idea: To test $H_0 : \theta \in \Theta_0 \longleftrightarrow H_1 : \theta \in \Theta_1$ known \vec{x} , examine the likelihood function $L(\theta; \vec{x})$ and **compare** $L_{\theta \in \Theta_0}$ and $L_{\theta \in \Theta}$ to see the likelihood that H_0 is true.

Def. **Likelihood Ratio (LR)**:

$$\lambda(\vec{x}) = \frac{\sup_{\theta \in \Theta_0} L(\theta; \vec{x})}{\sup_{\theta \in \Theta} L(\theta; \vec{x})}$$

Reject H_0 if $\lambda(\vec{x}) < \lambda_0$. Or equivalently

Reject H_0 if $-2 \ln \lambda(\vec{x}) > C (= -2 \ln \lambda_0)$.

where λ_0 (or equivalently $C = -2 \ln \lambda_0$) satisfies:

$$E_{\Theta_0}[\varphi(\vec{X})] \leq \alpha, \quad \forall \theta \in \Theta_0$$

LR and sufficient statistic: $\lambda(\vec{x})$ can be expressed as $\lambda(\vec{x}) = \lambda^*(T(\vec{x}))$, where $T(\vec{X})$ is sufficient statistic.

► Limiting Distribution of LR: Wilks' Thm.

If $\dim \Theta = k > \dim \text{span}\{\Theta_0\} = s^2$, then under $H_0 : \theta \in \Theta_0$:

$$\Lambda_{\theta \in \Theta_0}(\vec{x}) = -2 \ln \lambda(\vec{x}) \xrightarrow{\mathcal{L}} \chi_{k-s}^2$$

2.4.4 Uniformly Most Powerful Test

Idea: Neyman-Pearson Principle: control α , find $\min \beta$. i.e. control α , find $\max \pi(\theta)$

Def. **Uniformly Most Powerful Test (UMP)** φ_{UMP} with level of significance α satisfies

$$\pi_{\text{UMP}}(\theta) \geq \pi(\theta), \quad \forall \theta \in \Theta_1$$

Neyman-Pearson Lemma: For $\vec{X} = (X_1, X_2, \dots, X_n)$ i.i.d. from $f(\vec{x}; \theta)$.

Test hypothesis $H_0 : \theta = \theta_0 \longleftrightarrow H_1 : \theta = \theta_1$. Def. test function φ as:

$$\varphi(\vec{x}) = \begin{cases} 1, & \frac{f(\vec{x}; \theta_1)}{f(\vec{x}; \theta_0)} > C \\ r, & \frac{f(\vec{x}; \theta_1)}{f(\vec{x}; \theta_0)} = C \\ 0, & \frac{f(\vec{x}; \theta_1)}{f(\vec{x}; \theta_0)} < C \end{cases} \quad (2.2)$$

Then there exists C and r such that

²Here 'dimension' refers to 'degree of freedom'.

- $E[\varphi(\vec{x})|\theta_0] = P\left(\frac{f(\vec{x};\theta_1)}{f(\vec{x};\theta_0)} > C\right) + rP\left(\frac{f(\vec{x};\theta_1)}{f(\vec{x};\theta_0)} = C\right) = \alpha$
- This φ is UMP of level of significance α

Actually kind of 1-dimensional case of LRT.

Note: UMT exist for **simple** H_0, H_1 , otherwise may not exist.

UMP and sufficient statistics: Test function $\varphi(\vec{X})$ given by eqa.2.2 is function of sufficient statistics $T(\vec{X})$, i.e. $\varphi(\vec{X}) = \varphi^*(T(\vec{X}))$.

UMP and Exponential Family: For sample $\vec{X} = (X_1, X_2, \dots, X_n)$ from exponential family:

$$f(\vec{x}; \theta) = C(\theta)h(\vec{x}) \exp\{Q(\theta)T(\vec{x})\}$$

Test single hypothesis $H_0 : \theta = \theta_0 \longleftrightarrow H_1 : \theta = \theta_1$, (where $\theta_0 < \theta_1$). If

- θ_0 is inner point of Θ
- $Q(\theta)$ monotone increase with θ

Then UMP exists, in the form of:

$$\varphi(\vec{x}) = \begin{cases} 1, & T(\vec{x}) > C \\ r, & T(\vec{x}) = C \\ 0, & T(\vec{x}) < C \end{cases} \quad (2.3)$$

where C and r satisfies $E[\varphi(\vec{x})|\theta_0] = \alpha$.

Note: or take $Q(\theta)$ mono decreased, then in eqa.2.3, take opposite inequality operators.

► General Steps of UMP:

1. Find a point $\theta_0 \in \Theta_0$ and a point $\theta_1 \in \Theta_1$. (Note: **one** point)
2. Construct test function in the form of eqa.2.2, use $E[\varphi(\vec{x})|\theta_0] = \alpha$ to determine C and r .
3. Get R and $\varphi(\vec{x})$.
4. If φ does **not** depend on θ_1 , then H_1 can be generalized to $H_1 : \theta \in \Theta_1$.
5. If φ satisfies $E_{\theta \in \Theta_0}(\varphi) \leq \alpha$, then H_0 can be generalized to $H_0 : \theta \in \Theta_0$.

2.4.5 Duality of Hypothesis Testing and Interval Estimation

- Thm.: $\forall \theta_0 \in \Theta$ there exists hypothesis testing $H_0 : \theta = \theta_0 \longleftrightarrow H_1 : \theta \neq \theta_0$ of level α with rejection region R_{θ_0} . Then

$$C(\vec{X}) = \{\theta : \vec{X} \in R_{\theta}^C\}$$

is a $1 - \alpha$ confidence region for θ

- Thm.: $C(\vec{X})$ is a $1 - \alpha$ confidence region for θ . Then $\forall \theta_0 \in C(\vec{X})$, the rejection region of hypothesis testing $H_0 : \theta = \theta_0 \longleftrightarrow H_1 : \theta \neq \theta_0$ of level α satisfies

$$R_{\theta_0}^C = \{\vec{X} : \theta_0 \in C(\vec{X})\}$$

► Idea:

$$H_0 : \theta = \theta_0 \longleftrightarrow H_1 : \theta \neq \theta_0$$

$$\downarrow$$

$$P(R^C(\vec{X})|H_0) = P(R^C(\vec{X})|\theta_0) = 1 - \alpha$$

$$\downarrow$$

$$\text{Confidence Interval: } \theta_0 \in R^C(\vec{X})$$

Similar for Confidence Limit and One-Sided Testing.

2.4.6 Introduction to Non-Parametric Hypothesis Testing

Motivation: Usually distribution form unknown, cannot use parametric hypothesis testing.

Useful Method:

- Sign Test: Used for paired comparison $\vec{X} = (X_1, X_2, \dots, X_n)$, $\vec{Y} = (Y_1, Y_2, \dots, Y_n)$.

Take $Z_i = Y_i - X_i$ i.i.d., denote $E(Z) = \mu$. Test $H_0 : \mu = 0 \longleftrightarrow H_1 : \mu \neq 0$.

Denote $n_+ = \#(\text{positive } Z_i)$ and $n_- = \#(\text{negative } Z_i)$, $n_0 = n_+ + n_-$. Then $n_+ \sim B(n_0, \theta)$, test $H_0 : \theta = \frac{1}{2} \longleftrightarrow H_1 : \theta \neq \frac{1}{2}$

Then use Binomial Testing or large sample CLT Normal Testing.

Remark:

- Also can test $H_0 : \theta \leq \frac{1}{2} \longleftrightarrow H_1 : \theta > \frac{1}{2}$
- Drawback: ignores magnitudes.

- Wilcoxon Signed Rank Sum Test: Improvement of Sign Test. Base on order statistics.

Order Statistics of Z_i : $Z_{(1)} < Z_{(2)} < \dots < Z_{(n)}$, where each $Z_{(j)}$ corresponds to some Z_i , denote as $Z_i = Z_{(R_i)}$, then R_i is the rank of Z_i .³

Def. $\vec{R} = (R_1, R_2, \dots, R_n)$ is **Rank Statistics** of (Z_1, Z_2, \dots, Z_n)

Def. **Sum of Wilcoxon Signed Rank**:

$$W^+ = \sum_{i=1}^{n_0} R_i I_{Z_i > 0}$$

³If some X_i, X_j, \dots equal, then take same rank $R = \text{mean}\{R_i, R_j, \dots\}$.

Distribution of W^+ is complex. E and var of W^+ under H_0 :

$$E(W^+) = \frac{n_0(n_0 + 1)}{4} \quad var(W^+) = \frac{n_0(n_0 + 1)(2n_0 + 1)}{24}$$

Usually consider large sample CLT, construct normal approximation:

$$T = \frac{W^+ - E(W^+)}{\sqrt{var(W^+)}} \xrightarrow{\mathcal{L}} N(0, 1)$$

Rejection Region: $R = \{|T| > N_{\frac{\alpha}{2}}\}$

- Wilcoxon Two-Sample Rank Sum Test: Used for two independent sample comparison.

Assume $\vec{X} = (X_1, \dots, X_m)$ i.i.d. $\sim f(x)$; $\vec{Y} = (Y_1, \dots, Y_n)$ i.i.d. $\sim f(x - \theta)$, test $H_0 : \theta = 0 \longleftrightarrow H_1 : \theta \neq 0$.

Rank X_i and Y_i as:

$$Z_1 \leq Z_2 \leq \dots \leq Z_{m+n}$$

in which denote rank of Y_i as R_i , and def. **Wilcoxon two-sample rank sum**:

$$W = \sum_{i=1}^n R_i$$

E and var of W under H_0 :

$$E(W) = \frac{n(m+n+1)}{2} \quad var(W) = \frac{mn(n+m+1)}{12}$$

Use large sample approximation, construct CLT:

$$T = \frac{W - E(W)}{\sqrt{var(W)}} \xrightarrow{\mathcal{L}} N(0, 1)$$

- Goodness-of-Fit Test: For $\vec{X} = (X_1, X_2, \dots, X_n)$ i.i.d. from some certain population X . Test $H_0 : X \sim F(x)$.

where F is theoretical distribution, can be either parametric or non-parametric.

Idea: Define some *quantity* $D = D(X_1, \dots, X_n; F)$ to measure the difference between F and sample. And def. *Goodness-of-fit* when observed value of D (say d_0) is given:

$$p(d_0) = P(D \geq d_0 | H_0)$$

Goodness-of-Fit Test: Reject H_0 if $p(d_0) < \alpha$.

Pearson χ^2 Test: Usually used for discrete case.

Test $H_0 : P(X_i = a_i) = p_i, i = 1, 2, \dots, r$. Denote $\#(X_j = a_i) = \nu_i$, take D as:

$$K_n = K_n(X_1, \dots, X_n; F) = \sum_{i=1}^r \frac{(\nu_i - np_i)^2}{np_i} \quad (2.4)$$

Pearson Thm.: For K_n defined as eqa.2.4, then under H_0 :

$$K_n \xrightarrow{\mathcal{L}} \chi_{r-1-s}^2$$

Here s is number of unknown parameter, $r - 1 - s$ is the degree of freedom.

Note:

- a_i must **not** depend on sample.
- For continuous case, construct division:

$$\mathbb{R} \rightarrow (-\infty, a_1, a_2, \dots, a_{r-1}, \infty = a_r)$$

and test $H_0 : P(X \in I_j) = p_j$

Criterion: Pick proper interval so that np_i and ν_i both ≥ 5 .

- Contingency Table Independence & Homogeneity Test

- Independence Test:

Test a two-parameter sample and to see whether these two parameters(features) are independent.

Denote $Z = (X, Y)$ are some 'level' of sample, n_{ij} is number of sample with level (i, j)

Contingency Table:

X \ Y	Y					Σ
	1	...	j	...	s	
1	n_{11}	...	n_{1j}	...	n_{1s}	$n_{1\cdot}$
\vdots	\vdots	\ddots	\vdots	\ddots	\vdots	\vdots
i	n_{i1}	...	n_{ij}	...	n_{is}	$n_{i\cdot}$
\vdots	\vdots	\ddots	\vdots	\ddots	\vdots	\vdots
r	n_{r1}	...	n_{rj}	...	n_{rs}	$n_{r\cdot}$
Σ	$n_{\cdot 1}$...	$n_{\cdot j}$...	$n_{\cdot s}$	n

Test $H_0 : X \& Y$ are independent. i.e. $H_0 : P(X = i, Y = j) = P(X = i)P(Y = j) = p_i p_j$.

Construct χ^2 test statistic:

$$K_n = \sum_{i=1}^r \sum_{j=1}^s \frac{[n_{ij} - n(\frac{n_{i\cdot}}{n})(\frac{n_{\cdot j}}{n})]^2}{n(\frac{n_{i\cdot}}{n})(\frac{n_{\cdot j}}{n})} = n \left(\sum_{i=1}^r \sum_{j=1}^s \frac{n_{ij}^2}{n_{i\cdot} n_{\cdot j}} - 1 \right) \quad (2.5)$$

Then under H_0 , $K_n \xrightarrow{\mathcal{L}} \chi_{rs-1-(r+s-2)}^2 = \chi_{(r-1)(s-1)}^2$

Reject H_0 if $p(k_0) = P(K_n \geq k_0) < \alpha$

- Homogeneity Test:

Test R groups of sample with category rank, to see whether these groups has similar rank distribution.

Group \ Category	Category					Σ
	Category 1	...	Category j	...	Category C	
Group 1	n_{11}	...	n_{1j}	...	n_{1C}	$n_{1\cdot}$
\vdots	\vdots	\ddots	\vdots	\ddots	\vdots	\vdots
Group i	n_{i1}	...	n_{ij}	...	n_{iC}	$n_{i\cdot}$
\vdots	\vdots	\ddots	\vdots	\ddots	\vdots	\vdots
Group R	n_{R1}	...	n_{Rj}	...	n_{RC}	$n_{R\cdot}$
Σ	$n_{\cdot 1}$...	$n_{\cdot j}$...	$n_{\cdot C}$	n

Denote $P(\text{Category } j | \text{Group } i) = p_{ij}$. Test $H_0 : p_{ij} = p_j, \forall 1 \leq i \leq R$.

Construct χ^2 test statistic:

$$D = \sum_{i=1}^R \sum_{j=1}^C \frac{[n_{ij} - n(\frac{n_{i\cdot}}{n})(\frac{n_{\cdot j}}{n})]^2}{n(\frac{n_{i\cdot}}{n})(\frac{n_{\cdot j}}{n})} = n \left(\sum_{i=1}^R \sum_{j=1}^C \frac{n_{ij}^2}{n_{i\cdot} n_{\cdot j}} - 1 \right) \quad (2.6)$$

Then under H_0 , $D \xrightarrow{\mathcal{L}} \chi_{R(C-1)-(C-1)}^2 = \chi_{(R-1)(C-1)}^2$

- Test of Normality: normality is a good & useful assumption.

For $\vec{Y} = (Y_1, Y_2, \dots, Y_n)$,

Test H_0 : exists μ & σ^2 such that Y_i i.i.d. $\sim N(\mu, \sigma^2)$.

- Kolmogorov-Smirnov Test: Assume \vec{X} form population CDF $F(x)$, test $H_0 : F(x) = F_0(x)$ (where can take $F_0 = \Phi$ or some other known CDF).

use $F_n(x)$ (as defined in eqa.2.1) as approx. to $F(x)$, test

$$D_n = \sum_{-\infty < x < +\infty} |F_n(x) - F_0(x)|$$

Reject H_0 if $D_n > c$

or use goodness-of-fit: denote observed value of D_n as d_n . Reject H_0 if

$$p(d_n) = P(D_n > d_n | H_0) < \alpha$$

- Shapiro-Wilk Test:

Test H_0 : exists μ & σ^2 such that X_i i.i.d. $\sim N(\mu, \sigma^2)$.

Denote $Y_{(i)} = \frac{X_{(i)} - \mu}{\sigma}$, $m_i = E(Y_{(i)})$

Under H_0 , $(X_{(i)}, m_i)$ falls close to straight line. Test Statistic: Correlation

$$R^2 = \frac{(\sum_{i=1}^n (X_{(i)} - \bar{X})(m_i - \bar{m}))^2}{\sum_{i=1}^n (X_{(i)} - \bar{X})^2 \sum_{i=1}^n (m_i - \bar{m})^2}$$

Reject H_0 if $R^2 < c$

Shapiro-Wilk correction:

$$W = \frac{\left(\sum_{i=1}^{[n/2]} a_i (X_{(n+1-i)} - X_{(i)}) \right)^2}{\sum_{i=1}^n (X_{(i)} - \bar{X})^2}$$

► Summary: Useful Non-Parameter Hypothesis Testing.

