

# Climate change of sudden stratospheric warmings

# Structure

1 Introduction: The Polar Vortex

2 Sudden Stratospheric Warmings

3 Predicted Changes In Climate Models

3.1 General changes in the stratosphere

3.2 Changes in SSW frequency

3.3 Changes in other SSW characteristics

3.4 Reasons for predicted changes and uncertainties



3.5 Change in SSW frequency in the Southern hemisphere

4 Conclusion

# The stratospheric Polar Vortex

## Temperatures in the lower stratosphere:

- Minimum at equator
- Maximum at pole in the summer hemisphere
- Maximum at 45° in the winter hemisphere, colder on the pole
- Thermal wind balance: easterly winds in summer  
strong westerly winds in winter




# Sudden Stratospheric Warmings

## Forming:

- Planetary wave propagates from troposphere upward
- Wave dissipates and easterly momentum is released
- Winds weaken, abrupt warming due to thermal wind balance
- Consequence: polar vortex displacement (moving away from pole to lower latitudes), polar vortex split (polar vortex is split into two separate vortices)
- Mainly in Northern Hemisphere due to orography and land-sea distribution
- About 6 SSWs/decade in NH and only 1 recorded SSW in SH in about 60 years of observations

Disruption of stratospheric polar vortex in early January 2021



# Sudden Stratospheric Warmings

Classification:

**Major Warming:**

Reversal of zonal mean zonal wind in 10hPa at 60°(N)

WMO criterion:



additional reversal of temperature gradient between 60°N and pole

Minor warming:

Significant warming and weakening of wind without wind reversal

Final warming:


Polar vortex winds returning to easterly winds for the entire summer




© Baldwin et al. 2020

# Impacts on Stratosphere

- Colder upper stratosphere due to reversed meridional circulation and resulting upward movement
- Opposite anomalies in midlatitude and tropical stratosphere (not so pronounced)
- Changed circulation patterns lead to increased total ozone column over the poles, especially well visible during southern hemisphere SSW in 2002




Shading shows zonal mean temperature anomalies 0-30 days after SSW, arrows indicate meridional circulation



© Baldwin et al. 2020


# Stratosphere-Troposphere Coupling



- Impact on the troposphere can last for up to 2 months
- Anomalies propagate downward
- About 2/3 of SSW have measurable impact on the troposphere






# Impacts on Surface



- Higher sea level pressure over North Pole
- Lower pressure over mid-latitudes, especially Europe
- Results in a negative Northern Annular Mode and North Atlantic Oscillation
- Colder over Northern Eurasia, warmer over Eastern Canada and Greenland
- More precipitation over Southern and Western Europe, less over Northern Atlantic, polar regions, Northern Europe



# Predicted changes in climate models





© Manzini et al. 2014

General changes (NH winter, CMIP5, RCP 8.5):

- Strengthening of lower to midlatitude zonal winds
- No clear sign for change of wind speed in higher latitude stratosphere
- Temperature increase in the troposphere, at the surface especially in the Arctic, at the equator especially in upper troposphere
- Temperature decrease in stratosphere
- Lower sea level pressure over Arctic, higher pressure over midlatitudes



# Changes in polar vortex wind speed



© Simpson et al. 2018


- Large spread in climate models(CMIP5, RCP8.5)
- models with significant strengthening and weakening of polar vortex winds in CMIP5 ensemble, but majority of models show no significant change

# Changes in SSW frequency




- CMIP5: Large inter-model spread from almost no SSW to 1 per year
- Majority of models show less SSW than observed
- No clear signal for future changes: most models predict slight increase in frequency, but some also show a decrease

# Changes in SSW frequency



- CMIP6: large inter-model spread remains, generally closer to reanalysis
- Sign and magnitude of change (ensemble mean) similar compared to CMIP5 for RCP 4.5
- In RCP8.5 (SSP585) CMIP6 frequency decreases compared to RCP4.5, and no change compared to historical simulation

# Changes in SSW frequency





© Ayarzagüena et al. 2020

- CMIP 6, quadrupling of CO<sub>2</sub>, simulation using 12 models
- 4models significant decrease, 4models significant increase, 4models no significant change → majority of models predict significant change but no consensus on direction


# Changes in SSW frequency

- CMIP6, 1% CO<sub>2</sub> increase per year
  - 6/11 models show significant change in SSW frequency
  - Doubling of CO<sub>2</sub> between decade 6-7 (2060-2070 in RCP 8.5)
- change in SSW frequency unlikely before that






# Changes in seasonality of SSW



- Reanalysis: most SSW in mid-winter(Jan+Feb)
- Both CMIP5&6 simulate most SSW in March
- Significant increase shifts from Jan-Mar in CMIP5 to Dec-Feb in CMIP6
- No clear conclusion possible because of differences between models and differences compared to reanalysis

# Changes in tropospheric response

- CMIP5&6 simulate tropospheric response close to reanalysis
- No significant changes predicted in RCP4.5 or RCP8.5




Evolution of zonal mean zonal wind anomalies after SSW; contours show values and shading shows statistical significance (light:90%, dark:95%)

© Rao et al. 2021

# Changes in surface pressure response

- CMIP6: almost all models show the same pattern of pressure anomalies as in the reanalysis
- Magnitude of change varies between models
- High pressure anomalies in Northern Pacific are not represented in any model




Sea level pressure anomalies as contours and temperature anomalies as shading 15-60 after a SSW, green stippling shows significant differences of pressure to reanalysis

© Ayarzagüena et al. 2020

# Changes in surface pressure response

- Most models show no significant changes
- Some show stronger negative Northern Annular Mode (higher pressure over pole, lower over midlatitudes)
- Some show significant negative pressure anomalies over the Northern Pacific




Sea level pressure anomalies in preindustrial control simulations as contours and changes in 4xCO<sub>2</sub>-scenario as shading  
green stippling shows significant differences to preindustrial control simulation (95%)

© Ayarzagüena et al. 2020

# Changes in seasonal cycle of the polar vortex



- Historical models (green) simulate earlier vortex formation than in reanalysis
- All models show a longer persisting polar vortex with 4xCO<sub>2</sub> (brown)





# Changes in seasonal cycle of the polar vortex






Models with longest persisting vortex simulate less SSW in future





# Influence of polar vortex wind speed




- Models that simulate significantly less SSW show much stronger polar vortex winds in the future
- Change in SSW frequency not necessarily tied to mean wind speed: see models with increased SSW frequency



© Ayarzagüena et al. 2020



# Reasons for uncertainties in the models

- Negative correlation between SSW frequency and climate sensitivity of models (-0.33, not significant)
- Significant negative correlation between SSW frequency and change in lower tropospheric temperature (850hPa) over North pole
- Significant positive correlation between SSW frequency and change in temperature gradient in 850hPa
- Probably due to effects on wave propagation but still not well understood




# Role of sea ice in Barents and Kara-Sea

- Changes in sea ice concentration could alter meridional heat flux and with that wave propagation
- Non-linear response of polar vortex wind speeds to increasing temperatures depending on sea ice concentration
- With decreasing ice concentration wind speeds decrease
- Once Barents and Kara-Sea are ice free wind speed is increasing again



# SSWs in the Southern hemisphere

- Especially interesting for ozone hole projection and surface pressure response (could lead to more wildfires in Southern hemisphere)
- Historical simulation: 1SSW every 59 years (recorded: 1 in 63 years of observations)
- With increased CO<sub>2</sub> forcing (1120ppm): 1SSW every 883 years and significant strengthening of polar vortex



© Jucker et al. 2021

# Conclusion

- No consensus in models on change in SSW frequency in Northern hemisphere, most likely a slight increase considering multimodel mean
- Large spread in models probably due to multiple factors (non-linear response to sea ice loss, arctic amplification, climate sensitivity)
- No significant changes in stratosphere-troposphere coupling and surface pressure response predicted
- No clear sign for changes in seasonality of SSWs: climate models biased to more SSWs in late winter
- Much clearer for Southern hemisphere: significant decrease in SSW frequency and strengthening of the stratospheric polar vortex that will probably lead to a longer persisting ozone hole

# References

- Ayarzagüena et al., Uncertainty in the Response of Sudden Stratospheric Warmings and Stratosphere-Troposphere Coupling to Quadrupled CO<sub>2</sub> Concentrations in CMIP6 Models, DOI: 10.1029/2019JD032345, 2020
- Jucker et al., How Frequent Are Antarctic Sudden Stratospheric Warmings in Present and Future Climate, DOI: 10.1029/2021GL093215, 2021
- Kretschmer et al., The role of Barents-Kara sea ice loss in projected polar vortex changes, DOI: 10.5194/wcd-1-715-2020, 2020
- Manzini et al., Northern winter climate change: Assessment of uncertainty in CMIP5 projections related to stratosphere-troposphere coupling, DOI: 10.1002/2013JD021403, 2014
- Rao et al., CMIP5/6 models project little change in the statistical characteristics of sudden stratospheric warmings in the 21<sup>st</sup> century, DOI: 10.1088/1748-9326/abd4fe, 2021
- Simpson et al., The Downward influence of Uncertainty in the Northern Hemisphere Stratospheric Polar Vortex Response to Climate Change, DOI: 10.1175/JCLI-D-18-0041.1, 2018
- Baldwin et al., Sudden Stratospheric Warmings, DOI:10.1029/2020RG000708, 2020