

NCERT 9.7.1.5

This Question is from ncert class 9 chapter 7.Triangles

1. Line l is the bisector of an angle $\angle A$ and B is any point on l . BP and BQ are perpendiculars from B to the arms of $\angle A$. Show that:

(a) $\triangle APB \cong \triangle AQB$

(b) $BP = BQ$ or B is equidistant from the arms of $\angle A$.




Figure 1: $\triangle AQB$ and $\triangle APB$

solution:

Given l is the angular bisector of $\angle QAP$, so $\angle QAB = \angle PAB$ and

BP, BQ are perpendicular bisectors so angle $= 90^\circ$. let

$$\angle QAB = \angle PAB = \theta = 30^\circ \quad (1)$$

$$AP = r = 4 \quad (2)$$

$$AB = 5 \quad (3)$$

$$\mathbf{A} = \begin{pmatrix} 0 \\ 0 \end{pmatrix}, \mathbf{B} = \begin{pmatrix} 5 \\ 0 \end{pmatrix}, \mathbf{P} = \begin{pmatrix} r \cos(\theta) \\ -r \sin(\theta) \end{pmatrix}, \mathbf{Q} = \begin{pmatrix} r \cos(\theta) \\ r \sin(\theta) \end{pmatrix} \quad (4)$$

$$(5)$$

1. (a) By the definition of AAS (Angle Angle Side) congruency rule ,If two triangles have two equal angles and a pair of corresponding sides are equal then the two triangles said to be congruent. Here

$$\angle QAB = \angle PAB = 30^\circ \quad (l \text{ is angular bisector of } \angle QAP) \quad (6)$$

$$\angle AQB = \angle APB = 90^\circ \quad (BP \text{ and } BQ \text{ are perpendicular bisectors})$$

$$(7)$$

$$AB = 5 \quad (\text{common side for two triangles}) \quad (8)$$

\therefore By AAS congruency rule $\triangle APB \cong \triangle AQB$

(b)

$$\|\mathbf{B} - \mathbf{Q}\| = \sqrt{(\mathbf{B} - \mathbf{Q})^\top (\mathbf{B} - \mathbf{Q})} \quad (9)$$

$$\mathbf{B} - \mathbf{Q} = \begin{pmatrix} 5 \\ 0 \end{pmatrix} - \begin{pmatrix} r \cos(\theta) \\ r \sin(\theta) \end{pmatrix} \quad (10)$$

$$\mathbf{B} - \mathbf{Q} = \begin{pmatrix} 5 \\ 0 \end{pmatrix} - \begin{pmatrix} 3.464101 \\ 2 \end{pmatrix} \quad (11)$$

$$\mathbf{B} - \mathbf{Q} = \begin{pmatrix} 1.535899 \\ -2 \end{pmatrix} \quad (12)$$

$$(\mathbf{B} - \mathbf{Q})^\top = \begin{pmatrix} 1.535899 \\ -2 \end{pmatrix}^\top = \begin{pmatrix} 1.535899 & -2 \end{pmatrix} \quad (13)$$

$$(\mathbf{B} - \mathbf{Q})^\top (\mathbf{B} - \mathbf{Q}) = \begin{pmatrix} 1.535899 & -2 \end{pmatrix} \begin{pmatrix} 1.535899 \\ -2 \end{pmatrix} \quad (14)$$

$$= 2.3589857 + 4 \quad (15)$$

$$= 6.3589857 \quad (16)$$

$$\sqrt{(\mathbf{B} - \mathbf{Q})^\top (\mathbf{B} - \mathbf{Q})} = \sqrt{6.3589857} \quad (17)$$

$$\implies \|\mathbf{B} - \mathbf{Q}\| = 2.5217029 \quad (18)$$

$$\|\mathbf{B} - \mathbf{P}\| = \sqrt{(\mathbf{B} - \mathbf{P})^\top (\mathbf{B} - \mathbf{P})} \quad (19)$$

$$\mathbf{B} - \mathbf{P} = \begin{pmatrix} 5 \\ 0 \end{pmatrix} - \begin{pmatrix} r \cos(\theta) \\ -r \sin(\theta) \end{pmatrix} \quad (20)$$

$$\mathbf{B} - \mathbf{P} = \begin{pmatrix} 5 \\ 0 \end{pmatrix} - \begin{pmatrix} 3.464101 \\ -2 \end{pmatrix} \quad (21)$$

$$\mathbf{B} - \mathbf{P} = \begin{pmatrix} 1.535899 \\ 2 \end{pmatrix} \quad (22)$$

$$(\mathbf{B} - \mathbf{P})^\top = \begin{pmatrix} 1.535899 \\ 2 \end{pmatrix}^\top = \begin{pmatrix} 1.535899 & 2 \end{pmatrix} \quad (23)$$

$$(\mathbf{B} - \mathbf{P})^\top (\mathbf{B} - \mathbf{P}) = \begin{pmatrix} 1.535899 & 2 \end{pmatrix} \begin{pmatrix} 1.535899 \\ 2 \end{pmatrix} \quad (24)$$

$$= 2.3589857 + 4 \quad (25)$$

$$= 6.3589857 \quad (26)$$

$$\sqrt{(\mathbf{B} - \mathbf{P})^\top (\mathbf{B} - \mathbf{P})} = \sqrt{6.3589857} \quad (27)$$

$$\implies \|\mathbf{B} - \mathbf{P}\| = 2.5217029 \quad (28)$$

$\therefore BP = BQ$ Hence proved




Figure 2: $\triangle APB$ and $\triangle AQB$