

Contents

Chapter 1

Triangle

Consider a triangle with vertices

$$\mathbf{A} = \begin{pmatrix} -3 \\ -5 \end{pmatrix}, \mathbf{B} = \begin{pmatrix} 3 \\ -5 \end{pmatrix}, \mathbf{c} = \begin{pmatrix} -4 \\ -3 \end{pmatrix}, \quad (1.1)$$

1.1. Vectors

1.1.1. The direction vector of AB is defined as

$$\mathbf{B} - \mathbf{A} \quad (1.1.1.1)$$

Consider a triangle with vertices

$$\mathbf{A} = \begin{pmatrix} -3 \\ -5 \end{pmatrix} \quad (1.1.1.2)$$

$$\mathbf{B} = \begin{pmatrix} 3 \\ -5 \end{pmatrix} \quad (1.1.1.3)$$

$$\mathbf{C} = \begin{pmatrix} -4 \\ -3 \end{pmatrix} \quad (1.1.1.4)$$

The Direction Vector of AB is defined as

$$\mathbf{B} - \mathbf{A} \quad (1.1.1.5)$$

Question 1.1.1 : Find the Direction Vectors of AB, BC, CA .

Solution:

(a) The Direction vector of AB is

$$= \mathbf{B} - \mathbf{A} \quad (1.1.1.6)$$

$$= \begin{pmatrix} 3 - (-3) \\ -5 - (-5) \end{pmatrix} \quad (1.1.1.7)$$

$$= \begin{pmatrix} 6 \\ 0 \end{pmatrix} \quad (1.1.1.8)$$

(b) The Direction vector of BC

$$= \mathbf{C} - \mathbf{B} \quad (1.1.1.9)$$

$$= \begin{pmatrix} -4 - (3) \\ -3 - (-5) \end{pmatrix} \quad (1.1.1.10)$$

$$= \begin{pmatrix} -7 \\ 2 \end{pmatrix} \quad (1.1.1.11)$$

(c) The Direction vector of CA

$$= \mathbf{A} - \mathbf{C} \quad (1.1.1.12)$$

$$= \begin{pmatrix} -3 - (-4) \\ -5 - (-3) \end{pmatrix} \quad (1.1.1.13)$$

$$= \begin{pmatrix} 1 \\ -2 \end{pmatrix} \quad (1.1.1.14)$$

1.1.2. The length of side BC is

$$\|\mathbf{B} - \mathbf{A}\| \triangleq \sqrt{(\mathbf{B} - \mathbf{A})^\top \mathbf{B} - \mathbf{A}} \quad (1.1.2.1)$$

where

$$\mathbf{A}^\top \triangleq \begin{pmatrix} -3 & -5 \end{pmatrix} \quad (1.1.2.2)$$

Question 1.1.2 : Find the length of side AB, BC, CA.

Solution: Solving for BC Given,

$$\mathbf{A} = \begin{pmatrix} -3 \\ -5 \end{pmatrix}, \mathbf{B} = \begin{pmatrix} 3 \\ -5 \end{pmatrix}, \mathbf{C} = \begin{pmatrix} -4 \\ -3 \end{pmatrix} \quad (1.1.2.3)$$

$$\|\mathbf{B} - \mathbf{C}\| = \sqrt{(\mathbf{B} - \mathbf{C})^\top (\mathbf{B} - \mathbf{C})} \quad (1.1.2.4)$$

$$\mathbf{B} - \mathbf{C} = \begin{pmatrix} 3 \\ -5 \end{pmatrix} - \begin{pmatrix} -4 \\ -3 \end{pmatrix} \quad (1.1.2.5)$$

$$\mathbf{B} - \mathbf{C} = \begin{pmatrix} 7 \\ -2 \end{pmatrix} \quad (1.1.2.6)$$

$$(\mathbf{B} - \mathbf{C})^\top = \begin{pmatrix} 7 \\ -2 \end{pmatrix}^\top = \begin{pmatrix} 7 & -2 \end{pmatrix} \quad (1.1.2.7)$$

$$(\mathbf{B} - \mathbf{C})^\top (\mathbf{B} - \mathbf{C}) = \begin{pmatrix} 7 & -2 \end{pmatrix} \begin{pmatrix} 7 \\ -2 \end{pmatrix} \quad (1.1.2.8)$$

$$= 49 + 4 \quad (1.1.2.9)$$

$$= 53 \quad (1.1.2.10)$$

$$\sqrt{(\mathbf{B} - \mathbf{C})^\top (\mathbf{B} - \mathbf{C})} = \sqrt{53} \quad (1.1.2.11)$$

$$\Rightarrow \|\mathbf{B} - \mathbf{C}\| = \sqrt{53} \quad (1.1.2.12)$$

Solving for AB Given,

$$\|\mathbf{A} - \mathbf{B}\| = \sqrt{(\mathbf{A} - \mathbf{B})^\top (\mathbf{A} - \mathbf{B})} \quad (1.1.2.13)$$

$$\mathbf{A} - \mathbf{B} = \begin{pmatrix} -3 \\ -5 \end{pmatrix} - \begin{pmatrix} 3 \\ -5 \end{pmatrix} \quad (1.1.2.14)$$

$$\mathbf{A} - \mathbf{B} = \begin{pmatrix} -6 \\ 0 \end{pmatrix} \quad (1.1.2.15)$$

$$(\mathbf{A} - \mathbf{B})^\top = \begin{pmatrix} -6 \\ 0 \end{pmatrix}^\top = \begin{pmatrix} -6 & 0 \end{pmatrix} \quad (1.1.2.16)$$

$$(\mathbf{A} - \mathbf{B})^\top (\mathbf{A} - \mathbf{B}) = \begin{pmatrix} -6 & 0 \end{pmatrix} \begin{pmatrix} -6 \\ 0 \end{pmatrix} \quad (1.1.2.17)$$

$$= 36 + 0 \quad (1.1.2.18)$$

$$= 36 \quad (1.1.2.19)$$

$$\sqrt{(\mathbf{A} - \mathbf{B})^\top (\mathbf{A} - \mathbf{B})} = \sqrt{36} \quad (1.1.2.20)$$

$$\Rightarrow \|\mathbf{A} - \mathbf{B}\| = \sqrt{36} \quad (1.1.2.21)$$

Solving for CA Given,

$$\|\mathbf{C} - \mathbf{A}\| = \sqrt{(\mathbf{C} - \mathbf{A})^\top (\mathbf{C} - \mathbf{A})} \quad (1.1.2.22)$$

$$\mathbf{C} - \mathbf{A} = \begin{pmatrix} -4 \\ -3 \end{pmatrix} - \begin{pmatrix} -3 \\ -5 \end{pmatrix} \quad (1.1.2.23)$$

$$\mathbf{C} - \mathbf{A} = \begin{pmatrix} -1 \\ 2 \end{pmatrix} \quad (1.1.2.24)$$

$$(\mathbf{C} - \mathbf{A})^\top = \begin{pmatrix} -1 \\ 2 \end{pmatrix}^\top = \begin{pmatrix} -1 & 2 \end{pmatrix} \quad (1.1.2.25)$$

$$(\mathbf{C} - \mathbf{A})^\top (\mathbf{C} - \mathbf{A}) = \begin{pmatrix} -1 & 2 \end{pmatrix} \begin{pmatrix} -1 \\ 2 \end{pmatrix} \quad (1.1.2.26)$$

$$= 1 + 4 \quad (1.1.2.27)$$

$$= 5 \quad (1.1.2.28)$$

$$\sqrt{(\mathbf{C} - \mathbf{A})^\top (\mathbf{C} - \mathbf{A})} = \sqrt{5} \quad (1.1.2.29)$$

$$\Rightarrow \|\mathbf{C} - \mathbf{A}\| = \sqrt{5} \quad (1.1.2.30)$$

1.1.3. Points $\mathbf{A}, \mathbf{B}, \mathbf{C}$ are defined to be collinear if

$$\text{rank} \begin{pmatrix} 1 & 1 & 1 \\ \mathbf{A} & \mathbf{B} & \mathbf{C} \end{pmatrix} = 2 \quad (1.1.3.1)$$

Are the given points in (??) collinear?

Question 1.1.3 : Check the collinearity of $\mathbf{A}, \mathbf{B}, \mathbf{C}$

Solution: Given that,

$$\mathbf{A} = \begin{pmatrix} -3 \\ -5 \end{pmatrix} \quad \mathbf{B} = \begin{pmatrix} 3 \\ -5 \end{pmatrix} \quad \mathbf{C} = \begin{pmatrix} -4 \\ -3 \end{pmatrix} \quad (1.1.3.2)$$

Given that $\mathbf{A}, \mathbf{B}, \mathbf{C}$ are collinear if

$$\text{rank} \begin{pmatrix} 1 & 1 & 1 \\ \mathbf{A} & \mathbf{B} & \mathbf{C} \end{pmatrix} < 3 \quad (1.1.3.3)$$

Let

$$\mathbf{R} = \begin{pmatrix} 1 & 1 & 1 \\ -3 & 3 & -4 \\ -5 & -5 & -3 \end{pmatrix} \quad (1.1.3.4)$$

The matrix \mathbf{R} can be row reduced as follows,

$$\begin{pmatrix} 1 & 1 & 1 \\ -3 & 3 & -4 \\ -5 & -5 & -3 \end{pmatrix} \xleftrightarrow{R_3 \leftarrow R_3 + 5R_1} \begin{pmatrix} 1 & 1 & 1 \\ -3 & 3 & -4 \\ 0 & 0 & 2 \end{pmatrix} \quad (1.1.3.5)$$

$$\xleftrightarrow{R_2 \leftarrow R_2 + 3R_1} \begin{pmatrix} 1 & 1 & 1 \\ 0 & 6 & -1 \\ 0 & 0 & 2 \end{pmatrix} \quad (1.1.3.6)$$

There are no zero rows. So,

$$\text{rank} \begin{pmatrix} 1 & 1 & 1 \\ \mathbf{A} & \mathbf{B} & \mathbf{C} \end{pmatrix} = 3 \quad (1.1.3.7)$$

Hence, from (??) the points $\mathbf{A}, \mathbf{B}, \mathbf{C}$ are not collinear.

From Fig. ??, We can see that $\mathbf{A}, \mathbf{B}, \mathbf{C}$ are not collinear .




Figure 1.1: $\mathbf{A}, \mathbf{B}, \mathbf{C}$ plot

1.1.4. The parametric form of the equation of AB is

$$\mathbf{x} = \mathbf{A} + k\mathbf{m} \quad (1.1.4.1)$$

where

$$\mathbf{m} = \mathbf{B} - \mathbf{A} \quad (1.1.4.2)$$

is the direction vector of AB .

Question 1.1.4 :Find the parametric equation of AB, BC, CA .

Solution: The parametric equation for AB is given by

$$\mathbf{x} = \mathbf{A} + k\mathbf{m} \quad (1.1.4.3)$$

$$\text{where, } \mathbf{m} = \mathbf{B} - \mathbf{A} \quad (1.1.4.4)$$

$$= \begin{pmatrix} 3 \\ -5 \end{pmatrix} - \begin{pmatrix} -3 \\ -5 \end{pmatrix} \quad (1.1.4.5)$$

$$= \begin{pmatrix} 6 \\ 0 \end{pmatrix} \quad (1.1.4.6)$$

Hence we get,

$$AB : \mathbf{x} = \begin{pmatrix} -3 \\ -5 \end{pmatrix} + k \begin{pmatrix} 6 \\ 0 \end{pmatrix} \quad (1.1.4.7)$$

Similarly,

$$BC : \mathbf{x} = \begin{pmatrix} 3 \\ -5 \end{pmatrix} + k \begin{pmatrix} -7 \\ 2 \end{pmatrix} \quad (1.1.4.8)$$

$$CA : \mathbf{x} = \begin{pmatrix} -4 \\ -3 \end{pmatrix} + k \begin{pmatrix} 1 \\ -2 \end{pmatrix} \quad (1.1.4.9)$$

1.1.5. The normal form of the equation of AB is

$$\mathbf{n}^\top (\mathbf{x} - \mathbf{A}) = 0 \quad (1.1.5.1)$$

where

$$\mathbf{n}^\top \mathbf{m} = \mathbf{n}^\top (\mathbf{B} - \mathbf{A}) = 0 \quad (1.1.5.2)$$

$$\text{or, } \mathbf{n} = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix} \mathbf{m} \quad (1.1.5.3)$$

$$\mathbf{n}^\top (\mathbf{x} - \mathbf{A}) = 0 \quad (1.1.5.4)$$

where

$$\mathbf{n}^\top \mathbf{m} = \mathbf{n}^\top (\mathbf{B} - \mathbf{A}) = 0 \quad (1.1.5.5)$$

or,

$$\mathbf{n} = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix} \mathbf{m} \quad (1.1.5.6)$$

Question 1.1.5: Find the normal form of the equations of AB , BC and CA .

Solution: : The normal equation for the side AB is

$$\mathbf{n}^\top (\mathbf{x} - \mathbf{A}) = 0 \quad (1.1.5.7)$$

$$\implies \mathbf{n}^\top \mathbf{x} = \mathbf{n}^\top \mathbf{A} \quad (1.1.5.8)$$

Now our task is to find the \mathbf{n} so that we can find \mathbf{n}^\top . As given in the question

$$\mathbf{n} = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix} \mathbf{m} \quad (1.1.5.9)$$

Here $\mathbf{m} = \mathbf{B} - \mathbf{A}$ for side AB

$$\implies \mathbf{m} = \begin{pmatrix} 3 \\ -5 \end{pmatrix} - \begin{pmatrix} -3 \\ -5 \end{pmatrix} \quad (1.1.5.10)$$

$$= \begin{pmatrix} 6 \\ 0 \end{pmatrix} \quad (1.1.5.11)$$

Now as we have obtained vector \mathbf{m} , we can use this to obtain vector \mathbf{n}

$$\mathbf{n} = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix} \begin{pmatrix} 6 \\ 0 \end{pmatrix} = \begin{pmatrix} 0 \\ -6 \end{pmatrix} \quad (1.1.5.12)$$

The transpose of \mathbf{n} is

$$\mathbf{n}^\top = \begin{pmatrix} 0 & -6 \end{pmatrix} \quad (1.1.5.13)$$

Hence the normal equation of side AB is

$$\begin{pmatrix} 0 & -6 \end{pmatrix} \mathbf{x} = \begin{pmatrix} 0 & -6 \end{pmatrix} \begin{pmatrix} -3 \\ -5 \end{pmatrix} \quad (1.1.5.14)$$

$$\implies \begin{pmatrix} 0 & -6 \end{pmatrix} \mathbf{x} = 30 \quad (1.1.5.15)$$

The normal equation for the side BC is

$$\mathbf{n}^\top (\mathbf{x} - \mathbf{B}) = 0 \quad (1.1.5.16)$$

$$\implies \mathbf{n}^\top \mathbf{x} = \mathbf{n}^\top \mathbf{B} \quad (1.1.5.17)$$

Now our task is to find the \mathbf{n} so that we can find \mathbf{n}^\top . As given in the question

$$\mathbf{n} = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix} \mathbf{m} \quad (1.1.5.18)$$




Figure 1.2: The line **AB** plotted using python

Here $\mathbf{m} = \mathbf{C} - \mathbf{B}$ for side \mathbf{BC}

$$\Rightarrow \mathbf{m} = \begin{pmatrix} -4 \\ -3 \end{pmatrix} - \begin{pmatrix} 3 \\ -5 \end{pmatrix} \quad (1.1.5.19)$$

$$= \begin{pmatrix} -7 \\ 2 \end{pmatrix} \quad (1.1.5.20)$$

Now as we have obtained vector \mathbf{m} . we can use this to obtain vector \mathbf{n}

$$\mathbf{n} = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix} \begin{pmatrix} -7 \\ 2 \end{pmatrix} = \begin{pmatrix} 2 \\ 7 \end{pmatrix} \quad (1.1.5.21)$$

The transpose of \mathbf{n} is

$$\mathbf{n}^T = \begin{pmatrix} 2 & 7 \end{pmatrix} \quad (1.1.5.22)$$

Hence the normal equation of side BC is

$$\begin{pmatrix} 2 & 7 \end{pmatrix} \mathbf{x} = \begin{pmatrix} 2 & 7 \end{pmatrix} \begin{pmatrix} 3 \\ -5 \end{pmatrix} \quad (1.1.5.23)$$

$$\Rightarrow \begin{pmatrix} 2 & 7 \end{pmatrix} \mathbf{x} = -29 \quad (1.1.5.24)$$




Figure 1.3: The line **BC** plotted using python

The normal equation for the side CA is

$$\mathbf{n}^\top (\mathbf{x} - \mathbf{C}) = 0 \quad (1.1.5.25)$$

$$\implies \mathbf{n}^\top \mathbf{x} = \mathbf{n}^\top \mathbf{C} \quad (1.1.5.26)$$

Now our task is to find the \mathbf{n} so that we can find \mathbf{n}^\top . As given in the question

$$\mathbf{n} = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix} \mathbf{m} \quad (1.1.5.27)$$

Here $\mathbf{m} = \mathbf{A} - \mathbf{C}$ for side \mathbf{CA}

$$\implies \mathbf{m} = \begin{pmatrix} -3 \\ -5 \end{pmatrix} - \begin{pmatrix} -4 \\ -3 \end{pmatrix} \quad (1.1.5.28)$$

$$= \begin{pmatrix} 1 \\ -2 \end{pmatrix} \quad (1.1.5.29)$$

Now as we have obtained vector \mathbf{m} , we can use this to obtain vector \mathbf{n}

$$\mathbf{n} = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix} \begin{pmatrix} 1 \\ -2 \end{pmatrix} = \begin{pmatrix} -2 \\ -1 \end{pmatrix} \quad (1.1.5.30)$$

The transpose of \mathbf{n} is

$$\mathbf{n}^\top = \begin{pmatrix} -2 & -1 \end{pmatrix} \quad (1.1.5.31)$$




Figure 1.4: The line **CA** plotted using python

Hence the normal equation of side CA is

$$\begin{pmatrix} -2 & -1 \end{pmatrix} \mathbf{x} = \begin{pmatrix} -2 & -1 \end{pmatrix} \begin{pmatrix} -4 \\ -3 \end{pmatrix} \quad (1.1.5.32)$$

$$\Rightarrow \begin{pmatrix} -2 & -1 \end{pmatrix} \mathbf{x} = 11 \quad (1.1.5.33)$$

1.1.6. The area of $\triangle ABC$ is defined as

$$\frac{1}{2} \|(\mathbf{A} - \mathbf{B}) \times \mathbf{A} - \mathbf{C}\| \quad (1.1.6.1)$$

where

$$\mathbf{A} \times \mathbf{B} \triangleq \begin{vmatrix} 1 & -4 \\ -1 & 6 \end{vmatrix} \quad (1.1.6.2)$$

Question 1.1.6: Find the area of $\triangle ABC$.

Solution: Given,

$$\mathbf{A} = \begin{pmatrix} -3 \\ -5 \end{pmatrix}; \mathbf{B} = \begin{pmatrix} 3 \\ -5 \end{pmatrix}; \mathbf{C} = \begin{pmatrix} -4 \\ -3 \end{pmatrix} \quad (1.1.6.3)$$

$$\mathbf{A} - \mathbf{B} = \begin{pmatrix} -3 \\ -5 \end{pmatrix} - \begin{pmatrix} 3 \\ -5 \end{pmatrix} = \begin{pmatrix} -6 \\ 0 \end{pmatrix} \quad (1.1.6.4)$$

$$\mathbf{A} - \mathbf{C} = \begin{pmatrix} -3 \\ -5 \end{pmatrix} - \begin{pmatrix} -4 \\ -3 \end{pmatrix} = \begin{pmatrix} 1 \\ -2 \end{pmatrix} \quad (1.1.6.5)$$

$$\therefore (\mathbf{A} - \mathbf{B}) \times (\mathbf{A} - \mathbf{C}) = \begin{vmatrix} -6 & 1 \\ 0 & -2 \end{vmatrix} \quad (1.1.6.6)$$

$$= (-6) \times (-2) - 1 \times 0 \quad (1.1.6.7)$$

$$= 12 - 0 \quad (1.1.6.8)$$

$$= 12 \quad (1.1.6.9)$$

$$\Rightarrow \frac{1}{2} \|(\mathbf{A} - \mathbf{B}) \times (\mathbf{A} - \mathbf{C})\| = \frac{12}{2} = 6 \quad (1.1.6.10)$$

1.1.7. Question 1.1.7: Find the angles $\mathbf{A}, \mathbf{B}, \mathbf{C}$, given that

$$\cos A \triangleq \frac{(\mathbf{B} - \mathbf{A})^\top (\mathbf{C} - \mathbf{A})}{\|\mathbf{B} - \mathbf{A}\| \|\mathbf{C} - \mathbf{A}\|} \quad (1.1.7.1)$$

Solution:

From the given values of $\mathbf{A}, \mathbf{B}, \mathbf{C}$,

(a) Finding the value of angle A

$$\mathbf{B} - \mathbf{A} = \begin{pmatrix} 6 \\ 0 \end{pmatrix} \quad (1.1.7.2)$$

and

$$\mathbf{C} - \mathbf{A} = \begin{pmatrix} -1 \\ +2 \end{pmatrix} \quad (1.1.7.3)$$

also calculating the values of norms

$$\|\mathbf{B} - \mathbf{A}\| = \sqrt{36} = 6 \quad (1.1.7.4)$$

$$\|\mathbf{C} - \mathbf{A}\| = \sqrt{5} \quad (1.1.7.5)$$

and by doing matrix multiplication we get,

$$\begin{aligned}(\mathbf{B} - \mathbf{A})^\top (\mathbf{C} - \mathbf{A}) &= \begin{pmatrix} 6 & 0 \end{pmatrix} \begin{pmatrix} -1 \\ +2 \end{pmatrix} \\ &= -6\end{aligned}\tag{1.1.7.6}$$

so

$$\cos A = \frac{-6}{6\sqrt{5}}\tag{1.1.7.7}$$

$$= \frac{-1}{\sqrt{5}}\tag{1.1.7.8}$$

$$\implies A = \cos^{-1} \frac{-1}{\sqrt{5}}\tag{1.1.7.9}$$

(b) Finding the value of angle B

$$\mathbf{C} - \mathbf{B} = \begin{pmatrix} -7 \\ 2 \end{pmatrix}\tag{1.1.7.10}$$

and

$$\mathbf{A} - \mathbf{B} = \begin{pmatrix} -6 \\ 0 \end{pmatrix}\tag{1.1.7.11}$$

also calculating the values of norms

$$\|\mathbf{C} - \mathbf{B}\| = \sqrt{5}\tag{1.1.7.12}$$

$$\|\mathbf{A} - \mathbf{B}\| = \sqrt{53}\tag{1.1.7.13}$$

and by doing matrix multiplication we get,

$$\begin{aligned}(\mathbf{C} - \mathbf{B})^\top(\mathbf{A} - \mathbf{B}) &= \begin{pmatrix} -7 & 2 \end{pmatrix} \begin{pmatrix} -6 \\ 0 \end{pmatrix} \\ &= 42\end{aligned}\tag{1.1.7.14}$$

so

$$\cos B = \frac{42}{\sqrt{36}\sqrt{53}}\tag{1.1.7.15}$$

$$= \frac{7}{\sqrt{53}}\tag{1.1.7.16}$$

$$\implies B = \cos^{-1} \frac{7}{\sqrt{53}}\tag{1.1.7.17}$$

(c) Finding the value of angle C

$$\mathbf{A} - \mathbf{C} = \begin{pmatrix} 1 \\ -2 \end{pmatrix}\tag{1.1.7.18}$$

and

$$\mathbf{B} - \mathbf{C} = \begin{pmatrix} 7 \\ -2 \end{pmatrix}\tag{1.1.7.19}$$

also calculating the values of norms

$$\|\mathbf{A} - \mathbf{C}\| = \sqrt{5}\tag{1.1.7.20}$$

$$\|\mathbf{B} - \mathbf{C}\| = \sqrt{53}\tag{1.1.7.21}$$

and by doing matrix multiplication we get,

$$\begin{aligned}(\mathbf{A} - \mathbf{C})^T(\mathbf{B} - \mathbf{C}) &= \begin{pmatrix} 1 & -2 \end{pmatrix} \begin{pmatrix} 7 \\ -2 \end{pmatrix} \\ &= 11\end{aligned}\tag{1.1.7.22}$$

so

$$\cos C = \frac{11}{\sqrt{5}\sqrt{53}}\tag{1.1.7.23}$$

$$= \frac{11}{\sqrt{265}}\tag{1.1.7.24}$$

$$\implies C = \cos^{-1} \frac{11}{\sqrt{265}}\tag{1.1.7.25}$$

1.2. Median

1.3. Altitude

1.4. Perpendicular Bisector

1.5. Angle Bisector