
GSSSB · STAFF NURSE EXAM

Nutrition MCQ Bank

370 MCQs · GSSSB pattern · Answer key with explanation



10 Sections | 370 MCQs | Direct + Priority + Scenario | Match & Assertion-Reason | Calculations | Explanations

Hinglish explanation + English exam terms



Sister Asha
Aapki Nursing teacher

MCQ Bank – Sections

1. **Food Groups, Energy & Balanced Diet**
   30 MCQs · Nutrients, food groups, kcal values, BMR, SDA, RDA, reference man, My Plate, cooking & anti-nutritional factors
2. **Carbohydrates, Proteins, Fats, Fibre & Water**
   30 MCQs · Sugars, GI, fibre, amino acids, protein quality, fatty acids, EFA, cholesterol, trans fat, water & electrolytes
3. **Vitamins**
   30 MCQs · Fat-soluble A, D, E, K; B-complex & C – sources, functions, deficiencies, prophylaxis
4. **Minerals & Trace Elements**
   30 MCQs · Calcium, iron & anaemia, Anaemia Mukt Bharat, iodine & IDD, zinc, fluoride, selenium, copper
5. **Nutrition Through the Life Span**
   30 MCQs · Pregnancy, lactation, breastfeeding, breast milk, problems, complementary feeding, LBW, children, adolescents, elderly
6. **Malnutrition & Nutritional Assessment**
   30 MCQs · PEM, kwashiorkor vs marasmus, SAM/ MAM, NRC, WHO 10 steps, classifications, obesity & BMI, anthropometry, ABCD
7. **Therapeutic Diets & Diet in Diseases**
   30 MCQs · Hospital diets, diabetes, hypertension, cardiac, renal, liver, GI, fever, burns, gout, PKU, enteral & parenteral nutrition, drug-food interactions
8. **Food Hygiene, Safety, Preservation & Adulteration**
   30 MCQs · Food hygiene, milk pasteurisation, food poisoning, preservation, fortification, adulteration tests, FSSAI, labelling
9. **National Nutrition Programmes**
   30 MCQs · ICDS, POSHAN Abhiyaan, PM POSHAN, micronutrient programmes, NRC, NFSA, global targets, nutrition education
10. **GSSSB Pattern Mock Test – 100 MCQs (Full Nutrition Syllabus)**
    100 MCQs · 100 questions, 100 marks, 0.25 negative marking – complete Nutrition & Dietetics

Level 1 – Direct: facts, values, definitions

Level 2 – Priority: "first / best / immediate" action

Level 3 – Advanced: scenario, NOT/EXCEPT, match, A-R, calculation



MCQ kaise solve karein?

Pehle khud answer socho, phir options padho. Har section ke baad answer key aur explanation hai, galti wale questions ka number apni notebook me likho. Exam me galat answer par **0.25 marks** kat sakte hain, isliye bina soche guess mat karo.

1 Food Groups, Energy & Balanced Diet

? Level 1 – Direct

1. Most essential nutrient –
(a) Water (b) Vitamin (c) Carbohydrate (d) Protein

2. 1 g fat se energy –
(a) 2 kcal (b) 4 kcal (c) 7 kcal (d) 9 kcal

3. 1 g protein se energy –
(a) 4 kcal (b) 6 kcal (c) 7 kcal (d) 9 kcal

4. 1 g alcohol se energy –
(a) 0 kcal (b) 4 kcal (c) 7 kcal (d) 9 kcal

5. 1 kcal =
(a) 0.24 kJ (b) 1 kJ (c) 4.184 kJ (d) 10 kJ

6. Total energy expenditure ka sabse bada hissa –
(a) Growth (b) SDA (c) Physical activity (d) BMR

7. BMR per kg sabse zyada –
(a) Elderly (b) Adults (c) Infants (d) Adolescents

8. Highest SDA –
(a) Fibre (b) Carbohydrate (c) Fat (d) Protein

9. RDA meets needs of –
(a) 50 % (b) 75 % (c) 97.5 % population (d) 100 %

10. ICMR 2020 reference man weight –
(a) 55 kg (b) 60 kg (c) 65 kg (d) 70 kg

11. ICMR 2020 reference woman weight –
(a) 45 kg (b) 50 kg (c) 55 kg (d) 60 kg

12. Pregnancy extra energy (2nd–3rd trimester) –
(a) +150 kcal (b) +350 kcal (c) +600 kcal (d) +1000 kcal

13. Lactation extra energy (0–6 months) –
(a) +200 kcal (b) +350 kcal (c) +600 kcal (d) +900 kcal

14. Salt intake limit (WHO) –
(a) < 2 g/day (b) < 5 g/day (c) < 10 g/day (d) < 15 g/day

15. Cereals me limiting amino acid –
(a) Lysine (b) Tryptophan (c) Leucine (d) Methionine

16. Pulses me limiting amino acid –
(a) Methionine (b) Histidine (c) Valine (d) Lysine

17. Richest plant protein –
(a) Wheat (b) Rice (c) Groundnut (d) Soybean

18. Richest calcium cereal –
(a) Wheat (b) Ragi (c) Maize (d) Rice

19. Lathyrism –
(a) Khesari dal (b) Argemone oil (c) Groundnut (d) Maize

20. International Year of Millets –
(a) 2018 (b) 2020 (c) 2023 (d) 2025

? Level 2 – Priority & Application

21. Vitamin bachane ke liye best cooking –
(a) Adding soda (b) Deep frying (c) Boiling & discarding water (d) Pressure cooking/ steaming

22. Dal me soda (sodium bicarbonate) daalne se nash –
(a) Vitamin A (b) Thiamine & vitamin C (c) Calcium (d) Iron

23. Cereal-pulse combination (khichdi) ka fayda –
(a) Mutual supplementation of amino acids (b) More sugar (c) Less fibre (d) More fat

24. Raw egg white kaunse vitamin ko bind karta hai –
(a) Folate (b) Biotin (c) Vitamin A (d) B12

25. Germination/ sprouting se ↑ –
(a) Fat (b) Phytates (c) Vitamin C & B (d) Cholesterol

26. 38 °C → 39 °C fever me BMR ↑ –
(a) ~13 % (b) ~50 % (c) ~2 % (d) 0 %

27. ICMR My Plate – half plate –
(a) Vegetables + fruits (b) Cereals (c) Pulses (d) Fats

? Level 3 – Advanced (NOT / Match / Assertion-Reason / Calculation)

28. Protective food NAHI –
(a) Milk (b) Sugar (c) Fruits (d) GLV

29. Assertion (A): Vegetables ko kaatne se pehle dhona chahiye. Reason (R): Kaatne ke baad dhone se water-soluble vitamins nikal jaate hain.
(a) Both A and R true, R explains A (b) Both A and R true, R does not explain A (c) A true, R false (d) A false, R true

30. Diet: CHO 300 g, protein 60 g, fat 50 g – total energy –
(a) 1690 kcal (b) 1810 kcal (c) 1890 kcal (d) 2000 kcal

✓ Answer Key & Explanation

1-a Water.
2-d 9.
3-a 4.
4-c 7.
5-c 4.184.
6-d BMR.
7-c Infants.
8-d ~30 %.
9-c EAR + 2 SD.
10-c 65.
11-c 55.
12-b +350.
13-c +600.
14-b 5 g.
15-a Lysine.
16-a Methionine.
17-d ~40 %.
18-b Ragi.
19-a BOAA.
20-c 2023.
21-d Pressure.
22-b Alkali.
23-a Lysine + methionine.
24-b Avidin.

25-c Sprouts.
26-a 13 %/°C.
27-a Half.
28-b Sugar – empty calories.
29-a Dono sahi, R reason hai.
30-c 1200 + 240 + 450 = 1890.

2 Carbohydrates, Proteins, Fats, Fibre & Water

Level 1 – Direct

1. Sweetest sugar –
   (a) Lactose (b) Fructose (c) Sucrose (d) Glucose
2. Milk sugar –
   (a) Lactose (b) Galactose only (c) Maltose (d) Sucrose
3. Animal starch –
   (a) Glycogen (b) Pectin (c) Cellulose (d) Amylose
4. Brain ka main fuel –
   (a) Ketones always (b) Glucose (c) Fatty acids (d) Protein
5. Glucose ka GI –
   (a) 0 (b) 50 (c) 70 (d) 100
6. Soluble fibre ka effect –
   (a) ↑ Cholesterol (b) Constipation (c) ↓ Cholesterol & blood sugar (d) ↑ Fat absorption
7. Essential amino acids –
   (a) 5 (b) 9 (c) 12 (d) 20
8. Reference protein –
   (a) Egg (b) Milk (c) Meat (d) Soybean
9. Protein me nitrogen –
   (a) 6.25 % (b) 10 % (c) 16 % (d) 25 %
10. Best protein quality index –
    (a) PER only (b) NPU (c) BMI (d) GI
11. Most concentrated energy –
    (a) Carbohydrate (b) Fat (c) Fibre (d) Protein
12. Essential fatty acids –
    (a) Linoleic & α-linolenic acid (b) Palmitic & stearic (c) Oleic only (d) Lauric
13. EFA deficiency –
    (a) Phrynoderma (b) Rickets (c) Pellagra (d) Scurvy
14. Highest saturated fat oil –
    (a) Sunflower (b) Mustard (c) Coconut oil (d) Olive
15. ω-3 rich –
    (a) Coconut oil (b) Vanaspati (c) Ghee (d) Fish oil & flaxseed
16. Most harmful fat –
    (a) ω-3 (b) MUFA (c) PUFA (d) Trans fat
17. Cholesterol present only in –
    (a) Plant foods (b) Cereals (c) Animal foods (d) Fruits
18. Good cholesterol –
    (a) VLDL (b) HDL (c) LDL (d) Chylomicron

19. Body water (adult) –
(a) 25 % (b) 40 % (c) 60 % (d) 80 %

20. WHO ORS osmolarity –
(a) 150 mOsm/L (b) 245 mOsm/L (c) 311 mOsm/L (d) 500 mOsm/L

? Level 2 – Priority & Application

21. Diabetic patient ke liye better –
(a) High GI foods (b) Low GI foods (c) White bread (d) Sugar drinks

22. Lactose intolerant patient – better tolerated –
(a) Milk (b) Condensed milk (c) Ice cream (d) Curd/ buttermilk

23. Burns/ fever – nitrogen balance –
(a) Negative (b) Equilibrium (c) Positive (d) Zero

24. Fat malabsorption me oil –
(a) Coconut cream (b) Ghee (c) Vanaspati (d) MCT oil

25. CKD patient me avoid –
(a) Apple (b) Banana & coconut water (K^+) (c) Rice (d) Sugar

26. Pancreatitis/ gall bladder disease – diet –
(a) High fibre only (b) High sodium (c) Low fat (d) High fat

27. Constipation – advice –
(a) High fibre + fluids (b) Low fibre (c) Only milk (d) Fluid restriction

? Level 3 – Advanced (NOT / Match / Assertion-Reason / Calculation)

28. Monosaccharide NAHI –
(a) Sucrose (b) Fructose (c) Glucose (d) Galactose

29. Assertion (A): Carbohydrate ko protein-sparing kaha jata hai. Reason (R): Adequate carbohydrate hone par protein energy ke liye use nahi hota.
(a) Both A and R true, R explains A (b) Both A and R true, R does not explain A (c) A true, R false (d) A false, R true

30. Match: P. Sucrose Q. Lactose R. Maltose S. Glycogen – 1. Glucose + fructose 2. Glucose + galactose 3. Glucose + glucose 4. Animal starch
(a) P-2, Q-1, R-3, S-4 (b) P-4, Q-2, R-3, S-1 (c) P-1, Q-3, R-2, S-4 (d) P-1, Q-2, R-3, S-4

✓ Answer Key & Explanation

1-b Fructose.
2-a Lactose.
3-a Glycogen.
4-b Glucose.
5-d 100.
6-c Soluble.
7-b 9.
8-a Egg.
9-c 16.
10-b NPU.
11-b 9 kcal.
12-a EFA.
13-a Toad skin.
14-c Coconut.
15-d ω-3.
16-d Trans.
17-c Animal.
18-b HDL.
19-c 60 %.
20-b 245.
21-b Low GI.
22-d Fermented.

23-a Catabolism.
24-d MCT.
25-b Potassium.
26-c Low fat.
27-a Fibre.
28-a Disaccharide.
29-a Dono sahi, R reason hai.
30-d Sahi match.

3 Vitamins

Level 1 – Direct

1. 'Vitamine' term –
   (a) McCollum (b) Eijkman (c) Lind (d) Casimir Funk
2. Fat-soluble vitamins –
   (a) D, C (b) B, C (c) A, D, E, K (d) A, B, C
3. Earliest symptom of vitamin A deficiency –
   (a) Night blindness (b) Bitot's spots (c) Corneal ulcer (d) Keratomalacia
4. Bitot's spots –
   (a) Vitamin B2 (b) Vitamin C (c) Vitamin A (d) Vitamin D
5. Richest source of vitamin A –
   (a) Rice (b) Milk (c) Amla (d) Fish liver oil
6. Sunshine vitamin –
   (a) Vitamin K (b) Vitamin C (c) Vitamin A (d) Vitamin D
7. Active vitamin D formed in –
   (a) Skin (b) Liver (c) Kidney (d) Intestine
8. Rickets –
   (a) Vitamin K (b) Vitamin B1 (c) Vitamin D (d) Vitamin C
9. Antioxidant fat-soluble vitamin –
   (a) Vitamin E (b) Vitamin K (c) Vitamin A only (d) Vitamin D
10. Haemorrhagic disease of newborn –
    (a) Iron (b) Vitamin K deficiency (c) Vitamin C (d) Folic acid
11. Beriberi –
    (a) Thiamine (b) Riboflavin (c) Pyridoxine (d) Niacin
12. Angular stomatitis –
    (a) Vitamin D (b) Vitamin K (c) Riboflavin (d) Thiamine
13. Pellagra –
    (a) Thiamine (b) B12 (c) Vitamin C (d) Niacin
14. INH neuropathy prevention –
    (a) Vitamin C (b) Folate (c) Pyridoxine (d) Thiamine
15. Neural tube defects –
    (a) Vitamin K (b) Iron (c) Folic acid deficiency (d) Vitamin C
16. B12 present only in –
    (a) Animal foods (b) Vegetables (c) Cereals (d) Fruits
17. Scurvy –
    (a) Vitamin D (b) Vitamin E (c) Vitamin C (d) Vitamin A
18. Richest vitamin C –
    (a) Milk (b) Amla (c) Egg (d) Rice

19. Most easily destroyed vitamin –
(a) Vitamin A (b) Vitamin C (c) Vitamin D (d) Vitamin K

20. Vitamin K to newborn –
(a) 0.1 mg IV (b) 1 mg oral daily (c) 10 mg IM (d) 1 mg IM

? Level 2 – Priority & Application

21. Vitamin A prophylaxis 1st dose –
(a) 1 lakh IU at 9 months (b) 50,000 IU at 6 weeks (c) 2 lakh IU at birth (d) 2 lakh IU at 5 yrs

22. Infant me bulging fontanelle + vomiting after high-dose drops –
(a) Pellagra (b) Hypervitaminosis A (c) Scurvy (d) Rickets

23. Alcoholic – confusion, ataxia, ophthalmoplegia – pehle do –
(a) Glucose only (b) Thiamine before glucose (c) Vitamin C (d) Folate

24. Maize-eating population me deficiency –
(a) Scurvy (b) Rickets (c) Beriberi (d) Pellagra

25. Strict vegetarian (vegan) me risk –
(a) Vitamin C (b) Vitamin K (c) Folate (d) B12 deficiency

26. Earliest sign of rickets –
(a) Rickety rosary (b) Pigeon chest (c) Craniotabes (d) Bow legs

27. Breastfed infant supplement –
(a) Vitamin A 2 lakh IU daily (b) Vitamin C 500 mg (c) Vitamin D 400 IU (d) Iron 100 mg

? Level 3 – Advanced (NOT / Match / Assertion-Reason / Calculation)

28. Water-soluble vitamin jo liver me store hota hai –
(a) B2 (b) B1 (c) Vitamin C (d) B12

29. Assertion (A): B12 deficiency me sirf folic acid dena harmful hai. Reason (R): Folic acid anaemia theek kar deta hai par neurological damage badhta rehta hai.
(a) Both A and R true, R explains A (b) Both A and R true, R does not explain A (c) A true, R false (d) A false, R true

30. Match: P. Vitamin A Q. Vitamin D R. Vitamin C S. Niacin – 1. Xerophthalmia 2. Rickets 3. Scurvy 4. Pellagra
(a) P-1, Q-3, R-2, S-4 (b) P-4, Q-2, R-3, S-1 (c) P-1, Q-2, R-3, S-4 (d) P-2, Q-1, R-3, S-4

✓ Answer Key & Explanation

1-d Funk.
2-c ADEK.
3-a XN.
4-c A.
5-d Fish liver oil.
6-d D.
7-c Calcitriol.
8-c D.
9-a E.
10-b K.
11-a B1.
12-c B2.
13-d B3.
14-c B6.
15-c Folate.
16-a Animal.
17-c C.
18-b Amla.
19-b C.
20-d 1 mg.
21-a 9 months.
22-b Raised ICP.

23-b Wernicke.
24-d Niacin.
25-d B12.
26-c Craniotabes.
27-c D.
28-d 3–5 yrs.
29-a Dono sahi, R reason hai.
30-c Sahi match.

4 Minerals & Trace Elements

Level 1 – Direct

1. Most abundant mineral in body –
   (a) Iron (b) Calcium (c) Sodium (d) Phosphorus
2. Calcium RDA adult (ICMR 2020) –
   (a) 200 mg (b) 400 mg (c) 1000 mg (d) 2000 mg
3. Best source of calcium –
   (a) Milk (b) Rice (c) Potato (d) Apple
4. Iron storage form –
   (a) Transferrin (b) Myoglobin (c) Ferritin (d) Haemoglobin
5. Iron transport protein –
   (a) Ferritin (b) Transferrin (c) Ceruloplasmin (d) Albumin
6. Iron absorption site –
   (a) Colon (b) Ileum (c) Duodenum (d) Stomach
7. Iron absorption enhancer –
   (a) Tea (b) Phytates (c) Vitamin C (d) Calcium
8. Iron RDA adult woman (ICMR 2020) –
   (a) 10 mg (b) 19 mg (c) 29 mg (d) 50 mg
9. Earliest marker of iron depletion –
   (a) Hb (b) Serum iron (c) Serum ferritin (d) MCV
10. Spoon-shaped nails –
    (a) Iron deficiency (b) Zinc (c) Calcium (d) Iodine
11. Anaemia in pregnancy (WHO) –
    (a) Hb < 7 (b) Hb < 11 g/dl (c) Hb < 9 (d) Hb < 13
12. IFA tablet in pregnancy –
    (a) 20 mg + 100 µg (b) 45 mg + 400 µg (c) 60 mg iron + 500 µg folic acid (d) 100 mg + 5 mg
13. Iodine RDA adult –
    (a) 50 µg (b) 150 µg (c) 250 µg (d) 500 µg
14. Iodised salt at consumer level –
    (a) ≥ 5 ppm (b) ≥ 15 ppm (c) ≥ 30 ppm (d) ≥ 100 ppm
15. Goitrogen –
    (a) Cabbage (b) Apple (c) Milk (d) Rice
16. Optimum fluoride in water –
    (a) 0.1 ppm (b) 0.5–0.8 ppm (c) 2–3 ppm (d) 5 ppm
17. Defluoridation technique –
    (a) Chlorination (b) Baudouin (c) Lactometer (d) Nalgonda
18. Keshan disease –
    (a) Iodine (b) Copper (c) Zinc (d) Selenium deficiency

19. Wilson's disease –
(a) Copper excess (b) Iron excess (c) Calcium (d) Zinc deficiency

20. Cobalt part of –
(a) Haemoglobin (b) Vitamin B12 (c) Thyroxine (d) Vitamin C

? Level 2 – Priority & Application

21. IFA ke saath avoid –
(a) Amla (b) Tea/ coffee (c) Lemon juice (d) Water

22. Iron syrup child – teeth staining se bachne ke liye –
(a) Milk ke saath (b) Chewable (c) Straw/ dropper back of tongue (d) Brush na kare

23. Child with diarrhoea – zinc dose (> 6 months) –
(a) 5 mg × 30 days (b) 10 mg × 3 days (c) 20 mg × 14 days (d) 50 mg once

24. Adolescent WIFS tablet –
(a) Syrup biweekly (b) Blue, weekly (c) Pink, daily (d) Red, daily

25. Children 6–59 months AMB –
(a) IFA syrup 1 ml biweekly (b) Red tablet daily (c) Nothing (d) Blue weekly

26. Thyroidectomy ke baad tingling, carpopedal spasm –
(a) Hyponatraemia (b) Hypocalcaemia (c) Iron deficiency (d) Hyperkalaemia

27. Child with mottled teeth – region with high fluoride –
(a) Caries (b) Dental fluorosis (c) Scurvy (d) Rickets

? Level 3 – Advanced (NOT / Match / Assertion-Reason / Calculation)

28. Iron absorption ko ↓ NAHI karta –
(a) Calcium (b) Tea (c) Phytates (d) Vitamin C

29. Assertion (A): Iodine deficiency sabse common preventable cause of mental retardation hai.
Reason (R): Fetal brain development ke liye thyroid hormone zaroori hai.
(a) Both A and R true, R explains A (b) Both A and R true, R does not explain A (c) A true, R false (d) A false, R true

30. Match: P. Iron Q. Iodine R. Zinc S. Fluoride – 1. Anaemia 2. Goitre 3. Hypogeusia 4. Dental caries (deficiency)
(a) P-1, Q-3, R-2, S-4 (b) P-1, Q-2, R-3, S-4 (c) P-4, Q-2, R-3, S-1 (d) P-2, Q-1, R-3, S-4

✓ Answer Key & Explanation

1-b Calcium.
2-c 1000.
3-a Milk.
4-c Ferritin.
5-b Transferrin.
6-c Duodenum.
7-c Vitamin C.
8-c 29.
9-c Ferritin.
10-a Koilonychia.
11-b 11.
12-c Red tablet.
13-b 150.
14-b 15 ppm.
15-a Cabbage.
16-b 0.5–0.8.
17-d Nalgonda.
18-d Selenium.
19-a Copper.
20-b B12.
21-b Tannins.
22-c Staining.

23-c Zinc.
24-b WIFS.
25-a Syrup.
26-b Tetany.
27-b Fluorosis.
28-d Enhancer.
29-a Dono sahi, R reason hai.
30-b Sahi match.

5 Nutrition Through the Life Span

Level 1 – Direct

1. Breastfeeding initiation –
   (a) Within 1 hr of birth (b) After 6 hrs (c) After colostrum discard (d) After 24 hrs
2. Exclusive breastfeeding –
   (a) 1 year (b) 3 months (c) 4 months (d) 6 months
3. Breastfeeding continue till –
   (a) 1 year (b) 2 years or beyond (c) 5 years (d) 6 months
4. Colostrum rich in –
   (a) IgA & protein (b) Fat (c) Lactose (d) Iron
5. Highest lactose –
   (a) Human milk (b) Cow milk (c) Buffalo milk (d) Goat milk
6. Highest fat –
   (a) Goat milk (b) Cow milk (c) Buffalo milk (d) Human milk
7. Milk production hormone –
   (a) Prolactin (b) Oxytocin (c) FSH (d) Oestrogen
8. Milk ejection hormone –
   (a) GH (b) Oxytocin (c) Progesterone (d) Prolactin
9. Fat-rich milk at end of feed –
   (a) Transitional (b) Hindmilk (c) Foremilk (d) Colostrum
10. Breast milk is low in –
    (a) Lactose (b) Vitamin D & K (c) IgA (d) Protein (relatively)
11. Complementary feeding starts –
    (a) 1 year (b) 4 months (c) 6 months (d) 9 months
12. Honey avoided in < 1 yr –
    (a) Botulism risk (b) Allergy (c) Diabetes (d) Obesity
13. EBM in refrigerator –
    (a) 1 hr (b) 24 hrs (c) 7 days (d) 1 month
14. Bottle feeding major hazard –
    (a) Rickets (b) Obesity (c) Diarrhoea (d) Anaemia
15. LBW –
    (a) < 1500 g (b) < 2000 g (c) < 2500 g (d) < 3000 g
16. Normal weight gain pregnancy (normal BMI) –
    (a) 2–3 kg (b) 5–7 kg (c) 11.5–16 kg (d) 20–25 kg
17. Folic acid for NTD prevention –
    (a) 1 g (b) 5 mg always (c) 50 µg (d) 400 µg preconception
18. Calcium supplement pregnancy –
    (a) 2 g daily from conception (b) 500 mg BD from 14 weeks (c) 100 mg once (d) None

19. World Breastfeeding Week –
(a) 1–7 August (b) 14 November (c) 5 May (d) 1–7 September

20. IMS Act –
(a) 1975 (b) 1992 (c) 2006 (d) 2013

? Level 2 – Priority & Application

21. Sign of good attachment –
(a) Clicking sound (b) Nipple only in mouth (c) Lower lip everted, chin touching breast (d) Cheeks drawn in

22. Cracked nipples – main cause –
(a) Too frequent feeds (b) Infection only (c) Poor attachment (d) Small breast

23. Engorgement – management –
(a) Fluid restriction (b) Bind breasts (c) Stop feeding (d) Frequent feeding & expression

24. Mastitis – mother –
(a) Stop breastfeeding (b) Continue breastfeeding (c) Only formula (d) Only other breast forever

25. Adequate milk sign –
(a) No weight gain (b) Baby cries after feed (c) Urine ≥ 6 times/day (d) Hard stools

26. 32–34 weeks preterm – feeding –
(a) IV fluids only (b) Direct only (c) Bottle (d) EBM by paladai/ spoon

27. Complementary food energy density badhane ke liye –
(a) 1 tsp oil/ ghee per meal (b) Dal water only (c) Add water (d) Rice water

? Level 3 – Advanced (NOT / Match / Assertion-Reason / Calculation)

28. Breastfeeding contraindicated –
(a) Mastitis (b) Maternal hepatitis B (c) Maternal common cold (d) Galactosaemia in baby

29. Assertion (A): Mother anxious ho to milk flow kam lagta hai. Reason (R): Stress oxytocin reflex ko inhibit karta hai.
(a) Both A and R true, R explains A (b) Both A and R true, R does not explain A (c) A true, R false (d) A false, R true

30. Match: P. Foremilk Q. Hindmilk R. Colostrum S. Transitional – 1. Thirst 2. Satiety/ fat 3. First immunisation 4. Day 5–14
(a) P-1, Q-3, R-2, S-4 (b) P-2, Q-1, R-3, S-4 (c) P-1, Q-2, R-3, S-4 (d) P-4, Q-2, R-3, S-1

✓ Answer Key & Explanation

1-a 1 hr.
2-d 6 months.
3-b 2 yrs.
4-a IgA.
5-a Human.
6-c Buffalo.
7-a Prolactin.
8-b Oxytocin.
9-b Hindmilk.
10-b D & K.
11-c 6 months.
12-a Botulism.
13-b 24 hrs.
14-c Infection.
15-c 2500.
16-c IOM.
17-d 0.4 mg.
18-b MoHFW.
19-a August.
20-b IMS.
21-c Attachment.
22-c Attachment.

23-d Empty.
24-b Continue.
25-c Wet nappies.
26-d Paladai.
27-a Oil.
28-d Galactosaemia.
29-a Dono sahi, R reason hai.
30-c Sahi match.

6 Malnutrition & Nutritional Assessment

Level 1 – Direct

1. Kwashiorkor term –
   (a) Waterlow (b) Gomez (c) Cicely Williams (d) Jelliffe
2. Hallmark of kwashiorkor –
   (a) Hunger (b) Oedema (c) Severe wasting (d) Old-man face
3. Flag sign –
   (a) Scurvy (b) Kwashiorkor (c) Rickets (d) Marasmus
4. Marasmus age –
   (a) Adolescents (b) 1–4 yrs (c) 5–10 yrs (d) < 1 year
5. SAM by MUAC –
   (a) < 10 cm (b) < 11.5 cm (c) < 12.5 cm (d) < 13.5 cm
6. SAM by WFH –
   (a) < −1 SD (b) > +2 SD (c) < −3 SD (d) < −2 SD
7. MAM by MUAC –
   (a) < 11.5 cm (b) > 13.5 cm (c) 10–11 cm (d) 11.5–12.5 cm
8. Complicated SAM managed at –
   (a) AWC (b) Home only (c) NRC (d) School
9. ORS for SAM –
   (a) Plain water (b) Standard ORS (c) NS IV (d) ReSoMal
10. Initial feed in SAM –
    (a) F-100 (b) F-75 (c) Cow's milk (d) RUTF only
11. Refeeding syndrome hallmark –
    (a) Hypernatraemia (b) Hypercalcaemia (c) Hyperkalaemia (d) Hypophosphataemia
12. Stunting –
    (a) Low weight-for-height (b) Low weight-for-age (c) Low height-for-age (d) High BMI
13. Wasting –
    (a) High weight (b) Low head circumference (c) Low height-for-age (d) Low weight-for-height
14. Gomez classification based on –
    (a) BMI (b) MUAC (c) Weight-for-age (d) Height-for-age
15. BMI formula –
    (a) Weight/ age (b) Weight (kg) / height $(m)^2$ (c) Height/ weight (d) Weight × height
16. Indian obesity BMI –
    (a) ≥ 23 (b) ≥ 25 (c) ≥ 30 (d) ≥ 35
17. Indian women central obesity waist –
    (a) ≥ 80 cm (b) ≥ 88 cm (c) ≥ 90 cm (d) ≥ 102 cm
18. Skinfold thickness –
    (a) Shakir tape (b) Infantometer (c) Harpenden calipers (d) Lactometer

19. Most sensitive indicator of current nutrition –
(a) Head circumference (b) Chest (c) Height (d) Weight

20. Head = chest circumference at –
(a) 1 year (b) 6 months (c) 3 years (d) Birth

Level 2 – Priority & Application

21. SAM child – first 24 hrs main killers –
(a) Hypertension (b) Fever only (c) Hypoglycaemia & hypothermia (d) Obesity

22. Iron in SAM –
(a) After stabilisation (~2 weeks) (b) Only IV (c) Day 1 (d) Never

23. SAM child with dehydration – IV fluids only if –
(a) Mild dehydration (b) Never (c) Always (d) Shock

24. Good weight gain in rehabilitation –
(a) 0 (b) 1 g/kg/day (c) > 10 g/kg/day (d) 50 g/kg/day

25. MUAC age-independent between –
(a) Newborn (b) 10–15 yrs (c) Adults (d) 1–5 yrs

26. Commonest dietary survey method –
(a) Inventory (b) 24-hr recall (c) Diet history only (d) Weighment

27. Most accurate dietary survey –
(a) Questionnaire (b) FFQ (c) Weighment (d) 24-hr recall

Level 3 – Advanced (NOT / Match / Assertion-Reason / Calculation)

28. Kwashiorkor feature NAHI –
(a) Flaky-paint skin (b) Oedema (c) Fatty liver (d) Alert & hungry

29. Assertion (A): Kwashiorkor me weight loss kam dikhta hai. Reason (R): Oedema weight loss ko mask kar deta hai.
(a) Both A and R true, R explains A (b) Both A and R true, R does not explain A (c) A true, R false (d) A false, R true

30. Weight 70 kg, height 1.75 m – BMI –
(a) 20.0 (b) 22.9 (c) 25.0 (d) 40.0

Answer Key & Explanation

1-c Williams.
2-b Oedema.
3-b Hair.
4-d < 1.
5-b 11.5.
6-c −3 SD.
7-d MAM.
8-c NRC.
9-d ReSoMal.
10-b F-75.
11-d ↓ P.
12-c Chronic.
13-d Acute.
14-c Gomez.
15-b BMI.
16-b Asian.
17-a 80.
18-c Calipers.
19-d Weight.
20-a 1 yr.
21-c Steps 1-2.
22-a Not initially.
23-d Heart failure risk.
24-c Good.

25-d MUAC.
26-b Recall.
27-c Weighment.
28-d Marasmus feature.
29-a Dono sahi, R reason hai.
30-b 70/3.0625.

7 Therapeutic Diets & Diet in Diseases

Level 1 – Direct

1. Post-op first feed –
   (a) Soft diet (b) Full liquid with milk (c) Regular diet (d) Clear liquids
2. Clear liquid me NAHI –
   (a) Milk (b) Coconut water (c) Strained juice (d) Clear soup
3. DASH diet –
   (a) CKD (b) Gout (c) Diabetes (d) Hypertension
4. Diabetic diet –
   (a) Complex CHO, high fibre, low GI (b) Low protein only (c) High sugar (d) High fat
5. Hypertension me avoid –
   (a) Vegetables (b) Pickles & papad (c) Low-fat milk (d) Fruits
6. Nephrotic syndrome protein –
   (a) Nil (b) 0.3 g/kg (c) Very high (3 g/kg) (d) Normal (~1 g/kg)
7. Haemodialysis protein –
   (a) Nil (b) 0.4 g/kg (c) 3 g/kg (d) 1.2 g/kg
8. Renal stones – sabse important –
   (a) High salt (b) Fluid restriction (c) ↑ Fluids ≥ 2.5–3 L (d) High oxalate
9. Hepatitis diet –
   (a) No protein (b) High calorie, high CHO (c) High salt (d) Fat-free, low calorie
10. Cirrhosis with ascites –
    (a) Fluid push (b) Sodium restriction (c) Protein nil (d) High salt
11. Gall bladder disease –
    (a) Low fat diet (b) High protein (c) High fat (d) Low fibre
12. Coeliac disease –
    (a) Gluten-free diet (b) Lactose-free (c) Low purine (d) Low protein
13. Gluten present in –
    (a) Rice (b) Maize (c) Wheat (d) Ragi
14. Gout –
    (a) High protein (b) Organ meats (c) Low purine diet (d) High purine
15. PKU –
    (a) Low phenylalanine (b) Low lysine (c) Low sodium only (d) Low glucose
16. Galactosaemia –
    (a) Low protein (b) Gluten-free (c) Stop milk/ breastfeeding (d) High lactose
17. Refractory epilepsy diet –
    (a) Low fat (b) High CHO (c) Ketogenic (d) DASH
18. Preferred nutrition route –
    (a) Enteral (b) IV dextrose (c) Parenteral (d) None

19. Typhoid diet –
(a) Low calorie (b) High fibre (c) High calorie, high protein, low residue (d) Fat-rich

20. Burns diet –
(a) Low protein (b) Fluid restriction (c) Low calorie (d) High calorie & high protein

? Level 2 – Priority & Application

21. Tube feeding position –
(a) Flat supine (b) Prone (c) Head up 30–45° (d) Trendelenburg

22. NG tube placement confirm best –
(a) Whoosh test only (b) X-ray / pH ≤ 5.5 (c) Patient says OK (d) Smell

23. Commonest complication of tube feeding –
(a) Pneumothorax (b) Hyperkalaemia (c) Air embolism (d) Diarrhoea

24. TPN commonest serious complication –
(a) Catheter-related infection (b) Constipation (c) Hypertension (d) Rash

25. TPN bag delayed – nurse –
(a) Double the rate (b) Give bolus (c) Don't speed up to catch up (d) Stop & discard

26. Warfarin patient – advice –
(a) High vitamin K suddenly (b) Consistent vitamin K intake (c) No restriction (d) Avoid all vegetables

27. MAOI patient – avoid –
(a) Aged cheese/ tyramine (b) Milk (c) Banana (d) Rice

? Level 3 – Advanced (NOT / Match / Assertion-Reason / Calculation)

28. CKD diet me restrict NAHI –
(a) Sodium (b) Phosphorus (c) Energy (calories) (d) Potassium

29. Assertion (A): Dumping syndrome me fluids meals ke saath nahi dene chahiye. Reason (R): Fluids gastric emptying ko tez karte hain.
(a) Both A and R true, R explains A (b) Both A and R true, R does not explain A (c) A true, R false (d) A false, R true

30. Match: P. DASH Q. Gluten-free R. Low purine S. Low phenylalanine – 1. Hypertension 2. Coeliac 3. Gout 4. PKU
(a) P-2, Q-1, R-3, S-4 (b) P-1, Q-3, R-2, S-4 (c) P-1, Q-2, R-3, S-4 (d) P-4, Q-2, R-3, S-1

✓ Answer Key & Explanation

1-d Clear.
2-a Milk – full liquid.
3-d HTN.
4-a Diabetes.
5-b Sodium.
6-d Normal.
7-d ↑.
8-c Fluids.
9-b Hepatitis.
10-b Na.
11-a Low fat.
12-a Gluten.
13-c Wheat.
14-c Purine.
15-a PKU.
16-c Galactose.
17-c Keto.
18-a Gut.
19-c Typhoid.
20-d Hypermetabolism.
21-c Aspiration.
22-b Confirm.

23-d Diarrhoea.
24-a Infection.
25-c Constant.
26-b Vitamin K.
27-a Tyramine.
28-c Adequate energy.
29-a Dono sahi, R reason hai.
30-c Sahi match.

8 Food Hygiene, Safety, Preservation & Adulteration

? Level 1 – Direct

1. Temperature danger zone –
   (a) −18 °C (b) 0–4 °C (c) 70–100 °C (d) 5–60 °C
2. Holder pasteurisation –
   (a) 63 °C × 30 min (b) 72 °C × 30 min (c) 100 °C × 10 min (d) 121 °C × 15 min
3. HTST pasteurisation –
   (a) 63 °C × 30 min (b) 72 °C × 15–20 s (c) 80 °C × 1 min (d) 135 °C × 2 s
4. Pasteurisation adequacy test –
   (a) Baudouin (b) Phosphatase test (c) Lactometer (d) Methylene blue test
5. Shortest incubation food poisoning –
   (a) Salmonella (b) Staphylococcal (c) Cholera (d) Botulism
6. Food poisoning with paralysis –
   (a) Salmonella (b) Staph (c) B. cereus (d) Botulism
7. Fried/ reheated rice –
   (a) Shigella (b) Vibrio (c) Clostridium tetani (d) Bacillus cereus
8. Epidemic dropsy –
   (a) Ergot (b) Aflatoxin (c) Khesari dal (d) Argemone oil
9. Argemone test –
   (a) Iodine (b) Nitric acid (c) Baudouin (d) Lactometer
10. Fruit juice preservative –
    (a) Calcium carbide (b) Sodium benzoate (c) Salt (d) Sugar only
11. Jam preserved by –
    (a) Salt (b) Vinegar (c) High sugar (d) Freezing
12. Irradiated food symbol –
    (a) +F (b) ISI (c) Radura (d) Agmark
13. FSS Act –
    (a) 1954 (b) 1992 (c) 2006 (d) 2013
14. World Food Safety Day –
    (a) 7 April (b) 1 June (c) 16 October (d) 7 June
15. Milk water adulteration test –
    (a) Nitric acid (b) Iodine (c) Lactometer (d) HCl
16. Starch in milk –
    (a) Lactometer (b) Burning (c) HCl pink (d) Iodine turns blue
17. Vanaspati in ghee –
    (a) Iodine (b) Nitric acid (c) Magnet (d) Baudouin test
18. Metanil yellow in turmeric –
    (a) Magnet (b) Iodine blue (c) HCl turns pink (d) Water floats

19. Papaya seeds adulterate –
(a) Coffee (b) Chilli (c) Tea (d) Black pepper

20. Fortified rice nutrients –
(a) Iodine (b) Calcium (c) Vitamin A & D (d) Iron, folic acid, B12

? Level 2 – Priority & Application

21. Food handler with boil on hand –
(a) Exclude from food handling (b) Allow with gloves (c) Only cook (d) No issue

22. Cooked food room temp par max –
(a) 24 hrs (b) 2 hrs (c) 2 days (d) 8 hrs

23. +F oil & milk fortified with –
(a) Vitamins A & D (b) B12 (c) Iron (d) Iodine

24. Brick powder in chilli – test –
(a) Iodine (b) Water – brick settles (c) Lactometer (d) Nitric acid

25. Food poisoning outbreak investigation – first –
(a) Ban all food (b) Ignore (c) Give antibiotics to all (d) List cases & food histories

26. Fresh egg in water –
(a) Sinks (b) Floats (c) Dissolves (d) Stands upright at top

27. Veg packaged food symbol –
(a) Brown dot (b) Red triangle (c) Blue square (d) Green dot

? Level 3 – Advanced (NOT / Match / Assertion-Reason / Calculation)

28. Class I (natural) preservative NAHI –
(a) Vinegar (b) Sugar (c) Sodium benzoate (d) Salt

29. Assertion (A): Botulism ka toxin boiling se nasht ho sakta hai. Reason (R): Botulinum toxin heat-labile protein hai.
(a) Both A and R true, R explains A (b) Both A and R true, R does not explain A (c) A true, R false (d) A false, R true

30. Match: P. Khesari dal Q. Argemone oil R. Aflatoxin S. Ergot – 1. Lathyrism 2. Epidemic dropsy 3. Liver cancer 4. Ergotism
(a) P-1, Q-2, R-3, S-4 (b) P-4, Q-2, R-3, S-1 (c) P-1, Q-3, R-2, S-4 (d) P-2, Q-1, R-3, S-4

✓ Answer Key & Explanation

1-d Danger.
2-a Holder.
3-b Flash.
4-b Phosphatase.
5-b 1–6 hrs.
6-d Botulism.
7-d B. cereus.
8-d Argemone.
9-b Nitric.
10-b Benzoate.
11-c Osmotic.
12-c Radura.
13-c 2006.
14-d 7 June.
15-c Lactometer.
16-d Iodine.
17-d Baudouin.
18-c HCl.
19-d Pepper.
20-d FRK.
21-a Staph.
22-b 2 hrs.

23-a A & D.
24-b Sinks.
25-d Epidemiology.
26-a Fresh sinks.
27-d Green.
28-c Class II.
29-a Dono sahi, R reason hai.
30-a Sahi match.

9 National Nutrition Programmes

Level 1 – Direct

1. ICDS launched –
   (a) 2 October 1975 (b) 8 March 2018 (c) 15 August 1995 (d) 1985
2. ICDS ministry –
   (a) Health (b) Education (c) Women & Child Development (d) Agriculture
3. ICDS services –
   (a) 4 (b) 6 (c) 8 (d) 10
4. Supplementary nutrition child (6 m–6 yrs) –
   (a) 300 kcal, 8 g (b) 500 kcal, 12–15 g protein (c) 800 kcal, 25 g (d) 1000 kcal, 30 g
5. Pregnant/ lactating ICDS –
   (a) 300 kcal, 10 g (b) 500 kcal, 12 g (c) 600 kcal, 18–20 g protein (d) 800 kcal, 25 g
6. Severely malnourished child ICDS –
   (a) 500 kcal, 12 g (b) 600 kcal, 18 g (c) 800 kcal, 20–25 g (d) 1000 kcal
7. POSHAN Abhiyaan launched –
   (a) 1 April 2020 (b) 2 October 2014 (c) 8 March 2018 (d) 15 August 2016
8. POSHAN Abhiyaan launch place –
   (a) Varanasi (b) Jhunjhunu, Rajasthan (c) Ahmedabad (d) Delhi
9. Poshan Maah –
   (a) September (b) March (c) June (d) December
10. Mid-day meal started –
    (a) 1975 (b) 1995 (c) 2005 (d) 2013
11. MDM primary norm –
    (a) 300 kcal, 8 g (b) 450 kcal, 12 g protein (c) 600 kcal, 18 g (d) 700 kcal, 20 g
12. MDM upper primary norm –
    (a) 450 kcal, 12 g (b) 500 kcal, 15 g (c) 700 kcal, 20 g (d) 800 kcal, 25 g
13. Vitamin A prophylaxis programme –
    (a) 1970 (b) 1985 (c) 1992 (d) 2005
14. Anaemia Mukt Bharat –
    (a) 1970 (b) 2005 (c) 2013 (d) 2018
15. NIDDCP –
    (a) 1962 (b) 1975 (c) 1992 (d) 2006
16. NFSA –
    (a) 1997 (b) 2006 (c) 2013 (d) 2019
17. NFSA grains per person –
    (a) 2 kg/month (b) 5 kg/month (c) 10 kg/month (d) 35 kg/month
18. SDG 2 –
    (a) Good health (b) No poverty (c) Zero Hunger (d) Clean water

19. CFTRI located at –
(a) Delhi (b) Kolkata (c) Hyderabad (d) Mysuru

20. Best method to teach cooking –
(a) Lecture (b) Poster (c) Demonstration (d) Radio

? Level 2 – Priority & Application

21. Anganwadi worker job NAHI include karta –
(a) Pre-school education (b) Surgery (c) Supplementary nutrition (d) Growth monitoring

22. Mission Balam Sukham –
(a) Bihar roads (b) Delhi air (c) Kerala water (d) Gujarat child malnutrition

23. First 1000 days –
(a) Birth to 3 yrs (b) Pregnancy only (c) 0–1000 hours (d) Conception to 2 years

24. NFHS-5 under-5 stunting –
(a) 10 % (b) 19.3 % (c) 35.5 % (d) 67 %

25. NFHS-5 anaemia children 6–59 months –
(a) 25 % (b) 35.5 % (c) 67.1 % (d) 90 %

26. Most vulnerable group for malnutrition –
(a) Elderly men (b) Preschool children (c) Adolescent boys (d) Adult men

27. WHO global target – stunting reduction –
(a) 5 % (b) 10 % (c) 40 % (d) 100 %

? Level 3 – Advanced (NOT / Match / Assertion-Reason / Calculation)

28. ICDS beneficiary NAHI –
(a) Adult men (b) Lactating mothers (c) Pregnant women (d) Children 0–6

29. Assertion (A): Fortified rice PDS me diya ja raha hai. Reason (R): Fortification anaemia jaise micronutrient deficiencies ko kam karne ka cost-effective tarika hai.
(a) Both A and R true, R explains A (b) Both A and R true, R does not explain A (c) A true, R false (d) A false, R true

30. Match: P. ICDS Q. PM POSHAN R. NFSA S. POSHAN Abhiyaan – 1. 1975 2. 1995 3. 2013 4. 2018
(a) P-4, Q-2, R-3, S-1 (b) P-1, Q-2, R-3, S-4 (c) P-2, Q-1, R-3, S-4 (d) P-1, Q-3, R-2, S-4

✓ Answer Key & Explanation

1-a 1975.
2-c MWCD.
3-b 6.
4-b 500.
5-c 600.
6-c 800.
7-c 2018.
8-b Jhunjhunu.
9-a September.
10-b 1995.
11-b Primary.
12-c Upper primary.
13-a 1970.
14-d 2018.
15-c 1992.
16-c 2013.
17-b 5 kg.
18-c SDG 2.
19-d Mysuru.
20-c Demo.
21-b Not AWW.
22-d Gujarat.
23-d 1000 days.
24-c 35.5.

25-c 67.1.
26-b Preschool.
27-c 40 %.
28-a Men.
29-a Dono sahi, R reason hai.
30-b Sahi match.

10 GSSSB Pattern Mock Test – 100 MCQs (Full Nutrition Syllabus)



Mock test ke rules

Time: 90 minutes · **Marking**: +1 correct, −0.25 wrong · **Score guide**: 80+ Excellent · 65–79 Good · 50–64 Weak topics revise karo · < 50 Notes dobara padho.

? Mock Test – Part A (1–50)

1. Father of nutrition –
(a) McCarrison (b) Funk (c) Pasteur (d) Lavoisier

2. NIN located at –
(a) Hyderabad (b) Delhi (c) Pune (d) Mysuru

3. Body-building nutrient –
(a) Protein (b) Carbohydrate (c) Fat (d) Vitamin

4. Protective foods –
(a) Cereals (b) Vitamins & minerals (c) Fats (d) Sugars

5. Cheapest energy source –
(a) Carbohydrate (b) Vitamin (c) Protein (d) Fat

6. Kilocalorie heats 1 kg water by –
(a) 0.1 °C (b) 1 °C (c) 10 °C (d) 100 °C

7. BMR ↓ in –
(a) Hyperthyroidism (b) Infancy (c) Hypothyroidism (d) Fever

8. SDA of mixed diet –
(a) ~10 % (b) ~1 % (c) ~50 % (d) ~30 %

9. EAR covers –
(a) 25 % (b) 50 % population (c) 97.5 % (d) 100 %

10. Protein requirement adult (ICMR 2020) –
(a) 0.3 g/kg (b) 0.83 g/kg (c) 1.5 g/kg (d) 2 g/kg

11. Water requirement adult –
(a) ~5 ml/kcal (b) ~0.1 ml/kcal (c) ~1 ml/kcal (d) ~3 ml/kcal

12. Glycaemic index low –
(a) ≤ 55 (b) ≥ 70 (c) ≤ 90 (d) ≥ 100

13. Insoluble fibre –
(a) Cellulose (b) β-glucan (c) Pectin (d) Gums

14. Semi-essential amino acid –
(a) Lysine (b) Leucine (c) Arginine (d) Valine

15. Positive nitrogen balance –
(a) Burns (b) Fever (c) Growth & pregnancy (d) Starvation

16. MUFA rich oil –
(a) Coconut (b) Olive/ groundnut (c) Palm (d) Vanaspati

17. FSSAI trans fat limit –
(a) 0.1 % (b) 2 % (c) 10 % (d) 20 %

18. Desirable total cholesterol –
(a) < 100 mg/dl (b) < 200 mg/dl (c) < 300 mg/dl (d) < 400 mg/dl

19. LDL –
(a) Triglyceride (b) Bile (c) Good cholesterol (d) Bad cholesterol

20. Metabolic water maximum from –
(a) Carbohydrate (b) Vitamins (c) Fat (d) Protein

21. Main ICF cation –
(a) Calcium (b) Chloride (c) Sodium (d) Potassium

22. Xerophthalmia most characteristic sign –
(a) Fundus (b) Night blindness (c) Bitot's spots (d) Corneal scar

23. Keratomalacia –
(a) Vitamin A deficiency (emergency) (b) Vitamin D (c) Vitamin C (d) Vitamin K

24. Vitamin A total prophylactic doses –
(a) 3 (b) 5 (c) 9 (d) 12

25. Vitamin A 2nd dose –
(a) 1 lakh IU at 9 months (b) 2 lakh IU at 16–18 months (c) 5 lakh IU (d) 50,000 IU

26. Osteomalacia –
(a) Iron (b) Iodine (c) Vitamin D deficiency in adults (d) Vitamin C

27. Rickety rosary –
(a) Vitamin D (b) B12 (c) Vitamin C (d) Vitamin A

28. Vitamin D RDA adult –
(a) 50 IU (b) 100 IU (c) 600 IU (d) 5000 IU

29. Vitamin K factors –
(a) II, VII, IX, X (b) III, IV, XII (c) XI, XIII (d) I, V, VIII

30. Wet beriberi –
(a) Dermatitis (b) Diarrhoea (c) Heart failure (d) Neuritis

31. Magenta tongue –
(a) Vitamin C (b) Folate (c) Riboflavin (d) Niacin

32. Casal's necklace –
(a) Rickets (b) Beriberi (c) Pellagra (d) Scurvy

33. Niacin synthesised from –
(a) Glycine (b) Lysine (c) Methionine (d) Tryptophan

34. Megaloblastic anaemia –
(a) Iron (b) Vitamin K (c) Vitamin C (d) Folate/ B12 deficiency

35. Subacute combined degeneration –
(a) B1 (b) Folate (c) B12 (d) B6

36. Bleeding spongy gums –
(a) Pellagra (b) Beriberi (c) Rickets (d) Scurvy

37. Vitamin C ↑ absorption of –
(a) Calcium only (b) Iron (c) B12 (d) Fat

38. Infantile scurvy –
(a) Barlow's disease (b) Kwashiorkor (c) Rickets (d) Marasmus

39. Calcium absorption ↑ by –
(a) Fibre (b) Vitamin D (c) Phytates (d) Oxalates

40. Iron in haemoglobin –
(a) ~30 % (b) ~10 % (c) ~65–70 % (d) ~95 %

41. Haem iron absorbed –
(a) ~15–35 % (b) ~2 % (c) ~90 % (d) ~50 % always

42. Milk is poor source of –
(a) Iron & vitamin C (b) Protein (c) Calcium (d) Riboflavin

43. Reticulocyte response after iron –
(a) 1 day (b) 2 months (c) 5–10 days (d) 6 months

44. Severe anaemia pregnancy –
(a) Hb < 7 g/dl (b) Hb < 10 (c) Hb < 11 (d) Hb < 12

45. IFA pregnancy duration –
(a) 30 days (b) 90 days (c) 180 days (d) 365 days

46. Pink IFA tablet –
(a) Adolescents (b) Pregnant women (c) Infants (d) Children 5–9 yrs

47. Deworming day –
(a) 1 January (b) 5 May (c) 10 February & 10 August (d) 14 November

48. Iodine RDA pregnancy –
(a) 50 µg (b) 150 µg (c) 250 µg (d) 500 µg

49. Iodised salt at production –
(a) ≥ 5 ppm (b) ≥ 15 ppm (c) ≥ 30 ppm (d) ≥ 100 ppm

50. Best indicator of recent iodine intake –
(a) TSH only (b) Goitre rate (c) Hb (d) Urinary iodine

? Mock Test – Part B (51–100)

51. Skeletal fluorosis –
(a) Iron (b) High fluoride (c) Low fluoride (d) Iodine

52. Zinc deficiency –
(a) Goitre (b) Growth retardation, loss of taste (c) Scurvy (d) Rickets

53. Acrodermatitis enteropathica –
(a) Iron (b) Copper (c) Selenium (d) Zinc

54. Weight gain per week in 2nd–3rd trimester –
(a) ~0.1 kg (b) ~0.5 kg (c) ~2 kg (d) ~5 kg

55. Sudden excessive weight gain in pregnancy – suspect –
(a) Pre-eclampsia (b) Normal (c) Anaemia (d) GDM only

56. Pre-lacteal feeds –
(a) Honey essential (b) Necessary (c) Water essential (d) Harmful – avoid

57. Breast milk energy –
(a) ~5 kcal (b) ~70 kcal/100 ml (c) ~20 kcal (d) ~150 kcal

58. Whey:casein in breast milk –
(a) Only casein (b) Equal (c) More whey (d) More casein

59. Breastfeeding reduces in mother –
(a) Bone density (b) Iron needs to zero (c) PPH & breast cancer (d) Milk

60. LAM effective –
(a) Exclusive BF, amenorrhoea, < 6 months (b) Any feeding (c) Formula (d) After 1 year

61. Football hold useful –
(a) Never (b) Always worst (c) Only night (d) After LSCS/ twins

62. EBM fed with –
(a) Spoon only after 1 yr (b) Bottle (c) Cup/ paladai (d) Syringe IV

63. Complementary feeding 6–8 months frequency –
(a) 1 meal/week (b) 2–3 meals/day (c) Only breast milk (d) 6 meals

64. Cow's milk as main drink not before –
(a) 1 year (b) 2 years (c) 3 months (d) 6 months

65. Adolescent growth spurt needs ↑ –
(a) Calcium & iron (b) Only fat (c) Nothing (d) Only sugar

66. Elderly need extra –
(a) Fat (b) Sugar (c) Salt (d) Calcium, vitamin D, B12

67. Kwashiorkor meaning –
(a) Old age (b) Disease of deposed child (c) Night disease (d) Hunger disease

68. Marasmus face –
(a) Leonine (b) Mask face (c) Old-man face (d) Moon face

69. Fatty liver –
(a) Kwashiorkor (b) Scurvy (c) Marasmus (d) Rickets

70. SAM bilateral pitting oedema –
(a) Yes – SAM criterion (b) Normal (c) No (d) Only MAM

71. F-100 phase –
(a) Never (b) Day 1 (c) Stabilisation (d) Rehabilitation

72. RUTF energy –
(a) ~100 kcal (b) ~50 kcal (c) ~545 kcal/100 g (d) ~1000 kcal

73. IAP grade III –
(a) ≤ 50 % (b) 51–60 % (c) 61–70 % (d) 71–80 %

74. Underweight –
(a) Low height-for-age (b) Low weight-for-age (c) Low WFH (d) High BMI

75. NFHS-5 wasting –
(a) 3.4 % (b) 19.3 % (c) 35.5 % (d) 67 %

76. Asian normal BMI –
(a) 18.5–22.9 (b) 15–18 (c) 18.5–24.9 (d) 20–30

77. Broca's index –
(a) Height (cm) − 100 (b) Weight − 100 (c) Weight/ height (d) Height × 2

78. Short-term protein marker –
(a) Albumin (b) Globulin (c) Prealbumin (d) Ferritin

79. Clear liquid diet duration max –
(a) Lifelong (b) 2 weeks (c) 24–48 hrs (d) 1 month

80. Hypoglycaemia conscious –
(a) Wait 1 hr (b) 15 g fast sugar, recheck 15 min (c) Insulin (d) Water

81. CHF diet –
(a) Low sodium, fluid restriction (b) High fluids (c) High fat (d) High salt

82. CKD potassium leaching –
(a) Soak/ boil vegetables, discard water (b) Add salt (c) Raw eating (d) Fry

83. Calcium oxalate stones – avoid –
(a) Rice (b) Water (c) Spinach & chocolate (d) Milk restriction

84. Hepatic encephalopathy – preferred protein –
(a) Vegetable & dairy (b) Red meat (c) None ever (d) Only eggs

85. Peptic ulcer – avoid –
(a) Bananas (b) Water (c) Spicy food, caffeine, alcohol (d) Rice

86. Diarrhoea child – feeding –
(a) Only water (b) Continue feeding + ORS + zinc (c) Only juice (d) Stop all food

87. Curreri formula –
(a) Fluid in burns (b) BMI (c) Protein quality (d) Burns energy

88. PKU avoid sweetener –
(a) Stevia (b) Aspartame (c) Sugar (d) Jaggery

89. TPN route –
(a) Oral (b) Subcutaneous (c) Central vein (d) IM

90. Lipid emulsion –
(a) Yellow (b) Milky white (c) Clear (d) Red

91. Levodopa – avoid with dose –
(a) High protein meal (b) Rice (c) Water (d) Fruit

92. Lithium – maintain –
(a) Consistent salt & fluid (b) Low salt (c) No water (d) High sugar

93. UHT milk –
(a) 63 °C × 30 min (b) 72 °C × 15 s (c) 100 °C × 10 min (d) 135–140 °C × 2–4 s

94. Milk bacterial load test –
(a) Lactometer (b) Iodine (c) Methylene blue reduction (d) Phosphatase

95. Staph food poisoning food –
(a) Raw seafood (b) Canned food (c) Honey (d) Milk products/ sweets

96. Calcium carbide –
(a) Artificial fruit ripening (banned) (b) Preservative (c) Colour (d) Sweetener

97. AGMARK –
(a) Gold (b) Organic only (c) Agricultural products (d) Electrical

98. +F staples –
(a) Soft drinks (b) Sugar, tea (c) Biscuits (d) Salt, oil, milk, atta, rice

99. ICDS – Anganwadi worker serves –
(a) Only schools (b) Only men (c) Children 0–6, pregnant & lactating women (d) Only elderly

100. National Nutrition Week –

(a) 1–7 December (b) 1–7 August (c) 14–20 November (d) 1–7 September

Answer Key – Mock Test (with explanation)

1-d Lavoisier.
2-a Hyderabad.
3-a Protein.
4-b Protective.
5-a CHO.
6-b Definition.
7-c Hypothyroid.
8-a 10 %.
9-b EAR.
10-b 0.83.
11-c 1 ml/kcal.
12-a Low GI.
13-a Cellulose.
14-c Arginine.
15-c Positive.
16-b MUFA.
17-b 2 %.
18-b 200.
19-d LDL.
20-c Fat.
21-d K.
22-c Bitot's.
23-a A.
24-c 9.
25-b 2 lakh.
26-c Adults.
27-a D.
28-c 600.
29-a 1972.
30-c Wet.
31-c B2.
32-c Niacin.
33-d Tryptophan.
34-d Macrocytic.
35-c B12.
36-d C.
37-b Iron.
38-a Barlow.
39-b D.
40-c Hb.
41-a Haem.
42-a Milk.
43-c Response.
44-a Severe.
45-c 180.
46-d Pink.
47-c NDD.
48-c 250.
49-c 30.
50-d UIC.
51-b Fluoride.
52-b Zinc.
53-d Zinc.
54-b 0.5.
55-a PIH.
56-d Avoid.
57-b 70.
58-c Whey.
59-c Benefits.
60-a LAM.
61-d Football.
62-c Cup.
63-b 2–3.
64-a 1 yr.
65-a Growth.
66-d Elderly.
67-b Ga.
68-c Marasmus.
69-a Kwashiorkor.
70-a SAM.
71-d Catch-up.
72-c RUTF.
73-b IAP.
74-b WFA.
75-b 19.3.
76-a Asian.
77-a Broca.
78-c Prealbumin.
79-c Short.
80-b Rule of 15.
81-a CHF.
82-a Leaching.
83-c Oxalate.
84-a Veg/dairy.
85-c Ulcer.
86-b Feeding.
87-d Burns.
88-b Aspartame.
89-c Central.
90-b Lipid.
91-a Protein.
92-a Lithium.
93-d UHT.
94-c MBRT.
95-d Staph.
96-a Banned.
97-c AGMARK.
98-d +F.

99-c ICDS.
100-d Sept.

— Nutrition · GSSSB Staff Nurse Preparation · Padhte raho, revise karte raho! —