

Flowform[®] Eco-Technology results



Findings:

Treating animal effluent with Flowform technology resulted in:

- transformed dairy shed effluent into liquid fertilizer that can be sprayed out over paddocks
- lack of anaerobic decomposing smell
- no temporary nitrogen starvation of the foliage and consequently a nutrient deficiency for the animals
- faster absorption

- faster return of animals to paddocks
- healthy and clean pasture

Cow slurry treatment test

November 2012, Bonyhád, Hungary

During the test 0,8 m³ slurry has been treated with a Glonn 5 Flowform over a continuous 22 hour period, that resulted in the decrease of the COD (chemical oxygen demand) by 17% (in total of 4766 mg/l), decrease of total Carbon content by 23% (in total of 2277 mg/l) and decrease of total Nitrogen by 13% (in total of 197 mg/l).

The total COD decrease and 2/3 of the total carbon decrease measured has happened in the first 4-8 hours that justifies the initial very active aerobic process also leading to loss of smell after the first couple of hours. The nitrification process did not start due to the relatively short test period and low temperatures around 10 C, therefore nitrate content could not be measured.



1 -

Piggery le Rouget du Porc, France

Dear Ian

Re Flowform effectiveness

In 2009 we supplied Flowform to a commercial piggery called "le Rouget du Porc", in Nièvre, central France, owned by Patrick Ravery. His pigs suffered from chronic skin ailments, like eczema and itchy inflammation.

He has had good results after using Flowform water for the pigs, with much of the illnesses clearing up. The pigs were drinking more water as well.

11th June 2015

Best wishes,
Michaël

Atelier Coralis
18330 Saint Laurent
France
+33 0248 51 5818
www.fontainecoralis.com

Chicken Abattoir

Byron Bay, NSW. Australia

This abattoir slaughtered 26,000 chickens three to five times per week. The washing from the slaughter room went through a clarifier into a 4.3 megalitre pond. From there water is used for irrigation on an onsite turf farm. The stench caused neighbors to complain and the Environmental Protection Agency were about to close them down when the Flowform technology was set up. Dissolved Oxygen (DO) readings were zero.

Three sets of Vortex 7 Flowforms were installed and within two weeks of continuous running the pond's dissolved oxygen was 2.3 ppm. By the end of the sixth week DO reached 6.3 ppm. The neighbors rang to inquire where the smells had gone.

The Environmental Protection Agency has extended their license, with the added construction of gravel filters and reed beds.

Agriculture related measurement and observation results:

The influence of Flowform treatment on lettuce growth

Summer 2004

Healing Water Foundation

Forest Row

United Kingdom



Lettuce was grown in the field in the summer of 2004 to investigate the influence of Flowform treatment on irrigation water. The experiment included three lettuce groups where the Vortex Flowform model was used to treat water that has applied: (i) only at the greenhouse stage; (ii) during the whole growth period. Both the above were compared with (iii) controls treated with tap water.

The investigation included a series of observations (of colour, shape, leaf sequence, general appearance, germination rate, leaf size etc.) conducted once a week. After harvesting the lettuces were observed, tasted, weighed and tested for pH, and differences were detected in most parameters.

The weight of lettuce irrigated with Flowform-treated water increased by 32%, appeared more vibrant in colour and texture and had more taste. The lettuce that was grown in Flowform-treated water only at the greenhouse stage had characteristics that mostly combined those of the two other groups, though were closer to the lettuce treated throughout with Flowform water.

- 2 -

	FF Flowform water	FT Flowform water at	TT Tap water control
--	-------------------	----------------------	----------------------

		greenhouse stage	
Number of leaves	8,27 +/- 0,54 [SE]	8,20 +/- 0,34 [SE]	7,40 +/- 0,36 [SE]
Mean pH value of soil	6,8	7,1	6,8
Mean root length [cm]	21,67 +/- 0,73 [SE]	18,67 +/- 0,88 [SE]	17,83 +/- 1,42 [SE]
Mean weight [g]	178,4 +/- 8,6 [SE]	168,67 +/- [SE]	134,7 +/- 7,8 [SE]
Total weight [g]	7137	6748	5388
Increase in total weight compared to control TT [%]	32,5%	25,2%	-
Soluble content	1,44 +/- 0,19 [SE]	1,68 +/- 0,14 [SE]	2,04 +/- 0,17 [SE]
<i>(SE: Standard Error. Underlined numbers indicate a statistically significant difference with respect to the control)</i>			

The general impression was that the FF lettuces were larger in size and had more leaves than the FT samples, the TT plants being the smallest of all. No other apparent differences in color or texture could be detected. Roots appeared to be thicker in FF and FT plants, and the soil seemed to be harder to remove with tap watered plants.

Influence of Flowform water on wheat growth

Summer 2004

Healing Water Foundation

Forest Row

United Kingdom

This research project aimed to examine the effect of Flowform-treated water on plant growth. The first stages of cress germination and growth were examined, as it is known that during initial stages of growth the plant is very sensitive to its surroundings.

A number of researchers have established that the different phases of the moon, as well as other planetary influences, have a strong effect on the growth of plants (Lawrence Edwards 1993, Georg Schimdt 1984, Theodor Schwenk 1967, Maria Thun 2003).

Research by Freya Schikorr (1990) suggested that it is possible to enhance lunar influences on plant growth by treating water rhythmically in Flowform cascades.

It was found that Flowform treatment promoted root growth especially, and that the rhythmical movement of the water seems to have a positive effect on biological processes resulted from its enhanced sensitivity.

The cress-growth experiments were carried out using a method developed by Hiscia Cancer Institute in Arlesheim, Switzerland. A tap water control was used to compare with Flowform related results. A 38 vessel stackable Flowform cascade was used during the experiment.


The averages shown were calculated by using the germinated seeds only. The difference between Flowform treated and control samples was more pronounced in root (+9,5%) than in shoot lengths (+5,6%). However, this

increase in growth was not constant over the period investigated. The lunar phases were also scrutinized as possible influencing factors causing these fluctuations.

The experiment indicated that seedling growth shows a pattern according with moon phases.

- 3 -


Increase in growth compared to
Mean increase in cress-seedling
growth
10,00%



tap water control

Rotating table with seedlings
8,00% 6,00% 4,00% 2,00%
0,00%

shoot root total length



Tap water control Flowform treated

