The relationship of extracellular fields - neural oscillations and spikes

Cong Wang, Hoi Ming Ken Yip, Vicky Zhu, Weihao Sheng, Yongxiang Xiao Murky Stoats/ Lucky Coi 23/07/2021












Synchronization of unit activity: a key ingredient of neural oscillations

Timing within theta cycles




Our research question: Which cell type's spiking activities are better predictors of neural oscillations.


Our hypothesis: interneurons can better reconstruct theta oscillation than principal neurons.



Steinmetz dataset




Trial-averaged firing rates






Steinmetz, Nicholas A., et al. "Distributed coding of choice, action and engagement across the mouse brain." Nature 576.7786 (2019): 266-273.

Steinmetz dataset








Steinmetz, Nicholas A., et al. "Distributed coding of choice, action and engagement across the mouse brain." Nature 576.7786 (2019): 266-273.

RNN architecture & model evaluation



Spks

Noise

MD VISam

40

50

CA1

30


Brain Areas

frequency


DĠ




RNN architecture & model evaluation




Hyper-connected and under-connected training weights




Negatively hyper-connected neurons (n=20)
Positively hyper-connected neurons (n=20)
Under-connected neurons (n=20)





Principal neurons (PNs) and Interneurons (INs) training weights





Brain areas

Conclusions

- Developed RNN model with linear recurrent unit
- spiking activity -> LFP in θ band
- Findings:
 - Hyper-connected cells > under-connected cells
 - Interneurons > principal neurons



Acknowledgement and great teamwork!

Mentor



Andrea Hasenstaub

TA



Kai Chen



Nitin Anisetty

Our team



Cong Wang



Hoi Ming Ken Yip




Yongxiang Xiao




Vicky Zhu



Weihao Sheng



Unit functional connectivity vs. Cell types





Single-neuron firing rate vs. training weights

