

Generalized Rainbow Configurations

Mike Desgrottes¹, David Soukup¹, Renjun Zhu¹

August 3, 2017

¹This work was supported in part by NSF Grant DMS-1559911 

Table of Contents

1 Introduction

- Background
 - Rainbow Configuration
 - Definition
- Literature Review

2 Our Results

- Finite Case
 - FCT
 - Merging
 - Corollary
- Infinite Case
 - Pathological Coloring
 - Measure
 - ICT
 - Corollary
- w-subrainbow

Background

- We study Anti-Ramsey problems (rainbow problems). Ramsey Theory studies monochromatic substructure within a structure.

Background

- We study Anti-Ramsey problems (rainbow problems). Ramsey Theory studies monochromatic substructure within a structure.
 - Classical Ramsey Theory - Schur's Theorem:
Coloring the natural numbers into finite number of colors, then there must a color class contain a triple with $(x, y, x + y)$

Background

- We study Anti-Ramsey problems (rainbow problems). Ramsey Theory studies monochromatic substructure within a structure.
 - Classical Ramsey Theory - Schur's Theorem:
Coloring the natural numbers into finite number of colors, then there must a color class contain a triple with $(x, y, x + y)$
- **Goal:** find a condition on the coloring to guarantee a rainbow configuration.

What's a Rainbow Configuration?

Suppose we have a set $X = \{0, 1, 2, 3, 4, 5\}$

Q: Can you find $(x, y, x + y)$ such that x , y , and $x + y$ are different colors?

What's a Rainbow Configuration?

Suppose we have a set $X = \{0, 1, 2, 3, 4, 5\}$

Q: Can you find $(x, y, x + y)$ such that x , y , and $x + y$ are different colors?

Hint:

- $1 + 2 = 3$
- $2 + 3 = 5$
- $0 + 1 = 1$
- $2 + 2 = 4$
- $1 + 4 = 5$
- ...

What's a Rainbow Configuration?

Suppose we have a set $X = \{0, 1, 2, 3, 4, 5\}$

Q: Can you find $(x, y, x + y)$ such that x , y , and $x + y$ are different colors?

Hint:

- $1 + 2 = 3$
- $2 + 3 = 5$
- $0 + 1 = 1$
- $2 + 2 = 4$
- $1 + 4 = 5$

...

We also see $(1, 4, 5)$ is a triple with all distinct color elements.

Definition

Definition

A **coloring** of X is a function $f : X \rightarrow \mathcal{C}$ of colors. Here the preimages $\{f^{-1}(c)\}$ are the color classes from X .

Definition

Definition

A **coloring** of X is a function $f : X \rightarrow \mathcal{C}$ of colors. Here the preimages $\{f^{-1}(c)\}$ are the color classes from X .

Definition

The **configuration space** is the set $\mathcal{F} = \{(x_1, x_2, \dots, x_k) \in X^k\}$ for $k \geq 3$

Definition

Definition

A **rainbow tuple** is an element $f \in \mathcal{F}$ such that the elements $\pi_1(f), \pi_2(f) \dots \pi_k(f)$ all belong to distinct color classes. If no rainbow tuples exist, then our configuration space $\{X, \mathcal{F}, c\}$ is **rainbow-free**.

Definition

Definition

A **rainbow tuple** is an element $f \in \mathcal{F}$ such that the elements $\pi_1(f), \pi_2(f) \dots \pi_k(f)$ all belong to distinct color classes. If no rainbow tuples exist, then our configuration space $\{X, \mathcal{F}, c\}$ is **rainbow-free**.

- From our previous example, $X = \{0, 1, 2, 3, 4, 5\}$. The color classes are $\{2, 4\}$, $\{0, 5\}$, $\{1, 3\}$.

Definition

Definition

A **rainbow tuple** is an element $f \in \mathcal{F}$ such that the elements $\pi_1(f), \pi_2(f) \dots \pi_k(f)$ all belong to distinct color classes. If no rainbow tuples exist, then our configuration space $\{X, \mathcal{F}, c\}$ is **rainbow-free**.

- From our previous example, $X = \{0, 1, 2, 3, 4, 5\}$. The color classes are $\{2, 4\}$, $\{0, 5\}$, $\{1, 3\}$.
- Our configuration space $\mathcal{F} = \{(x, y, x + y) \in X^3\}$

Definition

Definition

A **rainbow tuple** is an element $f \in \mathcal{F}$ such that the elements $\pi_1(f), \pi_2(f) \dots \pi_k(f)$ all belong to distinct color classes. If no rainbow tuples exist, then our configuration space $\{X, \mathcal{F}, c\}$ is **rainbow-free**.

- From our previous example, $X = \{0, 1, 2, 3, 4, 5\}$. The color classes are $\{2, 4\}$, $\{0, 5\}$, $\{1, 3\}$.
- Our configuration space $\mathcal{F} = \{(x, y, x + y) \in X^3\}$
- Our coloring is **not** rainbow-free since there are rainbow triples, $(1, 4, 5)$, $(2, 3, 5)$, etc.

Known Results

Theorem: Ceja, Cook, and Hayden (2016)

If we let $X = \mathbb{Z}/p$, for large prime p , and $k \geq 3$, then for every partitioning of X into k colors with each of size $\lceil \frac{p}{k} \rceil$ or $\lfloor \frac{p}{k} \rfloor$, there must exist a rainbow tuple of the form $(x_1, x_2, x_1 + x_2)$.

Known Results

Theorem: Ceja, Cook, and Hayden (2016)

If we let $X = \mathbb{Z}/p$, for large prime p , and $k \geq 3$, then for every partitioning of X into k colors with each of size $\lceil \frac{p}{k} \rceil$ or $\lfloor \frac{p}{k} \rfloor$, there must exist a rainbow tuple of the form $(x_1, x_2, x_1 + x_2)$.

Theorem: Schönheim (1990)

For every 3-coloring of $1, 2, 3, \dots, n$, with every color class with at least $n/4$ elements, there exists a rainbow tuple $(x_1, x_2, x_1 + x_2)$.

Theorem: Senger (2017)

Fix $c > 0$. Given a coloring of \mathbb{F}_q^2 , with q sufficiently large, where no color class has size $\geq cq^2$, if unit equilateral triangles exist in \mathbb{F}_q^2 then there must be a rainbow unit equilateral triangle.

Constant M

Definition

Given $\mathcal{F} \subset X^k$, we define $M_{i,j}$ as the constant such that for any $x = (x_1, x_2, \dots, x_k) \in \mathcal{F}$, we have $|\{y \in \mathcal{F} : y_i = x_i, y_j = x_j\}| \leq M_{i,j}$

$$M = \sum_{i < j} M_{i,j}$$

- ① Given $\mathcal{F} = \{(x, y, x + y) \in X^3\}$, $M = M_{1,2} + M_{1,3} + M_{2,3} = 3$
 - If we fix any two coordinates, then we can uniquely determine the other coordinate. So, there's only one tuple in \mathcal{F} , namely $M_{1,2} = M_{2,3} = M_{1,3} = 1$

Constant M

Definition

Given $\mathcal{F} \subset X^k$, we define $M_{i,j}$ as the constant such that for any $x = (x_1, x_2, \dots, x_k) \in \mathcal{F}$, we have $|\{y \in \mathcal{F} : y_i = x_i, y_j = x_j\}| \leq M_{i,j}$

$$M = \sum_{i < j} M_{i,j}$$

- ① Given $\mathcal{F} = \{(x, y, x + y) \in X^3\}$, $M = M_{1,2} + M_{1,3} + M_{2,3} = 3$
 - If we fix any two coordinates, then we can uniquely determine the other coordinate. So, there's only one tuple in \mathcal{F} , namely $M_{1,2} = M_{2,3} = M_{1,3} = 1$
- ② Given $\mathcal{F} = \{(x, y, x + y, xy) \in X^4\}$,
 $M = M_{1,2} + M_{1,3} + M_{1,4} + M_{2,3} + M_{2,4} + M_{3,4} = 5 \cdot 1 + 2 = 7$

Finite Case Theorem(FCT)

Theorem

Let X be a finite set of size n , and $\mathcal{F} \subset X^k$ be a set of k -tuples. If no color class has size $\geq Cn$, where $|\mathcal{F}| \geq Dn^2$, and $C < \frac{2D}{9M}$, then there must be a rainbow k -tuple in \mathcal{F} .

Finite Case Theorem(FCT)

Theorem

Let X be a finite set of size n , and $\mathcal{F} \subset X^k$ be a set of k -tuples. If no color class has size $\geq Cn$, where $|\mathcal{F}| \geq Dn^2$, and $C < \frac{2D}{9M}$, then there must be a rainbow k -tuple in \mathcal{F} .

Sketch of the proof:

- 1 Assume it is a rainbow-free.
- 2 Merge color classes until attaining a minimum and maximum bound on all the color class sizes.

Finite Case Theorem(FCT)

Theorem

Let X be a finite set of size n , and $\mathcal{F} \subset X^k$ be a set of k -tuples. If no color class has size $\geq Cn$, where $|\mathcal{F}| \geq Dn^2$, and $C < \frac{2D}{9M}$, then there must be a rainbow k -tuple in \mathcal{F} .

Sketch of the proof:

- 1 Assume it is a rainbow-free.
- 2 Merge color classes until attaining a minimum and maximum bound on all the color class sizes.
- 3 Use $|\mathcal{F}|$ bounds to get a lower bound on the number of color classes.

Finite Case Theorem(FCT)

Theorem

Let X be a finite set of size n , and $\mathcal{F} \subset X^k$ be a set of k -tuples. If no color class has size $\geq Cn$, where $|\mathcal{F}| \geq Dn^2$, and $C < \frac{2D}{9M}$, then there must be a rainbow k -tuple in \mathcal{F} .

Sketch of the proof:

- 1 Assume it is a rainbow-free.
- 2 Merge color classes until attaining a minimum and maximum bound on all the color class sizes.
- 3 Use $|\mathcal{F}|$ bounds to get a lower bound on the number of color classes.
- 4 We use what we know about the color classes' size to reach a contradiction.

Merging

Definition

Define a coloring λ to be a **merging** of λ' if x, y are the same color under λ' , then they also have the same color under λ .




Figure: Merge color classes can destroy but not create rainbow configurations.

Proof.

- We merge the color classes until every color class has size between $\frac{1}{2}Cn$ and $\frac{3}{2}Cn$. Let s denote the number of color class after this process.

Proof.

- We merge the color classes until every color class has size between $\frac{1}{2}Cn$ and $\frac{3}{2}Cn$. Let s denote the number of color class after this process.
- Let n_i denote the size of color class i . Recall that M counts the number of tuples in \mathcal{F} that share a pair of coordinates. So there are at most Mn_i^2 elements in \mathcal{F} with at least two coordinates from color class i .

Proof.

- We merge the color classes until every color class has size between $\frac{1}{2}Cn$ and $\frac{3}{2}Cn$. Let s denote the number of color class after this process.
- Let n_i denote the size of color class i . Recall that M counts the number of tuples in \mathcal{F} that share a pair of coordinates. So there are at most Mn_i^2 elements in \mathcal{F} with at least two coordinates from color class i .
- $|\mathcal{F}| \leq M \sum_{i=1}^s n_i^2 \leq M \sum_{i=1}^s \left(\frac{3}{2}Cn\right)^2 \leq s \frac{9M}{4} C^2 n^2$

Proof.

- We merge the color classes until every color class has size between $\frac{1}{2}Cn$ and $\frac{3}{2}Cn$. Let s denote the number of color class after this process.
- Let n_i denote the size of color class i . Recall that M counts the number of tuples in \mathcal{F} that share a pair of coordinates. So there are at most Mn_i^2 elements in \mathcal{F} with at least two coordinates from color class i .
- $|\mathcal{F}| \leq M \sum_{i=1}^s n_i^2 \leq M \sum_{i=1}^s \left(\frac{3}{2}Cn\right)^2 \leq s \frac{9M}{4} C^2 n^2$
it implies $s \geq \frac{4}{9M} \frac{1}{C^2} \frac{|\mathcal{F}|}{n^2} \geq \frac{4D}{9MC^2}$

Proof.

- We merge the color classes until every color class has size between $\frac{1}{2}Cn$ and $\frac{3}{2}Cn$. Let s denote the number of color class after this process.
- Let n_i denote the size of color class i . Recall that M counts the number of tuples in \mathcal{F} that share a pair of coordinates. So there are at most Mn_i^2 elements in \mathcal{F} with at least two coordinates from color class i .
- $|\mathcal{F}| \leq M \sum_{i=1}^s n_i^2 \leq M \sum_{i=1}^s \left(\frac{3}{2}Cn\right)^2 \leq s \frac{9M}{4} C^2 n^2$
it implies $s \geq \frac{4}{9M} \frac{1}{C^2} \frac{|\mathcal{F}|}{n^2} \geq \frac{4D}{9MC^2}$
- Finally, X has at least $\left(\frac{1}{2}Cn\right) \cdot \frac{4D}{9MC^2} > n$ elements, for $C < \frac{2D}{9M}$

Contradiction!

Finite Case Corollary

Corollary 1

Suppose we have a coloring of a finite abelian group, G . If all color class has size $< \frac{2}{27}|G|$ then there must be a rainbow triple of the form $(x, y, x + y)$.

Finite Case Corollary

Corollary 1

Suppose we have a coloring of a finite abelian group, G . If all color class has size $< \frac{2}{27}|G|$ then there must be a rainbow triple of the form $(x, y, x + y)$.

- Applying FCT, here $X = G$, $k = 3$, and $\mathcal{F} = \{(x, y, x + y) \in G^3 | x, y \in G\}$.

Finite Case Corollary

Corollary 1

Suppose we have a coloring of a finite abelian group, G . If all color class has size $< \frac{2}{27}|G|$ then there must be a rainbow triple of the form $(x, y, x + y)$.

- Applying FCT, here $X = G$, $k = 3$, and $\mathcal{F} = \{(x, y, x + y) \in G^3 | x, y \in G\}$.
- so $M = 3$, $|\mathcal{F}| = n^2$

Finite Case Corollary

Corollary 1

Suppose we have a coloring of a finite abelian group, G . If all color class has size $< \frac{2}{27}|G|$ then there must be a rainbow triple of the form $(x, y, x + y)$.

- Applying FCT, here $X = G$, $k = 3$, and $\mathcal{F} = \{(x, y, x + y) \in G^3 | x, y \in G\}$.
- so $M = 3$, $|\mathcal{F}| = n^2$
- thus, $D = 1$, and $C < \frac{2D}{9M} = \frac{2}{27}$

Corollary 2

If we color \mathbb{F}_q such that each color class has size $< \frac{(2-o(1))}{63}q$, then there must be a rainbow quadruple of the form $(x, y, x + y, xy)$.

Corollary 2

If we color \mathbb{F}_q such that each color class has size $< \frac{(2-o(1))}{63}q$, then there must be a rainbow quadruple of the form $(x, y, x + y, xy)$.

Corollary 3

If we color a finite (additive) abelian group G , with no (nonidentity) element of order $< k$, then if there are no rainbow k -arithmetic progressions, at least one color class has size $\geq \frac{2}{9\binom{k}{2}}|G|$.

Corollary 2

If we color \mathbb{F}_q such that each color class has size $< \frac{(2-o(1))}{63}q$, then there must be a rainbow quadruple of the form $(x, y, x + y, xy)$.

Corollary 3

If we color a finite (additive) abelian group G , with no (nonidentity) element of order $< k$, then if there are no rainbow k -arithmetic progressions, at least one color class has size $\geq \frac{2}{9\binom{k}{2}}|G|$.

Corollary 4

If we color $[1..n]$ such that there are no rainbow triples of the form $(x, y, x + y)$, then at least one color class has size $\geq \frac{1-o(1)}{27}n$.

Pathological Coloring

Example

Suppose we have the following color classes on \mathbb{Q} , where $a, b \in \mathbb{Z}$ for $b \neq 0$

- color class 0: $\{\mathbb{Z}\} \cup \{\frac{a}{b} \in \mathbb{Q} : 2 \nmid a, b\}$

Pathological Coloring

Example

Suppose we have the following color classes on \mathbb{Q} , where $a, b \in \mathbb{Z}$ for $b \neq 0$

- color class 0: $\{\mathbb{Z}\} \cup \{\frac{a}{b} \in \mathbb{Q} : 2 \nmid a, b\}$
- color class 1: $\{\frac{a}{2^i b} : 2 \nmid a, b\}$

Pathological Coloring

Example

Suppose we have the following color classes on \mathbb{Q} , where $a, b \in \mathbb{Z}$ for $b \neq 0$

- color class 0: $\{\mathbb{Z}\} \cup \{\frac{a}{b} \in \mathbb{Q} : 2 \nmid a, b\}$
- color class 1: $\{\frac{a}{2^1 b} : 2 \nmid a, b\}$
- color class 2: $\{\frac{a}{2^2 b} : 2 \nmid a, b\}$

Pathological Coloring

Example

Suppose we have the following color classes on \mathbb{Q} , where $a, b \in \mathbb{Z}$ for $b \neq 0$

- color class 0: $\{\mathbb{Z}\} \cup \{\frac{a}{b} \in \mathbb{Q} : 2 \nmid a, b\}$
- color class 1: $\{\frac{a}{2^1 b} : 2 \nmid a, b\}$
- color class 2: $\{\frac{a}{2^2 b} : 2 \nmid a, b\}$
- ...
- color class i : $\{\frac{a}{2^i b} : 2 \nmid a, b\}$

If $x \in$ color class i , and $y \in$ color class j , then $x + y \in$ color class $\max(i, j)$. Here we have infinitely many classes, but still **rainbow-free** of the form $(x, y, x + y)$.

Pathological Coloring

Example

Suppose we color vector space \mathbb{R} over \mathbb{Q} as following, and let $\{x_i\}_{i \in I}$ be a well-ordered basis.

- Define A_j to be the set of linear combinations $\sum_{i \in I} c_i x_i$ with all but finitely many c_i equal to zero and $j = \min(i : c_i \neq 0)$

Pathological Coloring

Example

Suppose we color vector space \mathbb{R} over \mathbb{Q} as following, and let $\{x_i\}_{i \in I}$ be a well-ordered basis.

- Define A_j to be the set of linear combinations $\sum_{i \in I} c_i x_i$ with all but finitely many c_i equal to zero and $j = \min(i : c_i \neq 0)$
- The color classes are A_j . We see that if $a_i \in A_i$ and $a_j \in A_j$, then $a_i + a_j, a_i - a_j, a_j - a_i \in A_{\min(i,j)}$

Pathological Coloring

Example

Suppose we color vector space \mathbb{R} over \mathbb{Q} as following, and let $\{x_i\}_{i \in I}$ be a well-ordered basis.

- Define A_j to be the set of linear combinations $\sum_{i \in I} c_i x_i$ with all but finitely many c_i equal to zero and $j = \min(i : c_i \neq 0)$
- The color classes are A_j . We see that if $a_i \in A_i$ and $a_j \in A_j$, then $a_i + a_j, a_i - a_j, a_j - a_i \in A_{\min(i,j)}$

Here we have uncountably many color classes, but again it's a **rainbow-free** of the form $(x, y, x + y)$ in this coloring.

What is more do we need?

Measure!!!

Before we talk about the size of the set X , we can do that earlier because our sets were finite. However, now we can have base space and color classes be **infinite**. For that, we need a way to talk about the "size" of an infinite case.

Measure Space to the Rescue

Measure Space

- ① X : finite or infinite set
- ② Σ : **Measurable sets**
 - a collection of subsets of X , also known as σ -algebra, which is closed under its complement, countable unions, and intersections;
 - contains X , and \emptyset .
- ③ μ : **Measure**, a function map $\Sigma \rightarrow \mathbb{R}^+ \cup \{0, \infty\}$ such that:
 - $\mu(\emptyset) = 0$;
 - $\mu(\bigcup E_i) = \sum \mu(E_i)$, where $E_i \cap E_j = \emptyset$

(X, Σ, μ) is called measure space.

X is Finite: $\mu(X) < \infty$

Definition

Technique: **merging color classes**

Definition

Technique: **merging color classes**

Challenge: we don't want to merge measurable color classes and get an non-measurable color classes 😞

Definition

Technique: **merging color classes**

Challenge: we don't want to merge measurable color classes and get an non-measurable color classes 😞

Solve: we need to have a countably many different color classes. 😊

Definition

Technique: **merging color classes**

Challenge: we don't want to merge measurable color classes and get an non-measurable color classes 😞

Solve: we need to have a countably many different color classes. 😊

Definition

Let (X, μ) be a measure space, a coloring of X is **tractable** if every color class is a measurable subset of X , and there are only at most countably many color classes.

Infinite Case Theorem (ICT)

Let (X, μ) be a **finite measure space**, $n = \mu(X) < \infty$, and $\mathcal{F} \subset X^k$ be a set of k -tuples. If we have a **tractable coloring** where no color class has measure $\geq Cn$, with $m(\mathcal{F}) \geq D \binom{k}{2} n^2$, and $C < \frac{2D}{9M}$, then there must be a rainbow k -tuple in \mathcal{F} .

Infinite Case Theorem (ICT)

Let (X, μ) be a **finite measure space**, $n = \mu(X) < \infty$, and $\mathcal{F} \subset X^k$ be a set of k -tuples. If we have a **tractable coloring** where no color class has measure $\geq Cn$, with $m(\mathcal{F}) \geq D \binom{k}{2} n^2$, and $C < \frac{2D}{9M}$, then there must be a rainbow k -tuple in \mathcal{F} .

$$m(\mathcal{F}) = \sum_{i < j} m_{ij}(\mathcal{F})$$

where

$$m_{ij}(\mathcal{F}) = \iint |\{y \in \mathcal{F} : x_i = y_i, x_j = y_j\}| dx_i dx_j$$

Note: The finite case is considered μ as counting measure.

Merging

Remark

Suppose we have an original coloring, λ_0 . Let Λ be the set of all coloring that are merging of λ_0 and have no color classes of size $> Cn$.

Here, apply Zorn's lemma so that we can talk about maximum and minimum measure of the color class.

Sketch of ICT Proof

Proof.

- 1 Similar to the finite case, we start with assuming that our coloring is rainbow-free.
- 2 Merge color classes until the measure of each the color class, $\mu(A)$, is between $\frac{1}{2}Cn$ and $\frac{3}{2}Cn$
- 3 We obtain contradiction based on the number of coloring.



Corollary 1

Fix a probability measure μ on the unit circle in the complex plane. If we color the circle with at least 14 equally sized μ -measurable color classes, there must be a rainbow triple of the form (x, y, xy)

- We note that the measure space is finite since $\mu(X) = 1$.
- $\mathcal{F} = \{(x, y, xy) \in X^3 \mid x, y \in X\}$. We note that any two pairs of elements uniquely determine a triple in \mathcal{F} . So $m(\mathcal{F}) = 3$. Hence, $D = 1$.

Corollary 1

Fix a probability measure μ on the unit circle in the complex plane. If we color the circle with at least 14 equally sized μ -measurable color classes, there must be a rainbow triple of the form (x, y, xy)

- We note that the measure space is finite since $\mu(X) = 1$.
- $\mathcal{F} = \{(x, y, xy) \in X^3 \mid x, y \in X\}$. We note that any two pairs of elements uniquely determine a triple in \mathcal{F} . So $m(\mathcal{F}) = 3$. Hence, $D = 1$.
- We apply ICT with $D = 1$, $M = 3$, and $C \leq \frac{2D}{3M} = \frac{2}{27}$. We satisfy the color class because each color classes has μ -measure $1/14 < 2/27$.

Corollary 2

If we split any square into 104 equally sized Lebesgue measurable color classes, then it must contain three points x, y, z such that $|x - y| = |z - y| = |x - z|$ with the points being distinct colors.

Definition

Let X have a coloring, and $x \in X^k$. Then x is **w-subrainbow** if x has no w components of the same color.

$(x_1, x_2, x_3, x_4, x_5)$ is 3-subrainbow, but $(x_1, x_2, x_3, x_4, x_5)$ is not.

Definition

Let X have a coloring, and $x \in X^k$. Then x is **w-subrainbow** if x has no w components of the same color.

$(x_1, x_2, x_3, x_4, x_5)$ is 3-subrainbow, but $(x_1, x_2, x_3, x_4, x_5)$ is not.

Note: earlier cases are when $w = 2$, since every tuple has no 2 elements of the same color.

Definition

Let X have a coloring, and $x \in X^k$. Then x is **w-subrainbow** if x has no w components of the same color.

$(x_1, x_2, x_3, x_4, x_5)$ is 3-subrainbow, but $(x_1, x_2, x_3, x_4, x_5)$ is not.

Note: earlier cases are when $w = 2$, since every tuple has no 2 elements of the same color.

Definition

Let (X, μ) be a finite measure space, $\mathcal{F} \subset X^k$, and fix a $t \in [2..k-1]$. For any subset $S \subset [1..k]$, such that $|S| = t$, so write $S = \{s_1, s_2, \dots, s_t\}$. Define **t-bound** of \mathcal{F} be M_t , so $M_t = \sum_S M(S)$, where $M(S) = \sup_{(y_1, \dots, y_k) \in \mathcal{F}} |\{x \in \mathcal{F} : x_{s_j} = y_{s_j}\}|$

$$x = (x_1, x_2, x_3, \dots, x_t, \dots, x_k)$$

$$y = (y_1, y_2, y_3, \dots, y_t, \dots, y_k)$$

Similarly,

$$m = \sum_S m(S)$$

where

$$m(S) = \int \dots \int |\{y \in \mathcal{F} : x_{s_i} = y_{s_i}, i \in [1..t]\}| ds_1 \dots ds_t$$

Theorem(Generalized)

Let (X, μ) be a **finite measure space**, and define $n = \mu(X)$. Let $\mathcal{F} \subset X^k$ be measurable, and let the **t-bound** be M , and suppose $m(\mathcal{F}) \geq D \binom{k}{t} n^t$ and $2 \leq w \leq t < k$. Then for any **tractable coloring** of X , if $\mu(A) < Cn$ for all color classes A . Then \mathcal{F} must contain a w-subrainbow element, as long as $C^{w-1} < \frac{2^{w-1} D}{3^w M \binom{t}{w}}$

Proof.

We use the same technique as before



Corollary

Let G be an abelian group, $2, 3 \nmid |G|$, with some coloring. Then:

- 1 If no color class has size $\geq \frac{1}{135}|G|$, there must be a **rainbow** quintuple of the form $(x, y, z, x + y + z, x + 2y + 3z)$

Corollary

Let G be an abelian group, $2, 3 \nmid |G|$, with some coloring. Then:

- ① If no color class has size $\geq \frac{1}{135}|G|$, there must be a **rainbow** quintuple of the form $(x, y, z, x + y + z, x + 2y + 3z)$
- ② ...

For 1:

- Applying Generalized Theorem, here $X = G$, $k = 5$, and $\mathcal{F} = \{(x, y, z, x + y + z, x + 2y + 3z) \in G^5 \mid x, y, z \in G\}$.

Corollary

Let G be an abelian group, $2, 3 \nmid |G|$, with some coloring. Then:

- ① If no color class has size $\geq \frac{1}{135}|G|$, there must be a **rainbow** quintuple of the form $(x, y, z, x + y + z, x + 2y + 3z)$
- ② ...

For 1:

- Applying Generalized Theorem, here $X = G$, $k = 5$, and $\mathcal{F} = \{(x, y, z, x + y + z, x + 2y + 3z) \in G^5 \mid x, y, z \in G\}$.
- so $M_t = M_3 = \binom{5}{3} = 10$, $|\mathcal{F}| = |G|^3$

Corollary

Let G be an abelian group, $2, 3 \nmid |G|$, with some coloring. Then:

- 1 If no color class has size $\geq \frac{1}{135}|G|$, there must be a **rainbow** quintuple of the form $(x, y, z, x + y + z, x + 2y + 3z)$
- 2 ...

For 1:

- Applying Generalized Theorem, here $X = G$, $k = 5$, and $\mathcal{F} = \{(x, y, z, x + y + z, x + 2y + 3z) \in G^5 \mid x, y, z \in G\}$.
- so $M_t = M_3 = \binom{5}{3} = 10$, $|\mathcal{F}| = |G|^3$
- thus, $D = 1$, $w = 2$, and $C^{w-1} < \frac{2^{w-1}D}{3^w M\left(\frac{t}{w}\right)} = \frac{2 \cdot 1}{3^2 \cdot 10 \binom{3}{2}} = \frac{1}{135}$

Corollary

Let G be an abelian group, $2, 3 \nmid |G|$, with some coloring. Then:

- 1 ...
- 2 If no color class has size $\sqrt{\frac{2}{135}}|G|$, there must be a **3-subrainbow** quintuple of the form $(x, y, z, x + y + z, x + 2y + 3z)$

Corollary

Let G be an abelian group, $2, 3 \nmid |G|$, with some coloring. Then:

- 1 ...
- 2 If no color class has size $\sqrt{\frac{2}{135}}|G|$, there must be a **3-subrainbow** quintuple of the form $(x, y, z, x + y + z, x + 2y + 3z)$

For 2:

- Applying Generalized Theorem, here $X = G$, $k = 5$, and $\mathcal{F} = \{(x, y, z, x + y + z, x + 2y + 3z) \in G^5 \mid x, y, z \in G\}$.

Corollary

Let G be an abelian group, $2, 3 \nmid |G|$, with some coloring. Then:

- 1 ...
- 2 If no color class has size $\sqrt{\frac{2}{135}}|G|$, there must be a **3-subrainbow** quintuple of the form $(x, y, z, x + y + z, x + 2y + 3z)$

For 2:

- Applying Generalized Theorem, here $X = G$, $k = 5$, and $\mathcal{F} = \{(x, y, z, x + y + z, x + 2y + 3z) \in G^5 \mid x, y, z \in G\}$.
- so $M_t = M_3 = \binom{5}{3} = 10$, $|\mathcal{F}| = |G|^3$

Corollary






Let G be an abelian group, $2, 3 \nmid |G|$, with some coloring. Then:

- 1 ...
- 2 If no color class has size $\sqrt{\frac{2}{135}}|G|$, there must be a **3-subrainbow** quintuple of the form $(x, y, z, x + y + z, x + 2y + 3z)$





For 2:

- Applying Generalized Theorem, here $X = G$, $k = 5$, and $\mathcal{F} = \{(x, y, z, x + y + z, x + 2y + 3z) \in G^5 \mid x, y, z \in G\}$.
- so $M_t = M_3 = \binom{5}{3} = 10$, $|\mathcal{F}| = |G|^3$
- thus, $D = 1$, $w = 3$, and $C^{w-1} < \frac{2^{w-1}D}{3^w M \binom{t}{w}} = \frac{2^2 \cdot 1}{3^3 \cdot 10 \binom{3}{3}} = \frac{2}{135}$

Reference

-  M. Axenovich and D. Fon-Der-Flaass, *On rainbow arithmetic progressions*, Elec. J. Comb., 11:R1, (2004).
-  L. Babai, *An anti-Ramsey theorem*, Graphs Combin. 1 (1985), 23–28.
-  B. Green and T. Sanders, *Monochromatic sums and products*. Discrete Analysis, (2016) pages 1–43, 2016:5.
-  N. Hindman, *Partitions and sums and products of integers*. Trans. Amer. Math. Soc., 247:227–245, (1979).
-  N. Hindman, I. Leader, and D. Strauss, *Open problems in partition regularity*. Probably. Comput., 12(5-6):571–583, (2003). Special issue on Ramsey theory.

Reference

-  V. Jungić, J. Licht, M. Mahdian, J. Nešetřil, and R. Radoičić, *Rainbow Arithmetic Progressions and Anti-Ramsey Results*. *Combinatorics, Probability and Computing*, 12:599–620, (2003).
-  H. Lefmann and V. Rödl, *On canonical Ramsey numbers for complete graphs versus paths*, *J. Combin. Theory Ser. B*, 58:1–13, (1993).
-  S. Senger, *Rainbow triangles*. arXiv:1702.03043 (2017).
-  M. Young, *Rainbow arithmetic progressions in finite abelian groups*. arXiv:1603.08153 (2016).

Any Questions ??

Thank You 😊