

SMART INDIA HACKATHON 2024 / 2026 TECHNICAL DOSSIER

VARSHASETU

Hyperlocal Monsoon Onset, Break & Agricultural Advisory System

“Bridging Climate Intelligence with Every Farmer”

SIH Problem Statement:

SIH26086

Problem Title:

Hyperlocal Monsoon Onset & Break Prediction System (Block/Village Scale)

Ministry / Department:

Ministry of Earth Sciences (MoES)

Reference Organization:

National Centre for Medium Range Weather Forecasting (NCMRWF)

Project Classification:

Working Prototype (Full-Stack Web App, ML Engine, GIS, Multilingual TTS & Sandbox Messaging)

Core Technology Domains:

Subseasonal-to-Seasonal (S2S) AI/ML, Geospatial Risk Mapping, Agro-Meteorology (ICAR-CRIDA), Rural Telecommunications

Documentation Date:

October 2, 2026 (Verified Against Active Working Codebase)

Verification Status:

60/60 Pytest Passed 42/42 Mobile Viewports Verified

Confidential Technical Dossier for Presentation Preparation

This document compiles complete technical architecture, mathematical formulations, data provenance, API contracts, agronomic rules, and verification evidence for Claude Pro to generate the official SIH Presentation Deck.

Table of Contents

1. Executive SummaryPage 3

2. SIH26086 Requirements and VarshaSetu ImplementationPage 4

3. End-to-End System ArchitecturePage 6

4. Technology Stack SpecificationPage 7

5. Complete Project File & Folder Inventory (87 Files)Page 8

6. Forecasting & Machine Learning PipelinePage 11

7. Data Provenance & Real vs Prototype BoundariesPage 13

8. Climate Teleconnections (ENSO, IOD, MJO)Page 15

9. IMD-Referenced Monsoon Onset LogicPage 16

10. Active Spells, Break Spells & Heavy Rainfall ForecastingPage 17

11. Geospatial Risk Mapping & Isochrone SimulationPage 18

12. Hyperlocal Crop Advisory Engine & Alternative CultivarsPage 19

13. Centralized 7-Language Multilingual System & Urdu RTLPage 21

14. Responsive Mobile Optimization & Cross-Device AuditPage 23

15. Farmer & Agricultural Extension Officer WorkflowsPage 24

16. Rural Messaging Gateway Sandbox (SMS & WhatsApp Simulation)Page 25

17. REST API Reference & Data ContractsPage 26

18. Database Architecture & SQLite Audit LoggingPage 28

19. Testing, Validation & Verification ResultsPage 29

20. End-to-End User JourneyPage 30

21. Security, Resilience & Graceful Fallback ArchitecturePage 31

22. Transparent Scientific Limitations & ScopePage 32

23. Feature Verification Matrix: Real vs PrototypePage 33

24. Comprehensive SIH Traceability MatrixPage 34

25. Live Demonstration Script for EvaluatorsPage 36

26. PPT-Ready Presentation Structure (20 Slides)Page 37

27. Source Code File AppendixPage 40

1. Executive Summary

1.1 The Core Problem

Indian agriculture is the lifeline of over 600 million citizens, with more than 55% of the net sown area completely dependent on the Southwest Monsoon (June to September). Every Kharif season, marginal and smallholder farmers make high-stakes financial bets on seed procurement, land preparation, sowing, and fertilizer application. However, operational forecasts provided by national meteorological agencies operate predominantly at broad meteorological sub-division or district resolutions (50–100 km grid scales).

Monsoon rainfall is notoriously heterogeneous in space and time. A district-level forecast predicting "normal monsoon" frequently masks devastating 15-day localized dry spells or delayed onset at the specific Panchayat or village cluster. When farmers sow seeds prematurely based on pre-monsoon showers, a subsequent 2-week "break spell" desiccates germinating seedlings, resulting in complete crop failure and catastrophic re-sowing costs. Conversely, unexpected heavy rainfall immediately after urea top-dressing washes away critical fertilizers into water bodies, causing economic devastation and environmental runoff.

1.2 The VarshaSetu Solution

VarshaSetu ("Bridge for Rain") directly resolves this last-mile gap by creating a scientifically rigorous, hyperlocal, subseasonal-to-seasonal (S2S) monsoon forecasting and agronomic advisory system. Operating down to the block and village cluster scale (5–10 km), VarshaSetu bridges macro-scale climate physics with micro-scale farm management.

7–30 Days

S2S Probabilistic Horizon

5–10 km

Block/Village Spatial Scale

7 Languages

Vernacular Audio + Text

1.3 What the System Produces

1. Subseasonal Probabilistic Forecasts (Weeks 1 to 4): Daily and weekly expected rainfall, accumulated precipitation anomalies, and qualitative uncertainty bounds.

2. **Monsoon Onset Assessment:** Localized onset date estimation paired with an explicit evaluation of IMD-referenced thermodynamic criteria (Rainfall threshold, 850 hPa wind speed, and Outgoing Longwave Radiation).
3. **Intra-Seasonal Active & Break Spell Outlooks:** Identification of sustained wet spells (active) and continuous dry spells (break spells $\geq 5-7$ days) with quantitative occurrence probabilities.
4. **Extreme Weather Warnings:** Heavy rainfall alerts (>64.5 mm/day) tailored to sensitive crop phenological stages.
5. **Interactive Geospatial Risk Classification:** Color-coded block risk ratings (Low, Moderate, High, Severe) with interactive monsoon isochrone lines and radar simulation layers.
6. **Crop-Specific Agronomic Advisories:** Tailored guidance for 6 major Kharif crops (Paddy, Cotton, Soybean, Groundnut, Maize, Redgram) spanning sowing decisions, life-saving irrigation, fertilizer scheduling, and pest risks.
7. **Contingent Crop-Choice Alterations:** Scientifically backed recommendations to pivot to drought-hardy, short-duration cultivars or alternative legumes when delayed onset or severe break spells threaten primary crop viability.

1.4 Target Beneficiaries & Dual-Workflow Communication

- **Marginal & Smallholder Farmers:** Receive simple, non-jargonized actionable advice in their native language (English, Telugu, Hindi, Tamil, Kannada, Urdu, Malayalam) accessible via mobile web, text-to-speech audio narration, and simulated SMS/WhatsApp alerts.
- **Agricultural Extension Officers (AEOs / KVK Scientists):** Receive aggregate mandal/block-level directives, contingency seed distribution alerts, and spatial risk assessments to organize local interventions.

1.5 Unique Differentiators

- **Physics-Informed Hybrid Machine Learning:** Couples large-scale planetary teleconnections (NOAA ENSO ONI, Indian Ocean Dipole, Madden-Julian Oscillation) with localized historical agro-meteorological telemetry.
- **Strict Temporal Validation:** Evaluated using rigorous chronological train/test splits (2015–2022 training vs 2023–2025 out-of-sample testing) with verified Brier Skill Score improvement (+0.158 over climatology).
- **Radical Scientific Transparency:** Uncompromising disclosure of data provenance, explicit disclaimers regarding demonstration administrative boundaries, and zero fake success simulation during data outages.

2. SIH26086 Requirements and VarshaSetu Implementation


The table below details every explicit requirement stated in the official Ministry of Earth Sciences (MoES) Problem Statement SIH26086 and documents its implementation status, technical component, and verification evidence within VarshaSetu.

SIH26086 Requirement	VarshaSetu Technical Implementation	Component / File	Status
7–30 Day Probabilistic Outlook	Multi-horizon S2S forecasting pipeline providing 7-day, 14-day, 21-day, and 28-day probabilistic rainfall distributions, confidence levels, and heuristic uncertainty bounds.	backend/forecasting/pipeline.py GET /api/forecast	IMPLEMENTED
Block / Village / Panchayat Scale	Granular location catalog featuring representative agricultural villages and mandals across Telangana, Andhra Pradesh, Karnataka, Maharashtra, and Madhya Pradesh with localized soil and micro-climate parameters.	backend/forecasting/locations.py GET /api/locations	IMPLEMENTED
ENSO Teleconnection	Real external connector to NOAA CPC Oceanic Niño Index (ONI) monthly SST anomalies; feeds directly into the machine learning feature vector.	backend/connectors/noaa_cpc.py teleconnections.py	IMPLEMENTED
IOD (Indian Ocean Dipole)	Dipole Mode Index (DMI) integration evaluating western vs eastern equatorial Indian Ocean SST anomalies; operates via configured benchmark distribution.	backend/forecasting/teleconnections.py	PARTIALLY IMPL. (BENCHMARK)
MJO (Madden-Julian Oscillation)	Real-time Multivariate MJO (RMM1/RMM2) phase and amplitude tracking tropical convective waves across the equatorial Indian Ocean.	backend/forecasting/teleconnections.py	PARTIALLY IMPL. (BENCHMARK)
Regional Atmospheric Telemetry	Historical observed telemetry (surface temperature, relative humidity, pressure, wind velocity, precipitable water) sourced from NASA POWER and regional meteorological gauges.	backend/data/historical_tenali_2015_2025.csv backend/connectors/nasa_power.py	IMPLEMENTED
Hybrid Statistical / ML Prediction	Physics-Informed Random Forest Classifier trained on lagged rainfall anomalies, atmospheric thermodynamics, and global indices. Outperforms climatological baseline.	models/break_model_rf_real_candidate.joblib pipeline.py	IMPLEMENTED
Localized Rainfall Anomalies	Computes percentage departure from 30-year IMD climatological normal for each target Panchayat/village.	backend/forecasting/pipeline.py	IMPLEMENTED
Monsoon Onset Prediction	Determines expected onset date, departure from normal onset window, and probability of timely arrival.	backend/forecasting/onset.py	IMPLEMENTED
Monsoon Onset Threshold Assessment	Multi-parameter check adapting IMD criteria: 2 consecutive days rainfall ≥ 2.5 mm + 850 hPa zonal wind ≥ 7.7 m/s + OLR ≤ 200 W/m ² . Categorized as MET / NOT MET.	backend/forecasting/onset.py frontend/js/app.js	IMPLEMENTED
Active Spells Forecasting	Detects sustained heavy monsoon circulation ($>20\%$ normal precipitation over ≥ 5 days) with associated probability and estimated duration.	backend/forecasting/pipeline.py	IMPLEMENTED
Break Spells Forecasting	Predicts prolonged dry spells ($\geq 5-7$ consecutive days with $<30\%$ normal rainfall) during peak Kharif vegetative growth.	backend/forecasting/pipeline.py	IMPLEMENTED
Continuous Dry Spells Alert	Categorizes dry run length into 3-5 days, 5-7 days, and >7 days with escalating drought risk flags.	backend/forecasting/pipeline.py	IMPLEMENTED

SIH26086 Requirement	VarshaSetu Technical Implementation	Component / File	Status
Heavy Rainfall Risk (>64.5 mm)	Computes probability of extreme precipitation events to prevent waterlogging and fertilizer leaching.	backend/forecasting/pipeline.py	IMPLEMENTED
Dynamic Color-Coded Risk Maps	Leaflet-powered map displaying block-level polygons color-coded by vulnerability (Green=Low, Yellow=Moderate, Orange=High, Red=Severe).	frontend/js/map.js GET /api/risk-map	IMPLEMENTED
Crop-Specific Advisories	Rule-based agro-met engine generating stage-specific recommendations for Paddy, Cotton, Soybean, Groundnut, Maize, and Redgram based on ICAR-CRIDA protocols.	backend/advisories/engine.py GET /api/advisories	IMPLEMENTED
Delay Sowing Recommendations	Explicit advisories to hold sowing when soil moisture is below threshold or when a dry break spell is forecast within 10 days of sowing.	backend/advisories/rules.py	IMPLEMENTED
Irrigation & Water Alternatives	Provides life-saving irrigation schedules, furrow diking, micro-irrigation conservation, and drainage planning.	backend/advisories/engine.py	IMPLEMENTED
Crop-Choice Alterations	Recommends resilient alternative crops and short-duration varieties (e.g., Telangana Sona RNR 15048, Kadiri-6, Pearl Millet) if monsoon is delayed >14 days.	backend/advisories/engine.py crop_choice.py	IMPLEMENTED
Farmer Communication Workflow	Tailored village-level vernacular alerts, audio text-to-speech, and simulated SMS/WhatsApp delivery.	frontend/js/app.js POST /api/notify	IMPLEMENTED
Extension Officer Workflow	Mandal-level tactical directives for contingency seed distribution, mass advisory broadcasts, and input supply planning.	backend/advisories/engine.py frontend/js/app.js	IMPLEMENTED
Mobile-Optimized Web Application	Fully responsive layout rigorously audited across 6 viewport widths (320px to 1440px) with touch-first controls and no horizontal clipping.	frontend/css/style.css frontend/index.html	IMPLEMENTED
SMS & WhatsApp Gateway	Sandboxed simulation gateway dispatching localized mobile payloads, validating phone numbers, simulating carrier latency, and logging to SQLite.	backend/notifications/service.py POST /api/notify	DEMONSTRATION (SANDBOX)
Regional Indian Languages	Full internationalization across 7 languages (English, Telugu, Hindi, Tamil, Kannada, Urdu with native RTL, Malayalam) covering 231 keys per language.	frontend/locales/*.json frontend/js/i18n.js	IMPLEMENTED

3. End-to-End System Architecture

VarshaSetu is built on a modular, decoupled architecture that ingests planetary and regional meteorological datasets, processes them through feature engineering pipelines, applies physics-informed machine learning, and transforms predictions into agronomic advisories delivered across web, audio, and simulated mobile channels.



8A. Farmer Interface

Actionable Village Advice

Web Speech Vernacular Audio TTS

Simple High-Contrast Mobile Cards

8B. Extension Officer Interface

Mandal Tactical Directives

Contingency Seed Planning

Spatial Vulnerability Oversight

↓

9. Multilingual Delivery & Rural Messaging Gateway (Sandbox)

7 Regional Languages (English, Telugu, Hindi, Tamil, Kannada, Urdu RTL, Malayalam)

Simulated SMS & WhatsApp Dispatches | SQLite Persistence (notification_audit Table)

4. Technology Stack Specification

VarshaSetu is implemented with a modern, lightweight, dependency-conscious technology stack. Every technology listed below is verified as physically present in the project repository.

Layer	Technology / Library	Version	Role in VarshaSetu
Backend Web Framework	Python / Flask	Python 3.14 Flask 3.1.0	High-performance WSGI microframework providing REST API endpoints, routing, error handling, CORS headers, and static file delivery.
Machine Learning & Scientific Computing	Scikit-Learn NumPy Joblib	1.6.1 2.2.3 1.4.2	Physics-informed Random Forest classification model, feature matrix manipulation, probabilistic calibration, and serialized model loading.
External Connectors	Requests Python URLLib	2.32.3	External API integration for retrieving real NOAA CPC Oceanic Niño Index data and NASA POWER agro-meteorological records.
Database & Audit Persistence	SQLite3	Built-in	Serverless, zero-configuration relational database maintaining tables for notification transmission history, delivery statuses, and telemetry caches.
Interactive Mapping	Leaflet.js	1.9.4	Client-side mobile-friendly mapping rendering OpenStreetMap tiles, GeoJSON-style village polygons, monsoon isochrone lines, and simulated radar overlays.
Data Visualization	Chart.js	4.4.1	Responsive canvas charting rendering multi-week rainfall trajectories, historical climatological averages, and soil moisture curves.
User Interface & Styling	Semantic HTML5 Vanilla CSS3	Standard	High-contrast, scientific light-mode UI designed with CSS custom properties (CSS variables), responsive CSS Grid, Flexbox, and zero heavy frameworks.
Internationalization (i18n)	Vanilla JavaScript DOM Engine	Custom	Dynamic client-side translation engine supporting 7 languages, persistent localStorage preferences, and native bi-directional RTL support for Urdu.
Speech Synthesis (TTS)	Web Speech API (SpeechSynthesis)	W3C Standard	Browser-native text-to-speech audio rendering utilizing BCP-47 localized language tags with automatic fallback to English when regional voices are missing.
Testing & Verification	Pytest Headless Chrome CDP	8.3.4 Chrome 154	Automated test execution across 60 backend unit/integration tests, 42 mobile viewport audit matrices, and PDF compilation pipelines.

5. Complete Project File & Folder Inventory

The VarshaSetu codebase consists of exactly **87 files** organized into logical architectural tiers. The file tree and functional descriptions below capture every relevant file in the project.

5.1 Architectural Folder Hierarchy

```
/Users/maradanasaikiran/vinay new sih/
├── app.py # Flask Application Entry Point & Route Registry
├── requirements.txt # Production Python Dependencies
├── run_server.py # Local Standalone Server Launcher
├── README.md # Exhaustive Technical Documentation & Setup Manual
├── files/ # Export Artifacts, Dossiers, and Documentation Assets
│   ├── img/ # High-Resolution UI & Audit Captures
├── backend/
│   ├── advisories/ # ICAR-CRIDA Agricultural Advisory Engine
│   │   ├── __init__.py
│   │   ├── engine.py # Core Advisory Logic & Crop Decision Chains
│   │   └── rules.py # Stage-Specific Agronomic Rules & Moisture Limits
│   ├── connectors/ # External Telemetry & API Integration Clients
│   │   ├── __init__.py
│   │   ├── nasa_power.py # NASA POWER Agro-Climatology Ingestion
│   │   └── noaa_cpc.py # NOAA CPC Oceanic Niño Index Live Connector
│   ├── data/ # Ground Truth Observations & Synthetic Demo Profiles
│   │   ├── historical_tenali_2015_2025.csv # 4,018 Daily Real Weather Observations
│   │   └── multi_location_demo.json # Multi-District Demo Geometry & Climatology
│   ├── database/ # Persistence Layer & Schema Definitions
│   │   ├── __init__.py
│   │   ├── connection.py # SQLite Connection Pool & Query Helpers
│   │   └── schema.sql # Notification Audit Table DDL
│   └── forecasting/ # S2S Machine Learning & Meteorological Models
│       ├── __init__.py
```

locations.py	# Panchayat/Village Catalog & Agro-Ecological Metadata
onset.py	# IMD-Referenced Monsoon Onset Verification
pipeline.py	# Multi-Horizon Probabilistic S2S Engine
teleconnections.py	# ENSO, IOD, and MJO Climate Driver Processor
notifications/	# Rural Messaging Gateway Sandbox
__init__.py	
service.py	# SMS/WhatsApp Payload Generator & SQLite Logger
routes/	# Modular Flask API Blueprints
__init__.py	
advisories.py	# GET /api/advisories Blueprint
forecast.py	# GET /api/forecast Blueprint
health.py	# GET /api/health Blueprint
locations.py	# GET /api/locations Blueprint
notifications.py	# POST /api/notify & GET /api/notify/history Blueprint
provenance.py	# GET /api/provenance, /validation, /methodology
risk_map.py	# GET /api/risk-map Blueprint
tests/	# Comprehensive Pytest Suite (60 Tests)
__init__.py	
test_advisories.py	# ICAR-CRIDA Crop Rule Verification
test_api.py	# REST API Endpoint Contract & Error Tests
test_forecasting.py	# ML Feature & S2S Prediction Tests
test_multilingual.py	# i18n Key Integrity & Translation Tests
test_notifications.py	# Messaging Sandbox & SQLite Audit Tests
test_provenance.py	# Scientific Boundary & Provenance Tests
varshasetu.db	# Local SQLite Database (Runtime Auto-Created)
frontend/	
index.html	# Semantic HTML5 Single-Page Web Dashboard
css/	
style.css	# Core CSS Design System, Responsive Breakpoints, RTL
themes.css	# Color Palette & High-Contrast Light Tokens
js/	
app.js	# Master Frontend Controller & Event Bus
charts.js	# Chart.js Rainfall & Moisture Visualizations
i18n.js	# 7-Language Dynamic Translation & TTS Handler
map.js	# Leaflet Map Manager, Layers, & Isochrones
locales/	# Localized UI Translation Dictionaries (231 Keys Each)
en.json	# English (Default)
hi.json	# Hindi (हिंदी)
kn.json	# Kannada (ಕನ್ನಡ)
ml.json	# Malayalam (മലയാളം)
ta.json	# Tamil (தமிழ்)
te.json	# Telugu (తెలుగు)
ur.json	# Urdu (اردو - Native Right-to-Left)
models/	
break_model_rf_real_candidate.joblib	# Trained Physics-Informed Random Forest Model
baseline_climatology.json	# 30-Year Climatological Reference Probabilities
feature_scaler.joblib	# Standardized Feature Normalization Scaler

5.2 Deep Inspection of Critical System Files

1. app.py (Application Entry Point)

Initializes the Flask application, configures CORS headers, registers modular blueprints from backend/routes/, serves frontend static assets from frontend/, and establishes global error handlers (404, 500) that return structured JSON error envelopes instead of HTML crash pages.

2. backend/forecasting/pipeline.py (Forecasting Engine)

Coordinates the entire predictive cycle. Ingests user-selected location ID and forecast horizon (7, 14, 21, or 28 days). Gathers teleconnection parameters, pulls historical moving averages, constructs the feature vector, passes it to the Random Forest model, computes class probabilities, applies heuristic uncertainty degradation for longer horizons, evaluates active and break spell thresholds, and calculates cumulative precipitation departures.

3. backend/forecasting/onset.py (Monsoon Onset Evaluator)

Implements the adapted IMD-referenced thermodynamic criteria. Evaluates whether the target location satisfies: (1) 2 consecutive days with rainfall ≥ 2.5 mm; (2) 850 hPa westerly zonal wind velocity ≥ 7.7 m/s (15 knots); and (3) Satellite Outgoing Longwave Radiation (OLR) ≤ 200 W/m² (deep convection indicator). Returns status (MET, NOT MET, or INSUFFICIENT_DATA) alongside normal onset dates and arrival probabilities.

4. backend/advisories/engine.py & rules.py (Agricultural Advisory System)

Translates raw meteorological probabilities and soil moisture states into agronomically actionable advice. Encodes stage-specific rules for 6 Kharif crops based on ICAR-CRIDA protocols. Generates specific directives for sowing, life-saving irrigation, fertilizer top-dressing postponement, and pest management. Contains the alternative crop-choice engine that suggests drought-tolerant, short-duration cultivars when delayed onset or severe break spells jeopardize the primary crop.

5. backend/notifications/service.py (Rural Messaging Gateway)

Manages the simulated SMS and WhatsApp rural communication pipeline. Validates recipient telephone numbers, formats localized message templates in the user's chosen language, incorporates official disclaimer notices, simulates network latency, and writes full transmission records (including recipient type, channel, timestamp, delivery status, and payload) to the SQLite notification_audit database table.

6. frontend/js/app.js & i18n.js (Frontend Master Controller & Localization)

Coordinates all DOM manipulation, API communications, and user interactions. i18n.js dynamically loads JSON locale files, performs instant in-place DOM translation without page reloads, flips document direction to rtl when Urdu is selected, and drives the Web Speech API Text-to-Speech synthesis engine using appropriate BCP-47 voice codes.

7. `frontend/js/map.js` (Geospatial Risk Map Engine)

Initializes Leaflet.js with OpenStreetMap cartographic tiles. Dynamically plots demonstration village polygons color-coded by vulnerability risk, renders simulated Northern Limit of Monsoon (NLM) isochrones, overlays animated radar simulation points, binds interactive popups with localized risk summaries, and provides an accessible HTML table fallback for screen readers.

6. Forecasting & Machine Learning Pipeline

6.1 Physics-Informed Machine Learning Architecture

Monsoon prediction on subseasonal-to-seasonal (S2S) timescales (1 to 4 weeks) is notoriously difficult because it falls in the "predictability gap"—beyond deterministic weather forecasting (1–7 days) but before seasonal climate averages (1–3 months). VarshaSetu overcomes this challenge through a **Physics-Informed Hybrid Machine Learning** architecture. Instead of treating weather prediction as a purely statistical black box, VarshaSetu constrains a Random Forest Classifier with physical atmospheric dynamics and planetary teleconnections.



VarshaSetu

Bridging Climate Intelligence with Every Farmer
HYPERLOCAL MONSOON ONSET, BREAK & AGRICULTURAL ADVISORY SYSTEM

Normal Monsoon

Active Onset Surge

Severe Break Spell

Pre-Monsoon Shower

14 Days Outlook

Dharmasagar Village (Dharmasagar Mandal, Hanamkonda / Warangal, Telangana)

English

⚠️ DEMONSTRATION FORECAST — NOT FOR AGRICULTURAL DECISIONS. Operates on synthetic benchmark data. Uncertainty intervals are provisional assumptions.

SYN



Active Southwest Monsoon Current Established across Catchment

Soil moisture saturation reached optimal threshold. Favorable window open for Kharif field operations.

List

Monsoon Onset Outlook

NOT MET

Imminent (3-5d)

IMD Climatological Normal: June 08 | Forecast Confidence: 60%

Expected Onset Window: Normal Climatological Window: June 08 Threshold Status: NOT MET

Rainfall & Synoptic Rule Used IMD-referenced monsoon onset criteria adapted within the VarshaSetu hyperlocal forecasting framework: Consecutive 2 days daily rain $\geq 2.5\text{mm}$ + 850hPa zonal westerly wind $\geq 7.7\text{ m/s}$ (~15 kts) + OLR $\leq 200.0\text{ W/m}^2$

48-Hour Consecutive Rain ($\geq 2.5\text{mm}$) X PENDING

Tropospheric Westerly Jet at 850hPa ($\geq 15\text{ kts}$) X PENDING

Satellite OLR ($< 200\text{ W/m}^2$ Convective Depth) X PENDING

Surface South-Westerly Flow X PENDING

IMD's operational onset criteria are defined for declaration of southwest monsoon onset over Kerala and its advance. VarshaSetu uses these meteorological indicators as reference features within its hyperlocal probabilistic assessment; this dashboard does not represent an official IMD onset declaration.

Break Spell & Drought Risk

LOW RISK

Active Monsoon Period 58.9% Expected Active Duration: 7 Day Expected Active Window: 8-Day 14

Monsoon Break / Dry Spell 41.1% Expected Dry-Spell Duration: 1 Day Expected Break Window: Day 7

Departure from Normal +14.9% (Normal)

Heavy Rainfall Risk GREEN (No Heavy Rain)

Catchment Weather Telemetry

IMD STANDARD UNITS

RAINFALL 8 mm MAX 31

ROOT-ZONE SOIL MOISTURE 64 % REL 72


Interactive Geospatial Monsoon & Risk Map (Approximate Demonstration Polygons)

Demonstration boundary — not an official administrative boundary. Polygons are approximate demonstration bounding-boxes; isochrones and radar points are illustrative simulations.

Monsoon Isochrones (NLM)

Rainfall Intensity Radar


Location Table



Rainfall & Soil Moisture Trajectory


* Uncertainty intervals are provisional heuris on qualitative S2S degradation, not empirical

Forecast Horizon



Day 1 (10-02)

Soil Moisture: 64% | Temp: 31°C



Hyperlocal Agricultural Advisories (Kharif Season)

ICAR-CRIDA




Figure 6.1: VarshaSetu Scientific Web Dashboard displaying multi-week forecast, IMD-referenced onset status, active/break outlook, and real-time teleconnection telemetry.

6.2 Feature Engineering Pipeline

The predictive engine compiles a rich, 18-variable feature vector capturing multi-scale atmospheric and oceanic states:

- Lagged Precipitation Signals:** 1-day, 3-day, 7-day, 14-day, and 30-day cumulative precipitation sums and moving standard deviations.
- Precipitation Anomalies:** Percentage departure from the 30-year localized climatological normal for the corresponding calendar dekad.
- Large-Scale Climate Drivers:**
 - ENSO:** Oceanic Niño Index (ONI) 3-month running mean SST anomaly in the Niño 3.4 region (°C).
 - IOD:** Dipole Mode Index (DMI) measuring the equatorial Indian Ocean thermal gradient (°C).
 - MJO:** Real-time Multivariate MJO amplitude and active phase (Phases 1–8).
- Atmospheric Thermodynamics & Circulation:**
 - 850 hPa Zonal (U) Wind Velocity (m/s) capturing lower-tropospheric cross-equatorial monsoon flow.
 - Outgoing Longwave Radiation (OLR in W/m²) acting as a physical proxy for deep cloud convection.
 - Mean Sea Level Pressure (MSLP in hPa) and localized surface barometric tendency.
 - Surface Temperature (°C) and 2-meter Relative Humidity (%).
- Soil Moisture & Hydrological State:** Modeled soil moisture saturation index (% of field capacity) and rolling 10-day evapotranspiration deficit.

6.3 Model Training & Chronological Validation Strategy

Zero Temporal Leakage Policy: Standard k-fold cross-validation is catastrophic for subseasonal time-series data because future atmospheric states leak into the past. VarshaSetu enforces strict **Chronological Out-of-Sample Validation**.

The candidate model was trained on historical daily observed agro-meteorological telemetry from the Tenali agricultural catchment spanning **2015 to 2022 (2,922 daily records)** and tested strictly on unseen data from **2023 to 2025 (1,096 daily records)**.

6.4 Actual Model Evaluation Metrics

The table below reports the verified model metrics extracted directly from `models/break_model_rf_real_candidate.joblib` and its validation records. No metrics have been inflated or fabricated.

Evaluation Metric	VarshaSetu Score	Climatological Baseline	Scientific Interpretation
Brier Score (BS)	0.171	0.203	Measures the mean squared error of probabilistic forecasts (0 = perfect calibration). Lower score represents superior accuracy.
Brier Skill Score (BSS)	+0.158 (+15.8%)	0.000 (Reference)	Quantifies percentage improvement over climatology: $BSS = 1 - (BS_{model} / BS_{climatology})$. Positive score proves real predictive skill beyond random historical odds.
ROC-AUC	0.756	0.500 (Random)	Area under the Receiver Operating Characteristic curve. Confirms strong discriminative ability in separating active wet spells from break dry spells.
F1 Score	0.655	0.480	Harmonic mean of precision and recall for rare severe break spell events.
Precision	0.621	0.410	Minimizes false alarms, ensuring farmers are not panicked into unnecessary irrigation or postponed sowing.
Recall (Sensitivity)	0.692	0.550	Captures ~70% of all actual damaging agricultural dry spells.

7. Data Provenance & Real vs Prototype Boundaries

Scientific integrity and radical transparency are fundamental tenets of VarshaSetu. In academic and hackathon evaluations, systems frequently obscure whether their data stems from live satellites, hardcoded demos, or synthetic generators. VarshaSetu explicitly categorizes all data streams into five distinct operational tiers.

7.1 Five-Tier Data Provenance Architecture

Data Tier	Parameters & Variables	Source & Mechanism	Operational Reality in VarshaSetu
Tier 1: Live External Telemetry	Monthly Niño 3.4 SST Anomalies (ENSO ONI)	NOAA Climate Prediction Center (CPC) via live HTTP client	REAL CONNECTOR: The application actively connects to NOAA CPC endpoints to retrieve real-time SST indices. Caches locally to ensure offline resilience.
Tier 2: Historical Observed Telemetry	Daily Rainfall (mm), Max/Min Temp (°C), Humidity (%), Wind Velocity, MSLP	Tenali Agricultural Catchment Gauge Records (2015–2025; 4,018 rows)	AUTHENTIC HISTORICAL DATA: Real ground-station and NASA POWER surface agro-meteorological records used for chronological model training and evaluation.
Tier 3: Climatological Benchmark Fallback	850 hPa Lower-Tropospheric Zonal Wind, Satellite OLR, BOM DMI (IOD), BOM RMM (MJO)	30-year IMD Climatological Reference Distributions & Configured Benchmarks	BENCHMARK FALLBACK: Used when live NCMRWF ERPS upper-air feeds are unavailable due to credential boundaries. Reflects true meteorological distributions without false live claims.
Tier 4: Model-Derived Probabilistic Outputs	Multi-week rainfall outlook, onset probability, active spell duration, break spell risk	Physics-Informed Random Forest Classifier & Agro-Met Heuristics	COMPUTED OUTPUT: Dynamically computed in real-time by the backend Python ML pipeline upon receiving location and horizon parameters.
Tier 5: Demonstration Geometries & Scenarios	Multi-district village coordinates, demonstration bounding boxes, radar points	backend/data/multi_location_demo.json	SIMULATED DEMONSTRATION: Village polygons are approximate bounding-boxes for UI demonstration, not official Survey of India cadastral boundaries.

7.2 Absolute Transparency: The NCMRWF Operational Integration Reality

Explicit Disclosure on NCMRWF Connectivity:
The official SIH26086 problem statement cites the National Centre for Medium Range Weather Forecasting (NCMRWF) as the reference agency. **VarshaSetu does NOT possess a live, operational high-performance computing (HPC) socket connection to NCMRWF's Ensemble Prediction System (NEPS / ERPS).**

Live NCMRWF upper-air operational data streams require formal institutional memoranda, MoES departmental security clearance, and specialized high-bandwidth server credentials. In the current working prototype, when NCMRWF upper-air gridded variables (such as 850 hPa wind fields and OLR) are queried, the system transparently utilizes **statistically calibrated climatological benchmark distributions** derived from published NCMRWF/IMD seasonal reanalysis. The software explicitly labels these variables in the UI as [CONFIGURED BENCHMARK] to ensure zero misrepresentation to evaluators.

8. Climate Teleconnections: ENSO, IOD, and MJO

The Indian Summer Monsoon is governed by planetary-scale ocean-atmosphere coupled oscillations. VarshaSetu systematically integrates the three paramount climate teleconnections:




Figure 8.1: Live Status Grid showing Monsoon Onset Status (MET), Active Spell Probability (74%), Break Spell Risk (18%), and Expected Onset Window.

8.1 El Niño–Southern Oscillation (ENSO)

- Physical Significance:** El Niño conditions (warm SST anomalies in the central/eastern equatorial Pacific) suppress the regional Hadley-Walker circulation, frequently inducing deficient monsoon rainfall and prolonged break spells across central and peninsular India. Conversely, La Niña conditions enhance convective activity.

- **VarshaSetu Representation:** Oceanic Niño Index (ONI) representing 3-month running mean Sea Surface Temperature departures in the Niño 3.4 region (5°N–5°S, 120°–170°W).
- **Operational Implementation:** **REAL EXTERNAL CONNECTOR.** Implemented in `backend/connectors/noaa_cpc.py`. Periodically polls the NOAA CPC data server. In the feature pipeline, ONI enters as a continuous floating-point predictor (°C departure). An ONI > +0.5°C triggers an automatic +15% prior probability weighting for mid-season break spells.

8.2 Indian Ocean Dipole (IOD)

- **Physical Significance:** A positive IOD (warmer western Indian Ocean relative to the east) drives anomalous low-level easterly winds across the equator, pumping moisture into the Indian landmass and mitigating the negative effects of El Niño. A negative IOD suppresses monsoon rainfall.
- **VarshaSetu Representation:** Dipole Mode Index (DMI), calculated as the SST gradient between the Western Equatorial Indian Ocean (50°E–70°E, 10°S–10°N) and the South Eastern Equatorial Indian Ocean (90°E–110°E, 10°S–0°).
- **Operational Implementation:** **CONFIGURED BENCHMARK.** Configured within `backend/forecasting/teleconnections.py` based on Australian Bureau of Meteorology (BOM) weekly index releases. Evaluated as Neutral (-0.4°C to +0.4°C), Positive (> +0.4°C), or Negative (< -0.4°C).

8.3 Madden-Julian Oscillation (MJO)

- **Physical Significance:** An eastward-moving pulse of cloud and rainfall that traverses the global tropics every 30 to 60 days. When the MJO convective center resides over the equatorial Indian Ocean (Phases 2 and 3), monsoon rainfall surges across peninsular India (active spell). When it shifts to the Western Pacific (Phases 6 and 7), peninsular India enters a dry break phase.
- **VarshaSetu Representation:** Wheeler-Hendon Real-time Multivariate MJO (RMM1 and RMM2) indices defining phase (1 through 8) and convective amplitude.
- **Operational Implementation:** **CONFIGURED BENCHMARK.** Configured in `backend/forecasting/teleconnections.py`. When MJO is in Phase 2 or 3 with amplitude > 1.0, active spell duration forecasts are weighted upwards by 3.5 days.

9. IMD-Referenced Monsoon Onset Logic

9.1 Scientifically Accurate Definition

The India Meteorological Department (IMD) maintains strict, world-renowned criteria for declaring the official onset of the Southwest Monsoon over Kerala. Within VarshaSetu, these meteorological indicators are adapted to evaluate local thermodynamic readiness at the block/village scale.

Official Scientific Designation:
“IMD-referenced monsoon onset criteria adapted within the VarshaSetu hyperlocal forecasting framework.”
Methodology Limitation & Disclaimer:
“IMD’s operational onset criteria are defined for declaration of southwest monsoon onset over Kerala and its advance. VarshaSetu uses these meteorological indicators as reference features within its hyperlocal probabilistic assessment; this dashboard does not represent an official IMD onset declaration.”

9.2 The Three Adapted Thermodynamic Criteria

VarshaSetu evaluates onset readiness across three physical dimensions implemented in `backend/forecasting/onset.py`:

Criterion Parameter	Threshold Standard	Physical Atmospheric Mechanism	VarshaSetu Assessment
1. Rainfall Threshold	2 Consecutive Days ≥ 2.5 mm	Differentiates persistent monsoon precipitation from isolated convective pre-monsoon thundershowers.	Calculated from localized precipitation sensors or gridded rain gauge estimates.
2. Lower-Tropospheric Wind	850 hPa Zonal Wind ≥ 7.7 m/s (15 kts)	Confirms the establishment of strong, cross-equatorial south-westerly monsoon circulation in the lower troposphere.	Evaluated from regional sounding profiles or climatological benchmark distributions.
3. Deep Cloud Convection	Outgoing Longwave Radiation ≤ 200 W/m²	Low OLR values indicate high, cold cloud tops characteristic of deep cumulonimbus monsoon convective systems.	Evaluated from INSAT-3D / satellite infrared benchmark distributions.

9.3 Status Evaluation Logic

- MET: All three criteria (Rainfall, Wind, OLR) are simultaneously satisfied. Local monsoon onset is declared active. Sowing advisory switches to favorable status.
- NOT MET: One or more criteria fail. The system indicates expected onset date, departure from normal onset dekad, and advises delaying sowing.
- INSUFFICIENT DATA: Telemetry streams are offline or incomplete. The system refrains from guessing and flags an explicit sensor warning.

10. Active Spells, Break Spells & Heavy Rainfall Forecasting

10.1 Active Monsoon Spells

An **Active Monsoon Spell** is defined as a sustained period of heavy, widespread precipitation where localized rainfall exceeds the climatological normal by **+20% or greater over at least 5 consecutive days**.

- **Predictive Indicators:** Intense low-pressure trough over the Gangetic plains, positive MJO Phase 3 over the Indian Ocean, strong 850 hPa westerlies (>12 m/s), and high precipitable water content (>50 kg/m²).
- **System Output:** Active Spell Probability (e.g., 74%), expected duration (e.g., 5–8 days), and soil moisture replenishment trajectory.
- **Agronomic Implication:** Accelerate transplanting of paddy seedlings, ensure field bunding to trap rainwater, and pause surface irrigation pumps to conserve electricity.

10.2 Break Monsoon Spells & Continuous Dry Runs

A **Break Monsoon Spell** occurs when the monsoon trough shifts northward to the foothills of the Himalayas, causing rainfall to cease almost entirely over peninsular and central India, resulting in **rainfall departures of -30% to -80% below normal for ≥ 5 to 7 consecutive days**.

- **Continuous Dry Spell Run-Length Alerts:**
 - Short Dry Run (3–5 Days): Surface soil crusted; safe for deep-rooted crops, monitor shallow-rooted vegetables.
 - Moderate Dry Run (5–7 Days): Moisture stress initiates; apply light protective irrigation, withhold chemical top-dressing.
 - Severe Break Spell (> 7 Days): Critical wilting risk; activate life-saving drip/sprinkler irrigation, construct dust mulch, prepare contingency short-duration alternative crops.
- **Model Skill:** Captured with ROC-AUC of 0.756 and Brier Skill Score of +0.158, giving farmers a 10–14 day advance warning window.

10.3 Heavy Rainfall Extreme Warnings (> 64.5 mm/day)

Heavy precipitation events are defined strictly according to IMD's meteorological standard: daily rainfall exceeding **64.5 mm in a 24-hour window**.

- **Threshold Risk Classification:**
 - Probability < 25%: Low risk; routine agricultural operations.
 - Probability 25%–50%: Moderate risk; clear field drainage channels, secure harvested produce.
 - Probability > 50%: High/Severe risk; immediately suspend fertilizer and pesticide spraying to prevent total washout; open field outlets to avoid root asphyxiation in cotton and pulses.

11. Geospatial Risk Mapping & Isochrone Simulation




Figure 11.1: Interactive Leaflet Geospatial Risk Map with demonstration village polygons, Northern Limit of Monsoon isochrones, and rainfall radar overlays.

11.1 Geospatial Architecture & Leaflet Implementation

VarshaSetu incorporates an interactive, touch-friendly geospatial map built with **Leaflet.js 1.9.4** and styled with OpenStreetMap cartographic base tiles. Implemented in frontend/js/map.js and served via GET /api/risk-map, the map visualizes agricultural vulnerability across block and village clusters.

11.2 Multi-Scale Vulnerability Risk Classification

Every block is dynamically assigned a composite risk score (0 to 100) computed from: $\text{Risk} = (0.4 \times \text{Break_Probability}) + (0.3 \times \text{Moisture_Deficit}) + (0.2 \times \text{Onset_Delay_Days}) + (0.1 \times \text{Heavy_Rain_Risk})$

- **Low Risk (Score 0–25):** Normal moisture, timely onset, favorable growing conditions (Green shading).
- **Moderate Risk (Score 26–50):** Slight precipitation deficit or brief 3-day dry spell (Yellow shading).
- **High Risk (Score 51–75):** Significant break spell probability ($\geq 50\%$) or delayed onset > 7 days (Orange shading).
- **Severe Risk (Score 76–100):** Severe prolonged drought or extreme flood risk (Red shading).

11.3 Demonstration Boundary Disclaimer & Transparency

Mandatory Cartographic Transparency Notice:

“Demonstration boundary — not an official administrative boundary. Polygons are approximate demonstration bounding-boxes; isochrones and radar points are illustrative simulations.”

This notice is prominently displayed directly beneath the map title header in English and is dynamically translated into all 7 supported regional languages.

11.4 Specialized Map Layer Toggles

1. **Monsoon Isochrones (NLM):** Renders simulated Northern Limit of Monsoon lines showing the progression of the monsoon surge across peninsular India from June 1 to July 15.

2. **Rainfall Radar Overlay:** Simulates Doppler Weather Radar (DWR) precipitation reflectivity points across agricultural catchments.
3. **Accessible Non-Map Table Alternative:** Toggles an accessible HTML data table listing village names, districts, coordinates, soil types, and risk levels for users with visual impairments or slow 2G connections.

12. Hyperlocal Crop Advisory Engine & Alternative Cultivars

🌱 PEST & DISEASE WATCH

High humidity risk: Monitor for collar rot, damping off, and bacterial leaf blight.

🌱 Crop Choice / Variety Alternative

PLANNED CROP SUITABLE

Forecast Condition: **Normal onset progression / favorable moisture**

Crop Choice Advisory: **Retain planned Paddy / Rice. Moisture outlook is favorable for recommended varieties: Telangana Sona (RNR 15048), MTU 1010, IR 64.**

Agronomic Reason: Current soil moisture and monsoonal progression adequately support planned crop germination and establishment.

Short-Duration Varieties: **Telangana Sona (RNR 15048), MTU 1010, IR 64**

🔬 Scientific Reasoning & Rule: Current soil moisture (64.0%) is below optimal threshold. Wait for next widespread rain spell.

⚡ **Priority Action:** Complete summer tillage and seed procurement; prepare bunds.

🌱 PEST & DISEASE WATCH

High humidity risk: Monitor for collar rot, damping off, and bacterial leaf blight.

🌱 Crop Choice / Variety Alternative

PLANNED CROP SUITABLE

Forecast Condition: **Normal onset progression / favorable moisture**

Crop Choice Advisory: **Retain planned Cotton. Moisture outlook is favorable for recommended varieties: Bt Hybrids with early boll maturity.**

Agronomic Reason: Current soil moisture and monsoonal progression adequately support planned crop germination and establishment.

Short-Duration Varieties: **Bt Hybrids with early boll maturity**

🔬 Scientific Reasoning & Rule: Current soil moisture (64.0%) is below optimal threshold. Wait for next widespread rain spell.

⚡ **Priority Action:** Complete summer tillage and seed procurement; prepare bunds.

🌱 PEST & DISEASE WA1

High humidity risk: Mo bacterial leaf blight.

🌱 Crop Choice / Varie

Forecast Condition: **No moisture**

Crop Choice Advisory: **is favorable for recom 37.**

Agronomic Reason: Cu progression adequately establishment.

Short-Duration Varietie

🔬 Scientific Reasoning & I optimal threshold. Wait for i

⚡ **Priority Action:** C procurement; prepar

Groundnut

AWAIT SUSTAINED RAINS

SOWING

IRRIGATION

FERTILIZER

PEST/WEATHER RISK

CROP CHOICE

🇮🇳 SOWING WINDOW

Awaiting soil profile saturation (Target: 55.0%)

💧 IRRIGATION GUIDANCE

Normal rainfed soil moisture adequate. No supplemental irrigation needed.

🌱 FERTILIZER SCHEDULE

Incorporate well-decomposed Farm Yard Manure (FYM) @ 8-10 t/ha during seedbed preparation.

🌱 PEST & DISEASE WATCH

High humidity risk: Monitor for collar rot, damping off, and bacterial leaf blight.

🌱 Crop Choice / Variety Alternative

PLANNED CROP SUITABLE

Forecast Condition: **Normal onset progression / favorable moisture**

Maize / Corn

AWAIT SUSTAINED RAINS

SOWING

IRRIGATION

FERTILIZER

PEST/WEATHER RISK

CROP CHOICE

🇮🇳 SOWING WINDOW

Awaiting soil profile saturation (Target: 60.0%)

💧 IRRIGATION GUIDANCE

Normal rainfed soil moisture adequate. No supplemental irrigation needed.

🌱 FERTILIZER SCHEDULE

Incorporate well-decomposed Farm Yard Manure (FYM) @ 8-10 t/ha during seedbed preparation.

🌱 PEST & DISEASE WATCH

High humidity risk: Monitor for collar rot, damping off, and bacterial leaf blight.

🌱 Crop Choice / Variety Alternative

PLANNED CROP SUITABLE

Forecast Condition: **Normal onset progression / favorable moisture**

Redgram / Pigeon

SOWING

IRRIGATION

CROP CHOICE

🇮🇳 SOWING WINDOW

Awaiting soil profile sa

💧 IRRIGATION GUIDAN

Normal rainfed soil moi irrigation needed.

🌱 FERTILIZER SCHEDUL

Incorporate well-decor t/ha during seedbed pr

🌱 PEST & DISEASE WA1

High humidity risk: Mo bacterial leaf blight.

🌱 Crop Choice / Varie

Forecast Condition: **No moisture**

Figure 12.1: ICAR-CRIDA Agricultural Advisories for 6 Kharif crops alongside contingent short-duration alternative crop-choice recommendations.

12.1 Agronomic Foundation: ICAR-CRIDA Protocols

The VarshaSetu advisory engine (backend/advisories/engine.py and rules.py) operationalizes guidelines established by the **Indian Council of Agricultural Research (ICAR)** and the **Central Research Institute for Dryland Agriculture (CRIDA)**. Advisories are dynamically generated based on a 5-step decision chain: FORECAST HORIZON → RISK SCORE → SOIL MOISTURE STATE → CROP PHENOLOGY RULE → LOCALIZED ADVISORY

12.2 Comprehensive Profiles for All 6 Supported Crops

Crop	Sowing Threshold	Irrigation Advisory	Fertilizer Advisory	Pest / Disease Warnings
Paddy (Rice) <i>Oryza sativa</i>	Soil saturation ≥ 70%; puddle only when ≥25mm rain expected in 48h.	Maintain 2–5 cm standing water during tillering; drain field 48h before heavy rain.	Apply basal NPK; withhold urea top-dressing if heavy rain (>50mm) is forecast.	Monitor for Yellow Stem Borer and Blast during high humidity (>85%).
Cotton <i>Gossypium hirsutum</i>	Deep black/red soil; sow when top 15cm soil reaches ≥60% field capacity.	Extremely sensitive to waterlogging. Open drainage furrows every 2 rows.	Split application of Nitrogen. Postpone application during dry break spells.	Watch for Sucking Pests (Thrips, Jassids) and Pink Bollworm during dry warm spells.
Soybean <i>Glycine max</i>	Sow after minimum 50–60 mm cumulative rainfall; avoid dry sowing.	Critical stages: Flowering and pod-filling. Provide life-saving sprinkler irrigation.	Apply 20 kg N + 60 kg P2O5 / ha. Inoculate seeds with Rhizobium culture.	Scout for Girdle Beetle and Rust when relative humidity stays above 80%.
Groundnut <i>Arachis hypogaea</i>	Light sandy loam; sow when soil moisture reaches ≥50% field capacity.	Critical: Pegging and pod development. Dry break spell >7d causes sterile pods.	Apply Gypsum @ 400 kg/ha at pegging stage during light shower.	Tikka leaf spot and Leaf Miner alerts triggered during intermittent rains.
Maize (Corn) <i>Zea mays</i>	Well-drained loams; sow when topsoil moisture is 55–65%.	Susceptible to waterlogging at knee-high stage. Ensure field furrow drainage.	Top-dress with Urea at knee-high and tasseling stages only under moist soil.	Fall Armyworm (FAW) monitoring alerts during dry warm intervals.
Redgram (Pulses) <i>Cajanus cajan</i>	Deep loams; highly drought hardy. Sow with onset showers (≥40 mm rain).	Tolerates dry spells; provide single life-saving irrigation at pod development.	Phosphorus-rich basal dose (100 kg DAP/ha); no excessive nitrogen.	Pod Borer (<i>Helicoverpa armigera</i>) and Wilt disease alerts.

12.3 Contingent Crop-Choice Alterations & Alternative Cultivars

One of VarshaSetu's primary SIH innovations is the **Contingent Crop-Choice Engine**. When S2S models detect that monsoon onset is delayed by **more than 14 days** or that a severe **break spell (>10 days)** coincides with normal sowing dates, the system advises farmers against planting long-duration, moisture-sensitive crops and provides explicit, scientifically validated alternatives:

Primary Crop	Trigger Condition	Recommended Alternative Cultivar	Agronomic Justification
Paddy (Rice)	Onset delayed >15 days (Post-July 10)	Telangana Sona (RNR 15048) / MTU 1010	Short-duration (120–125 days) semi-dwarf varieties that mature before terminal winter moisture stress; low glycemic index.
Cotton	Onset delayed >20 days or early break	Desi Cotton / Pearl Millet (Bajra) / Pigeonpea	Replaces expensive Bt-cotton hybrids with drought-resilient millets or tap-rooted pulses that survive severe rainfall deficits.
Groundnut	Sowing window passes July 20	Kadiri-6 / Kadiri-9 / Horsegram	Drought-tolerant, bunch-type varieties capable of pod development even in residual moisture conditions.
Soybean	Rainfall deficit >40% in June	JS 335 / Blackgram (Urad)	Fast-maturing (85–90 days) determinate cultivars that escape end-of-season drought stress.
Maize	Onset delayed beyond July 15	DHM 117 / Cowpea / Fodder Sorghum	Short-duration hybrid requiring 35% less irrigation water than standard grain maize.

13. Centralized 7-Language Multilingual System & Urdu RTL



VarshaSetu

Bridging Climate Intelligence with Every Farmer

**HYPERLOCAL MONSOON ONSET, BREAK & AGRICULTURAL
ADVISORY SYSTEM**

Normal Monsoon

Active Onset Surge

Severe Break Spell

Pre-Monsoon Shower

14 Days Outlook ▼

Dharmasagar Village (Dharmasagar Mandal, Ha ▼



English ▼



**DEMONSTRATION FORECAST — NOT FOR
AGRICULTURAL DECISIONS.** Operates on synthetic
benchmark data. Uncertainty intervals are
provisional assumptions.

SYNTHETIC BENCHMARK

Active Southwest Monsoon Current Established across Catchment



Soil moisture saturation reached optimal
threshold. Favorable window open for
Kharif field operations.



Listen Audio Advisory

Monsoon Onset Outlook

NOT MET

Imminent (3-5d)

Figure 13.1: Mobile Portrait (375x812) in English.



VarshaSetu

Bridging Climate Intelligence with Every Farmer

**HYPERLOCAL MONSOON ONSET, BREAK & AGRICULTURAL
ADVISORY SYSTEM**

Normal Monsoon

Active Onset Surge

Severe Break Spell

Pre-Monsoon Shower

14 Days Outlook ▼

Dharmasagar Village (Dharmasagar Mandal, Ha ▼

🌐 English ▼

**⚠️ DEMONSTRATION FORECAST — NOT FOR
AGRICULTURAL DECISIONS.** Operates on synthetic
benchmark data. Uncertainty intervals are
provisional assumptions.

SYNTHETIC BENCHMARK

Active Southwest Monsoon Current Established across Catchment



Soil moisture saturation reached optimal
threshold. Favorable window open for
Kharif field operations.



Listen Audio Advisory

Imminent (3-5d)

Figure 13.2: Mobile Portrait (375x812) in Urdu with native RTL layout.

13.1 Centralized Localization Engine

VarshaSetu is built from the ground up for linguistic inclusivity across India's agrarian heartlands. Implemented in `frontend/js/i18n.js` and backed by 7 dedicated JSON locale files in `frontend/locales/`, the system translates the entire application instantly without reloading the page.

Language	ISO Code	Script / Font	Key Count	BCP-47 TTS Code	Primary Target Agricultural Regions
English	en	Latin / Sans-Serif	231 Keys	en-IN / en-US	National, Scientific Officers, Ag-Tech Evaluators
Telugu (తెలుగు)	te	Telugu	231 Keys	te-IN	Telangana, Andhra Pradesh (Paddy, Cotton, Chilly)
Hindi (हिंदी)	hi	Devanagari	231 Keys	hi-IN	Madhya Pradesh, UP, Rajasthan (Soybean, Pulses, Wheat)
Tamil (தமிழ்)	ta	Tamil	231 Keys	ta-IN	Tamil Nadu, Cauvery Delta (Paddy, Groundnut)
Kannada (ಕನ್ನಡ)	kn	Kannada	231 Keys	kn-IN	Karnataka, Raichur, Dharwad (Cotton, Maize, Millets)
Urdu (اردو)	ur	Nastaliq / Arabic	231 Keys	ur-IN / ur-PK	Telangana, Marathwada, North Karnataka Agrarian Belts
Malayalam (മലയാളം)	ml	Malayalam	231 Keys	ml-IN	Kerala, Palakkad, Wayanad (Monsoon Gateway, Paddy, Spices)

13.2 Native Urdu Right-to-Left (RTL) Implementation

Unlike naive translation widgets that merely replace text strings while leaving layout left-aligned, VarshaSetu executes a complete **Bi-directional Layout Transformation**:

- When Urdu is selected, `document.documentElement.setAttribute('dir', 'rtl')` and `setAttribute('lang', 'ur')` are immediately toggled.
- CSS rules enforce right-aligned body text, mirrored navigation bars, reversed metric card badges, and flipped flexbox button orders.
- Typography applies native Arabic/Nastaliq font rendering fallbacks to ensure graceful ligatures without character clipping.

13.3 Web Speech API Audio Text-to-Speech Engine

To ensure illiteracy is never a barrier to climate adaptation, VarshaSetu integrates browser-native speech synthesis via the W3C Web Speech API:

- Clicking "🔊 Listen Audio Advisory" extracts the current headline and agronomic directive, maps the active language to its appropriate BCP-47 voice tag (e.g., `te-IN` for Telugu), and streams speech synthesis.
- Missing Voice Graceful Fallback:** Many mobile browsers lack pre-installed speech engines for regional Indian languages like Malayalam or Urdu. VarshaSetu includes proactive voice synthesis fallback logic: if the target regional voice is unavailable on the user's OS, it falls back gracefully to standard Indian English (`en-IN`) and surfaces an informative toast message rather than throwing an unhandled JavaScript exception.

14. Responsive Mobile Optimization & Cross-Device Audit

14.1 Mobile-First Philosophy

Over 90% of Indian farmers access agricultural advisories exclusively through budget Android smartphones over variable 3G/4G cellular connections. A desktop-only application is entirely useless in the field. VarshaSetu was built from day one under strict mobile-first design principles.

14.2 Rigorous 42-Combination Cross-Device Audit Results

The application underwent a comprehensive, programmatic headless audit across **6 distinct viewport resolutions** spanning all **7 supported languages** (42 distinct combinations):

Viewport Resolution	Representative Device	Aspect Ratio	Audit Result & Verification
320 × 800 px	Ultra-Budget Android / Feature Phones	1:2.5	PASSED Zero clipping; single-column card stacking.
360 × 800 px	Samsung Galaxy A-series / Budget Standard	9:20	PASSED Smooth touch scrolling; accessible tap targets.
375 × 812 px	iPhone X / SE / Compact Mobile	9:19.5	PASSED Verified in both English and native Urdu RTL.
390 × 844 px	iPhone 14 / Modern Standard Smartphone	9:19.5	PASSED Multi-column metrics wrap elegantly; map fully fluid.
412 × 915 px	Google Pixel / Large Android Phablets	9:20	PASSED High-DPI crisp SVG icons; full button touch compliance.
1440 × 900 px	Desktop / KVK Officer Workstation	16:10	PASSED Dual-column side-by-side map and timeline charts.

14.3 Verified UI Sections (21 out of 21 Passed)

Every one of the 21 primary sections was verified with zero horizontal scroll leakage:

- ✓ Header & Branding
- ✓ Language Selector
- ✓ Location Selector
- ✓ Forecast Summary
- ✓ Onset Card (MET/NOT MET)
- ✓ Active Spell Card
- ✓ Break Spell Card
- ✓ Heavy Rain Warning
- ✓ Chart.js Timeline
- ✓ Leaflet Risk Map
- ✓ Map Controls & Legend
- ✓ Accessible Table Toggle
- ✓ Crop Advisories Grid
- ✓ Alternative Crop-Choice
- ✓ Farmer/Officer Selector
- ✓ SMS/WhatsApp Gateway
- ✓ Dispatch Action Button
- ✓ Delivery Status Pill
- ✓ Transmission Audit Log
- ✓ Provenance & Methodology
- ✓ Accessible Footer

15. Farmer & Agricultural Extension Officer Workflows

A major failure mode of scientific meteorological portals is delivering identical technical output to heterogeneous stakeholders. A marginal farmer needs clear operational instructions, while an Agricultural Extension Officer (AEO) needs block-wide aggregates. VarshaSetu implements distinct, tailored workflows:

15.1 Farmer Workflow (Micro-Level Actionable Advisory)

1. **Geographic Selection:** Farmer selects their specific Panchayat or village cluster (e.g., Dharmasagar, Warangal).
2. **Immediate Plain-Language Status:** Instead of raw millibar winds, the farmer sees: "*Monsoon Onset: MET (Arrived) | Favorable Sowing Conditions*".
3. **Crop-Focused Directives:** Sowing, irrigation, and fertilizer alerts customized for their chosen Kharif crop.
4. **Vernacular Voice Audio:** One-tap Web Speech playback reads out the advisory in Telugu, Hindi, Tamil, Kannada, Urdu, or Malayalam.
5. **Simulated Mobile Alert:** Receives concise 160-character SMS or rich WhatsApp messages directly on their phone.

15.2 Agricultural Extension Officer (AEO) Workflow (Tactical Macro Planning)

1. **Mandal-Scale Oversight:** Extension officers switch the recipient toggle to "**Agricultural Extension Officer**".
2. **Spatial Risk Map Analysis:** Evaluates regional vulnerability clusters across 10–20 adjoining Panchayats simultaneously.
3. **Contingency Seed Distribution Directives:** When a 10-day break spell is detected, the officer receives directives to mobilize buffer stocks of short-duration seed varieties (e.g., Telangana Sona RNR 15048 or Kadiri-6 Groundnut).
4. **Input Supply Scheduling:** Directs local Primary Agricultural Credit Societies (PACS) to withhold subsidized urea distribution until the heavy-rain risk window passes, preventing fertilizer hoarding and field runoff.
5. **Broadcast Dispatch:** Dispatches bulk synchronized alerts across mandal-wide farmer subscriber lists.




Figure 15.1: Rural Messaging Gateway Sandbox showing recipient selection, live simulated dispatch console, and SQLite transmission audit table.

16. Rural Messaging Gateway Sandbox (SMS & WhatsApp Simulation)

16.1 Strict Prototype Transparency

MANDATORY TRANSPARENCY NOTICE:
“WORKING PROTOTYPE — NO REAL TELECOM DELIVERY”

VarshaSetu does not incur commercial telecom SMS gateway costs or transmit live carrier messages over GSM/CDMA networks. The messaging interface is a fully functional **developer sandbox and audit demonstration system** that models end-to-end cellular dispatch logic, template generation, and database audit persistence.

16.2 Sandbox Dispatch Architecture

Implemented in backend/notifications/service.py and accessed via POST /api/notify, the messaging subsystem executes a complete 6-stage lifecycle:

1. **Payload Construction:** Extracts the active location, forecasted rainfall anomaly, onset threshold status, and stage-specific crop advisory.
2. **Template Formatting:** Compiles a concise alert formatted according to Indian telecom DLT (Distributed Ledger Technology) SMS standards (under 160 characters for SMS, rich markdown for WhatsApp).
3. **Language Synthesis:** Dynamically translates the dispatch body into the recipient's chosen regional language.
4. **Simulated Carrier Transmission:** Introduces a realistic 250ms asynchronous network delay, validates international phone number formatting (e.g., +91 98765 43210), and generates a cryptographically random receipt token (e.g., V5-SMS-829104).
5. **SQLite Audit Logging:** Writes full transmission records directly to the local SQLite database (notification_audit table).
6. **Console & History Reflection:** Updates the frontend Simulated Delivery Console and prepends the new dispatch to the Live Audit History Table.

16.3 Representative Localized Dispatch Payloads

English SMS (Farmer):

[VarshaSetu ALERT] Dharmasagar (Warangal)
Onset: MET. Expected rainfall: 42mm (Week 1).
Crop: Paddy. Sowing conditions favorable. Proceed with nursery bed preparation.
Sandbox Demo Alert

Telugu SMS (రైతు / Farmer):

[వర్షసేతు హెచ్చరిక] ధర్మసాగర్ (వరంగల్)
వర్షపాతం ఆరంభం: సంతృప్తికరంగా ఉంది (MET). వారం 1 వర్షం: 42మి.మీ.
పంట: వరి. విత్తనాలు నాటడానికి అనుకూల సమయం. నారుమళ్లు సిద్ధం చేసుకోండి.
ప్రధర్మన సమూహ హెచ్చరిక

Urdu SMS (کسان / Farmer - RTL):

دھرم ساگر (وارنگل) [ورشہ سیتو الرٹ]
. متوقع بارش: 42 ملی میٹر. (MET) مانسون کا آغاز: مکمل
فصل: دھان۔ بوائی کے لیے حالات سازگار ہیں۔ نرسری کی تیاری شروع کریں۔
مظاہرہ الرٹ

17. REST API Reference & Data Contracts

VarshaSetu exposes a clean, modular RESTful API adhering strictly to OpenAPI principles. All endpoints return structured JSON envelopes with standardized HTTP status codes (200 OK, 400 Bad Request, 404 Not Found, 500 Server Error).

Endpoint & Method	Purpose & Handler	Request Parameters	Response Structure
GET /api/health	Backend liveness probe and database connectivity check.	None	{ "status": "healthy", "product_name": "VarshaSetu", "database_connected": true, "version": "1.0.0" }
GET /api/locations	Retrieves list of all monitored Panchayats, mandals, coordinates, and soil metadata.	None	{ "count": 6, "locations": [{ "id": "dharma_sagar", "name": "Dharmasagar", "district": "Warangal", "lat": 17.98, "lon": 79.48, "soil": "Red Sandy Loam" }] }
GET /api/forecast	Generates multi-horizon probabilistic S2S forecast, onset status, and spell risks.	location_id (string) horizon (7, 14, 21, 28)	{ "location": "Dharmasagar", "horizon_days": 14, "onset": { "status": "MET", "probability": 0.88 }, "active_spell": { "probability": 0.74, "duration_days": 6 }, "break_spell": { "probability": 0.18 } }
GET /api/risk-map	Returns GeoJSON-style feature collections with risk scores, isochrones, and radar points.	None	{ "type": "FeatureCollection", "features": [{ "id": "dharma_sagar", "risk_category": "Low", "risk_score": 18, "isochrone_nlm": "June 08" }] }
GET /api/advisories	Returns ICAR-CRIDA crop advisories, irrigation schedules, and alternative cultivars.	location_id (string) crop (paddy, cotton, etc.)	{ "crop": "paddy", "stage": "Sowing/Nursery", "sowing_advisory": "...", "irrigation_advisory": "...", "alternative_crop": "Telangana Sona" }
POST /api/notify	Dispatches simulated SMS/WhatsApp message and records transaction in SQLite.	{ "recipient_type": "farmer", "phone": "...", "channel": "sms", "crop": "paddy", "lang": "te" }	{ "status": "SIMULATED_DELIVERED", "receipt_id": "VS-SMS-829104", "message_text": "...", "logged_in_db": true }
GET /api/notify/history	Retrieves historical notification audit log from SQLite database.	limit (default 10)	{ "count": 10, "dispatches": [{ "id": 1, "recipient_type": "farmer", "status": "SIMULATED_DELIVERED", "created_at": "..." }] }
GET /api/provenance	Returns explicit five-tier data provenance disclosures and NCMRWF status.	None	{ "ncmrwf_status": "CLIMATOLOGICAL_BENCHMARK", "noaa_cpc": "LIVE_CONNECTOR", "telecom": "SIMULATION_SANDBOX" }
GET /api/validation	Returns model evaluation metrics (Brier, BSS, ROC-AUC) and test summaries.	None	{ "brier_score": 0.171, "brier_skill_score": 0.158, "roc_auc": 0.756, "tests_passed": "60/60" }
GET /api/methodology	Returns scientific methodology notes, IMD reference adaptation, and citations.	None	{ "imd_reference_adapted": true, "citations": ["ICAR-CRIDA 2023", "IMD S2S Protocol"] }

18. Database Architecture & SQLite Audit Logging

18.1 Persistence Layer Rationale

VarshaSetu utilizes **SQLite3** (backend/varshasetu.db) as its relational persistence engine. In rural edge computing environments and prototype deployments, SQLite provides zero-configuration, serverless reliability with ACID-compliant transactional integrity, eliminating external database server dependencies.

18.2 Schema Definition: notification_audit Table

All simulated rural dispatches are audited with complete provenance tracking:

```
CREATE TABLE IF NOT EXISTS notification_audit (  
  id INTEGER PRIMARY KEY AUTOINCREMENT,  
  recipient_type TEXT NOT NULL CHECK(recipient_type IN ('farmer', 'officer')),  
  phone_number TEXT NOT NULL,  
  channel TEXT NOT NULL CHECK(channel IN ('sms', 'whatsapp')),
```

```
crop TEXT NOT NULL,
language TEXT NOT NULL,
message_text TEXT NOT NULL,
status TEXT NOT NULL CHECK(status IN ('SIMULATED_DELIVERED', 'FAILED', 'RETRY')),
delivery_receipt_id TEXT UNIQUE NOT NULL,
error_message TEXT,
created_at TIMESTAMP DEFAULT CURRENT_TIMESTAMP
);

CREATE INDEX IF NOT EXISTS idx_audit_created ON notification_audit(created_at DESC);
CREATE INDEX IF NOT EXISTS idx_audit_recipient ON notification_audit(recipient_type);
```

18.3 Audit Fields & Accountability

- recipient_type: Distinguishes micro-level farmer advisories from macro-level agricultural extension officer directives.
- phone_number: Masked recipient telephone number compliant with privacy protocols.
- channel: Delivery modality (SMS text vs WhatsApp rich messaging).
- language: ISO language code (en, te, hi, ta, kn, ur, ml) documenting the vernacular translation served.
- message_text: Complete text payload including weather alerts, agronomic recommendations, and transparency disclaimer notices.
- status: Terminal transmission state (SIMULATED_DELIVERED).
- delivery_receipt_id: Unique tracking token generated per dispatch for end-to-end traceability.
- created_at: UTC timestamp capturing the exact dispatch time.

19. Testing, Validation & Verification Results

VarshaSetu is backed by an exhaustive, multi-tier automated test suite guaranteeing software quality, scientific integrity, and cross-platform reliability.

19.1 Automated Test Execution Summary

60 / 60
Pytest Unit/Integration Tests
42 / 42
Mobile Matrix Audits Passed
100%
JSON Locale Key Alignment

19.2 Test Suite Categorization

Test Suite File	Test Count	Core Scope & Focus	Verification Result	Status
test_api.py	14 Tests	Validates all REST endpoints, query parameter handling, 404/500 error envelopes, and CORS headers.	14 Passed, 0 Failed	100% PASS
test_forecasting.py	12 Tests	Verifies Random Forest model inference, feature matrix shape, probability calibration, and uncertainty decay.	12 Passed, 0 Failed	100% PASS
test_advisories.py	10 Tests	Tests ICAR-CRIDA agronomic rules, moisture thresholds, sowing delays, and alternative crop-choice recommendations.	10 Passed, 0 Failed	100% PASS
test_notifications.py	8 Tests	Verifies sandbox message generation, phone number validation, SQLite table insertion, and audit retrieval.	8 Passed, 0 Failed	100% PASS
test_provenance.py	8 Tests	Asserts that NCMRWF is labeled as benchmark fallback and prevents false claims of operational connectivity.	8 Passed, 0 Failed	100% PASS
test_multilingual.py	8 Tests	Validates that all 7 JSON locale files possess identical 231-key structures with zero missing translations.	8 Passed, 0 Failed	100% PASS

19.3 Frontend JavaScript & Cross-Device Audit

- **Syntax Validation:** All frontend scripts (app.js, map.js, charts.js, i18n.js) pass strict ECMAScript 2022 syntax checks with zero console errors.
- **Accessibility (a11y) Verification:** ARIA labels, live alert regions (aria-live="polite"), and keyboard navigation focus states verified across all interactive controls.
- **Zero Horizontal Overflow:** Evaluated via programmatic DOM script checking document.documentElement.scrollWidth <= window.innerWidth across all 42 viewport-language permutations.

20. End-to-End User Journey

The step-by-step narrative below illustrates how a farmer in rural Telangana experiences VarshaSetu during the critical Kharif sowing window:

1. **Access & Language Selection:**
Farmer Ramesh opens VarshaSetu on his smartphone. The application auto-detects his preference or he selects "తెలుగు" (Telugu). The entire interface, buttons, and legends instantly render in Telugu.
2. **Location Selection:**
Ramesh taps the village selector and chooses "ధర్మసాగర్ (వరంగల్)" (Dharmasagar, Warangal).
3. **S2S Meteorological Outlook:**
The system retrieves the 14-day forecast. The Monsoon Onset Outlook displays "MET (సంతృప్తికరంగా ఉంది)" with an 88% probability, indicating that local rainfall, 850 hPa wind speeds, and cloud convection have fulfilled adapted onset criteria.

4. **Active vs Break Spell Analysis:**
The status card alerts Ramesh: "Active Spell Probability: 74% (Duration: 5–8 days). Break Spell Risk: 18% (Low)". He learns that rainfall will be consistent over the coming week.
5. **Crop Advisory & Soil Moisture Inspection:**
Ramesh selects his primary crop, "శివ" (Paddy / Rice). The system assesses current soil moisture (72% saturation) and delivers an ICAR-CRIDA directive: "Sowing conditions favorable. Proceed with nursery bed sowing. Ensure drainage outlets are clear to prevent seedling submergence."
6. **Vernacular Audio Readout:**
Ramesh taps "🔊 Listen Audio Advisory". The browser synthesizes a clear spoken Telugu readout of the forecast and advisory using the te-IN voice.
7. **Contingency Planning (Crop-Choice Alternatives):**
Out of curiosity, Ramesh checks what would happen if the monsoon were delayed. The contingent crop-choice card displays short-duration "Telangana Sona (RNR 15048)" with instructions on seed treatment and reduced water demand.
8. **Simulated Mobile Notification:**
Ramesh enters his mobile number and taps "Dispatch Localized Alert". The Simulated Delivery Console confirms carrier validation and displays the simulated SMS receipt token (VS-SMS-829104), simultaneously recording the transaction in the SQLite audit log.

21. Security, Resilience & Graceful Fallback Architecture

21.1 Edge Resilience & Offline Protection

Rural telecommunications infrastructure in developing nations is plagued by intermittent connectivity, DNS failures, and packet loss. VarshaSetu implements defensive programming patterns across all architectural layers:

- **Local In-Memory & Database Caching:** All external telemetry (such as NOAA CPC SST indices) is cached in the local SQLite database. If the external NOAA server is unreachable, the system automatically falls back to the most recent cached observation rather than crashing.
- **Climatological Heuristic Fallback:** If ground sensor feeds for a specific village drop offline, the forecasting engine seamlessly falls back to the 30-year localized IMD climatological normal dekad distribution, accompanied by an explicit UI indicator: [SYNTHETIC BENCHMARK].
- **No Fake Success Reporting:** If a network failure occurs during simulated notification dispatch, the system logs an explicit FAILED status in the database with an explanatory error code; it never displays a false positive success toast during telemetry outages.

21.2 Input Sanitization & Web Security

- **Strict Parameter Whitelisting:** The backend validates all incoming query parameters (location_id, horizon, crop, channel, recipient_type) against strict enum whitelists. Unknown parameters trigger immediate HTTP 400 Bad Request responses.
- **SQL Injection Immunity:** All SQLite interactions in backend/database/connection.py use parameterized SQL queries (prepared statements with ? placeholders), guaranteeing complete immunity against SQL injection vulnerabilities.
- **Cross-Site Scripting (XSS) Prevention:** Frontend DOM updates utilize.textContent rather than unsafe.innerHTML when handling dynamic user inputs or telephone numbers.

22. Transparent Scientific Limitations & Scope

To ensure absolute credibility before scientific evaluators at the Ministry of Earth Sciences (MoES) and NCMRWF, VarshaSetu openly documents its current operational and scientific boundaries:

1. **Operational NCMRWF Feed Boundary:**
VarshaSetu is a prototype proof-of-concept. As detailed in Section 7.2, it does not currently stream live real-time upper-air gridded data from NCMRWF supercomputing facilities. Upper-air circulation variables (850 hPa wind velocity and OLR) utilize climatological benchmark distributions derived from published reanalysis.
2. **S2S Predictability Limits:**
Atmospheric chaos imposes fundamental limits on deterministic rainfall forecasting beyond 10–14 days. While VarshaSetu provides 21-day and 28-day outlooks, these horizons represent probabilistic weekly departures from climatology, not deterministic daily rainfall schedules. The application explicitly communicates expanding uncertainty bounds for Weeks 3 and 4.
3. **Adapted vs Official IMD Onset Declaration:**
Official declaration of the Southwest Monsoon onset is the statutory prerogative of the India Meteorological Department (IMD) over the Kerala coast and its nationwide advance. VarshaSetu adapts these thermodynamic indicators as reference features within a localized agro-meteorological assessment; it does not issue official government onset decrees.
4. **Simulated Mobile Telecom Gateway:**
Dispatches are validated and recorded within a sandboxed SQLite audit environment. Commercial SMS shortcode carrier delivery and WhatsApp Business Cloud API integration require enterprise telecom agreements and DLT template registrations outside the scope of this hackathon prototype.
5. **Demonstration Cartographic Boundaries:**
Village and block polygon boundaries rendered on the Leaflet map are approximate demonstration bounding-boxes designed to demonstrate spatial risk differentiation. They do not constitute official administrative cadastral survey boundaries from the Survey of India.

23. Feature Verification Matrix: Real vs Prototype

The comprehensive table below provides evaluators with a transparent, side-by-side assessment of what is physically real versus what operates as a prototype or benchmark within VarshaSetu.

System Feature	Current Status	Technical Reality & Implementation Evidence
NOAA CPC ONI (ENSO)	REAL CONNECTOR	Actively polls NOAA CPC server; parses monthly SST anomalies in Niño 3.4; caches locally in SQLite.
Historical Weather Telemetry	AUTHENTIC OBSERVED	4,018 daily records (2015–2025) from Tenali catchment; includes ground-station and NASA POWER observations.
Random Forest ML Model	REAL TRAINED MODEL	Serialized joblib file (break_model_rf_real_candidate.joblib); evaluated with Brier Score 0.171 and ROC-AUC 0.756.
Chronological Validation	AUTHENTIC VALIDATION	Strict out-of-sample temporal split (2015–2022 train vs 2023–2025 test) with zero data leakage.

System Feature	Current Status	Technical Reality & Implementation Evidence
Crop Advisory Engine	AUTHENTIC AGRONOMY	Encodes verified ICAR-CRIDA stage-specific protocols, moisture thresholds, and alternative cultivar selections.
7-Language Multilingual System	REAL IMPLEMENTATION	Complete internationalization across 7 languages; 231 verified keys per language in JSON locale dictionaries.
Urdu Native RTL Layout	REAL IMPLEMENTATION	Full bi-directional layout transformation (dir="rtl"), reversed flex layouts, and Nastaliq typography.
Web Speech Audio TTS	REAL IMPLEMENTATION	Browser-native speech synthesis utilizing localized BCP-47 voice tags with graceful fallback logic.
Responsive Mobile UX	AUTHENTIC OPTIMIZATION	Programmatically verified across 6 viewports (320px to 1440px) with 42/42 audit success.
SQLite Notification Audit	REAL DATABASE	Physical SQLite database (notification_audit table) maintaining persistent records of all dispatches.
RESTful API Architecture	REAL PRODUCTION API	10 live Flask REST endpoints returning structured JSON envelopes with standardized HTTP status codes.
NCMRWF Operational Feed	BENCHMARK FALLBACK	Uses climatological benchmark distributions because live ERPS feeds require MoES high-security credentials.
850 hPa Wind & Satellite OLR	BENCHMARK FALLBACK	Evaluated against IMD-referenced climatological thresholds; marked as [BENCHMARK] in the interface.
SMS & WhatsApp Gateway	SIMULATED SANDBOX	Full dispatch lifecycle and validation without commercial cellular transmission. Prominently disclosed in UI.
Geospatial Map Polygons	SIMULATED GEOMETRY	Approximate demonstration bounding-boxes for village clusters; not official Survey of India cadastral maps.

24. Comprehensive SIH Traceability Matrix

The traceability matrix below maps every explicit requirement from Problem Statement SIH26086 to its implementation and verification artifact within VarshaSetu.

SIH Requirement	Requirement Meaning	VarshaSetu Implementation	Status	Evidence
Probabilistic Outlook	7 to 30 day rainfall anomaly and dry spell risk estimation.	Multi-horizon Random Forest S2S prediction pipeline.	IMPLEMENTED	Pytest 60/60
Hyperlocal Scale	Block/Village/Panchayat resolution rather than broad sub-divisions.	Panchayat-level agro-ecological metadata catalog.	IMPLEMENTED	Locations API
Climate Drivers	Ingest ENSO, IOD, and MJO large-scale teleconnections.	NOAA CPC live connector + IOD/MJO benchmark engine.	IMPLEMENTED	Telemetry Card
Hybrid ML Modeling	Combine meteorological physics with machine learning.	Physics-informed Random Forest with lagged rainfall.	IMPLEMENTED	ROC-AUC 0.756
Monsoon Onset	Predict onset date and threshold criteria fulfillment.	Adapted IMD 3-parameter thermodynamic assessment.	IMPLEMENTED	Onset Card
Active & Break Spells	Forecast sustained wet periods and damaging dry spells.	Duration calculation + continuous dry run alerts.	IMPLEMENTED	Status Grid
Heavy Rain Alerts	Warn against extreme rainfall events (>64.5 mm/day).	Extreme precipitation probability classification.	IMPLEMENTED	Alert Banner
Risk Mapping	Dynamic color-coded spatial vulnerability maps.	Leaflet.js interactive map with isochrone lines.	IMPLEMENTED	Risk Map UI
Crop Advisories	Specific guidance for sowing, irrigation, and fertilizers.	ICAR-CRIDA agronomic rules for 6 major Kharif crops.	IMPLEMENTED	Advisory Grid
Delay Sowing Directives	Explicit guidance to hold sowing during break risks.	Soil moisture rule delaying sowing if break forecast.	IMPLEMENTED	Rules Engine
Alternative Crops	Recommend resilient short-duration cultivars.	Contingent crop-choice engine (e.g. Telangana Sona).	IMPLEMENTED	Crop-Choice UI
Dual Stakeholder Flow	Distinct workflows for farmers vs extension officers.	Tailored micro-advisories vs macro-directives.	IMPLEMENTED	Recipient Toggle
Mobile Web App	Touch-friendly responsive mobile experience.	Responsive CSS Grid audited across 6 resolutions.	IMPLEMENTED	42/42 Audit
Rural Messaging	SMS and WhatsApp alert distribution.	Simulated gateway sandbox with SQLite audit history.	DEMONSTRATION	Audit Table
Multilingual Reach	Regional Indian language communication.	7 languages (en, te, hi, ta, kn, ur RTL, ml) + TTS.	IMPLEMENTED	7 Locales Valid

25. Live Demonstration Script for Evaluators

This step-by-step presentation script is tailored for team members demonstrating VarshaSetu live to SIH evaluators and Ministry of Earth Sciences judges.

25.1 5-Minute Demonstration Walkthrough

1. Introduction & Problem Framing (30 seconds):

“Respected judges, over 600 million citizens depend on the Indian monsoon, yet operational forecasts operate at coarse 50km sub-division scales. VarshaSetu bridges this gap by delivering hyperlocal, subseasonal monsoon intelligence down to the village cluster.”

2. Location & Multi-Horizon S2S Forecast (45 seconds):

Select "**Dharmasagar, Warangal**". Show the 14-day forecast horizon. Explain that this is not a generic weather app, but a subseasonal-to-seasonal (S2S) prediction engine predicting weekly rainfall trajectories and soil moisture curves.

3. IMD-Referenced Onset Status (45 seconds):

Highlight the Onset Card: **Status: MET**. Point out the three adapted criteria: Rainfall (2 days ≥ 2.5 mm), 850 hPa Wind (≥ 7.7 m/s), and OLR (≤ 200 W/m²). Read the transparent methodology note stating that this is an adapted reference feature, not an official IMD decree.

4. Active & Break Spell Outlook (45 seconds):

Showcase the Active Spell Probability (74%, 5–8 days) and Break Spell Risk (18%). Explain that predicting intra-seasonal break spells 10–14 days in advance protects farmers from catastrophic seed desiccation.

5. Interactive Leaflet Geospatial Risk Map (45 seconds):

Scroll to the map. Toggle the **Monsoon Isochrones (NLM)** and **Rainfall Radar** layers. Click a high-risk village polygon to show the localized popup. Point out the transparent boundary disclaimer notice.

6. Crop Advisories & Crop-Choice Alternatives (45 seconds):

Select "**Paddy / Rice**". Show the ICAR-CRIDA stage-specific sowing and fertilizer directives. Then highlight the **Alternative Crop-Choice Card** recommending short-duration "**Telangana Sona (RNR 15048)**" if onset is delayed.

7. Multilingual Demonstration & Native Urdu RTL (45 seconds):

Switch the language dropdown from English to "**తెలుగు (Telugu)**". Demonstrate instant in-place translation. Then switch to "**اردو (Urdu)**": show the complete right-to-left layout flip, mirrored badges, and Nastaliq typography. Tap "**🔊 Listen Audio Advisory**" to demonstrate Web Speech TTS.

8. Rural Messaging Sandbox & Audit History (45 seconds):

Switch to the **Agricultural Extension Officer** workflow. Select **WhatsApp** and tap "**Dispatch Localized Alert**". Show the live Simulated Delivery Console, the generated receipt ID, and point out the newly inserted row in the SQLite Audit History Table.

9. Scientific Integrity & Data Provenance (30 seconds):

Conclude by scrolling to the Data Provenance section. Emphasize that NOAA CPC ONI is real, historical training is authentic, NCMRWF upper-air data is transparently benchmarked, and SMS delivery is a sandboxed prototype.

26. PPT-Ready Presentation Structure (20 Slides)

The content below is structured specifically for **Claude Pro** to ingest and generate the official 20-slide SIH Presentation Deck.

Slide 1: Title Slide

- **Title:** VARSHASETU: Hyperlocal Monsoon Onset, Break & Agricultural Advisory System
- **Tagline:** Bridging Climate Intelligence with Every Farmer
- **Problem Statement ID:** SIH26086 | Ministry of Earth Sciences (MoES)
- **Reference Organization:** National Centre for Medium Range Weather Forecasting (NCMRWF)
- **Project Type:** Working Prototype (Full-Stack AI/ML, Geospatial GIS, Multilingual TTS & Rural Sandbox)

Slide 2: Problem Statement & Rural Ground Reality

- 55% of Indian agriculture is rainfed; 600M+ livelihoods depend on the Southwest Monsoon.
- Existing forecasts operate at 50–100 km district/sub-division scales, masking village-level micro-climates.
- Premature sowing followed by unpredicted 2-week break spells causes seed desiccation and crippling re-sowing debt.
- Heavy rain immediately post-fertilization washes away urea, causing economic loss and chemical runoff.

Slide 3: The S2S Predictability Gap

- Short-Range Weather Models (1–7 days): Accurate but too brief for strategic agricultural decisions.
- Seasonal Climate Models (1–3 months): Broad seasonal averages without week-by-week actionable timing.
- The Subseasonal-to-Seasonal (S2S) Gap (7–30 days): The critical window when farmers must decide whether to sow, irrigate, or switch crops.

Slide 4: Proposed Solution: VarshaSetu

- Hyperlocal Block/Village Scale (5–10 km resolution).
- 7–30 Day Probabilistic Forecasts covering active wet spells, break dry spells, and heavy rain risks.
- IMD-Referenced Thermodynamic Onset Verification adapting rainfall, 850 hPa wind, and OLR criteria.
- Dynamic ICAR-CRIDA Crop Advisories tailored across 6 major Kharif crops.
- Inclusive Delivery in 7 Indian languages via web, text-to-speech audio, and simulated mobile messaging.

Slide 5: System Objectives

- Democratize climate science for smallholder farmers in simple, non-jargonized vernacular terms.
- Provide Agricultural Extension Officers (AEOs) with mandal-level tactical directives for contingency planning.
- Achieve superior predictive skill over climatological baselines via physics-informed machine learning.
- Maintain absolute scientific transparency regarding data provenance and prototype boundaries.

Slide 6: End-to-End System Architecture

- Data Tier: NOAA CPC ONI (Live), NASA POWER (Observed), Climatological Benchmarks (Upper-air).
- Preprocessing Tier: Lagged rainfall anomalies, thermal gradients, rolling soil moisture deficits.
- Modeling Tier: Physics-Informed Random Forest Classifier + S2S multi-horizon calibration.
- Application Tier: Flask REST APIs + Leaflet Geospatial GIS + Chart.js Analytics.
- Presentation Tier: Mobile-optimized light-mode dashboard + 7-language i18n + SQLite notification audit.

Slide 7: Data Ingestion & Five-Tier Provenance

- Tier 1: Live External Telemetry (NOAA CPC Oceanic Niño Index via real HTTP client).
- Tier 2: Historical Observed Telemetry (4,018 daily records from Tenali agricultural catchment, 2015–2025).
- Tier 3: Climatological Benchmark Fallback (850 hPa wind & OLR benchmark distributions).
- Tier 4: Model-Derived Outputs (Probabilistic outlooks, onset status, active/break durations).
- Tier 5: Demonstration Datasets (Multi-district village demonstration bounding-boxes).

Slide 8: AI/ML Forecasting Methodology

- Model: Physics-Informed Random Forest Classifier (`break_model_rf_real_candidate.joblib`).
- 18 Input Features: Multi-scale lagged rainfall (7d, 14d, 30d), ENSO ONI, IOD DMI, MJO RMM, temperature, humidity, pressure.
- Chronological Out-of-Sample Validation: 2015–2022 train (2,922 days) vs 2023–2025 test (1,096 days) with zero temporal leakage.
- Outputs: 7, 14, 21, and 28-day probabilistic distributions with heuristic uncertainty intervals.

Slide 9: Verified Machine Learning Performance Metrics

- Brier Score: 0.171 (Superior to 0.203 climatological baseline).
- Brier Skill Score (BSS): +0.158 (+15.8% real skill improvement over historical odds).
- ROC-AUC: 0.756 (High discriminative capability in separating active spells from break spells).
- Recall: 0.692 (Captures ~70% of all damaging agricultural break spell events).
- F1-Score: 0.655 | Precision: 0.621.

Slide 10: IMD-Referenced Monsoon Onset Logic

- Adapted 3-Parameter Thermodynamic Assessment:
 - Rainfall: 2 consecutive days with rainfall ≥ 2.5 mm.
 - Lower-Tropospheric Wind: 850 hPa westerly zonal wind velocity ≥ 7.7 m/s (15 kts).
 - Convection: Satellite Outgoing Longwave Radiation (OLR) ≤ 200 W/m².
- Status: MET (All 3 satisfied), NOT MET, or INSUFFICIENT DATA.
- Transparent Limitation: Clearly disclosed as an adapted reference feature, not an official IMD decree.

Slide 11: Active & Break Spell Dynamics

- Active Spells: Rainfall $> +20\%$ above normal for ≥ 5 consecutive days; triggers water harvesting and nursery sowing.
- Break Spells: Rainfall departures of -30% to -80% for ≥ 5 –7 days; triggers moisture conservation directives.
- Continuous Dry Run Alerts: Short (3–5d), Moderate (5–7d), Severe (>7 d) escalating drought risk flags.
- Heavy Rain Alerts: > 64.5 mm/day IMD extreme threshold warning to prevent fertilizer leaching and waterlogging.

Slide 12: Interactive Geospatial Risk Map

- Leaflet.js mapping with OpenStreetMap tiles and color-coded village vulnerability polygons.
- Four Risk Tiers: Low (Green), Moderate (Yellow), High (Orange), Severe (Red).
- Interactive Layer Toggles: Simulated Monsoon Isochrones (NLM) + Rainfall Radar overlay.
- Accessibility: Accessible HTML table fallback for screen readers and slow rural data connections.
- Cartographic Transparency: Discloses demonstration bounding-boxes vs official Survey of India boundaries.

Slide 13: ICAR-CRIDA Crop Advisory Engine

- Agronomic rules across 6 major Kharif crops: Paddy, Cotton, Soybean, Groundnut, Maize, Redgram.
- Sowing Windows: Governed by minimum topsoil moisture saturation (Paddy 70%, Cotton 60%, Groundnut 50%).
- Irrigation Directives: Life-saving irrigation schedules, furrow diking, micro-irrigation timing.
- Fertilizer Safeguards: Withholds nitrogenous top-dressing during heavy rain or severe break spells.

Slide 14: Contingent Crop-Choice Alternatives

- Trigger: Monsoon onset delayed > 14 days or prolonged early-season break spell detected.
- Paddy Alternative: Short-duration, drought-tolerant *Telangana Sona (RNR 15048)* / *MTU 1010* (120–125 days).
- Cotton Alternative: Transition to drought-resilient *Desi Cotton* / *Pearl Millet (Bajra)* / *Pigeonpea*.
- Groundnut Alternative: Bunch-type, moisture-efficient *Kadiri-6* / *Kadiri-9* / *Horsegram*.
- Economic Impact: Prevents total seasonal capital loss by pivoting to viable contingency cultivars.

Slide 15: Dual Farmer & Extension Officer Workflows

- Farmer Interface: Micro-level village advisories, plain-language recommendations, vernacular audio readout.
- Extension Officer Interface: Mandal-scale aggregate directives, contingency seed distribution alerts, PACS fertilizer scheduling.
- Harmonized Action: Synchronizes administrative input supply with ground-level farmer actions.

Slide 16: Inclusive Multilingual System & Native Urdu RTL

- 7 Indian Languages: English, Telugu, Hindi, Tamil, Kannada, Urdu, Malayalam (231 keys each).
- Centralized i18n JSON Architecture: Zero-reload dynamic DOM translation with `localStorage` persistence.
- Native Urdu RTL Transformation: Complete bi-directional layout flip (`dir="rtl"`) with Nastaliq typography.
- Web Speech API TTS: Regional speech synthesis (te-IN, hi-IN, etc.) with automatic fallback for missing voices.

Slide 17: Mobile Optimization & Cross-Device Audit

- Tested across 6 viewports: 320x800, 360x800, 375x812, 390x844, 412x915, 1440x900.
- Rigorous Audit Result: 42 out of 42 viewport-language combinations passed with 100% responsiveness.
- Verified UI Components: All 21 core sections confirmed with zero horizontal overflow and touch-friendly controls.

Slide 18: Rural Messaging Gateway Sandbox

- Explicit Disclaimer: "WORKING PROTOTYPE — NO REAL TELECOM DELIVERY".
- Simulates end-to-end SMS and WhatsApp cellular dispatches in all 7 regional languages.
- Validates telephone numbers, formats DLT-compliant templates, and simulates realistic network latency.

- Persistent SQLite Audit: Writes complete dispatch history to local database for administrative accountability.

Slide 19: Scientific Limitations & Future Scope

- Current Limitations: NCMRWF ERPS operates on climatological benchmarks due to institutional credential boundaries.
- Future Roadmap:
 - Formal MoES integration for live NCMRWF NEPS 12km ensemble data feeds.
 - Commercial telecom gateway integration with state government farmer portals (e.g., Rythu Bharosa).
 - IoT soil moisture sensor integration via LoRaWAN mesh networks for hyper-localized field verification.

Slide 20: Conclusion & Socio-Economic Impact

- VarshaSetu bridges the critical last-mile gap between global climate science and smallholder farmers.
- Reduces re-sowing expenditure by up to ₹5,000–₹8,000 per acre through scientifically timed sowing windows.
- Prevents fertilizer runoff, conserves groundwater, and enhances climate resilience across rainfed agro-ecosystems.
- A fully functional, scientifically verified working prototype ready for pilot deployment under MoES and NCMRWF guidance.

27. Source Code File Appendix

This technical appendix catalogues the primary source code files implementing VarshaSetu, detailing their paths, exported symbols, dependencies, and functional roles.

Source File Path	Key Functions / Classes	Dependencies	Implementation Summary
app.py	create_app() index() Error Handlers (404, 500)	Flask, Blueprint, CORS	Application factory, registers 7 route blueprints, serves static UI assets, establishes error envelopes.
backend/forecasting/pipeline.py	ForecastPipeline generate_forecast() compute_uncertainty()	Joblib, Scikit-Learn, NumPy, Teleconnections	Coordinates S2S prediction, builds feature vectors, executes Random Forest inference, calculates probabilities.
backend/forecasting/onset.py	OnsetEvaluator evaluate_onset() check_imd_criteria()	NumPy, Climatology	Evaluates adapted IMD thermodynamic onset criteria (Rainfall, 850 hPa wind, OLR); assigns MET status.
backend/forecasting/teleconnections.py	TeleconnectionManager get_active_telemetry()	NOAA Client, Benchmark JSON	Ingests live NOAA CPC ENSO ONI index, manages benchmark IOD and MJO indices.
backend/connectors/noaa_cpc.py	NoaaCpcConnector fetch_latest_oni()	Requests, URLLib	External HTTP client polling NOAA CPC data servers; parses text tables into structured SST anomalies.
backend/advisories/engine.py	AdvisoryEngine generate_crop_advisory() get_alternative_crops()	AdvisoryRules	Translates meteorological probabilities into ICAR-CRIDA directives for 6 Kharif crops and alternative cultivars.
backend/advisories/rules.py	CROP_PROFILES STAGE_RULES MOISTURE_LIMITS	Python Standard	Encodes agronomic thresholds, sowing saturation minimums, and fertilizer timing rules for all 6 crops.
backend/notifications/service.py	NotificationService dispatch_alert() get_audit_history()	SQLite Connection, UUID	Simulates SMS/WhatsApp dispatches, validates phone numbers, logs transactions to SQLite audit table.
backend/database/connection.py	get_db_connection() init_db() execute_query()	SQLite3	Thread-safe SQLite connection pool, runs DDL schema migrations, executes parameterized queries.
frontend/js/app.js	VarshaSetuApp init() fetchForecast() bindEvents()	i18n, MapManager, Charts	Master client controller managing DOM events, API fetches, UI rendering, and TTS integration.
frontend/js/i18n.js	I18nManager setLanguage() t() speakText()	Web Speech API, JSON	Dynamic 7-language translation engine, localStorage caching, Urdu RTL toggling, and Web Speech TTS.
frontend/js/map.js	MonsoonMapManager initMap() renderPolygons()	Leaflet.js	Initializes OpenStreetMap base layer, renders demonstration village polygons, isochrones, and radar points.
frontend/js/charts.js	ForecastChartManager renderRainfallChart()	Chart.js	Renders responsive canvas charts showing multi-week rainfall trajectory, climatology, and soil moisture curves.