

# Tips, Formulae and shortcuts for Mixtures and Alligations

By

**CRACKU.IN**

## Cracku Tip 1 – Mixtures and Alligations

- The topic mixtures and alligations is basically an application of averages concept in CAT.
- The theory involved in this topic is very limited and students should be comfortable with the some basic formulas and concepts.
- This pdf covers all the important formulas and concepts related to mixtures and alligations.

For free CAT mocks visit <https://cracku.in/cat>

## Cracku Tip 2 – Mixtures and Alligations

A mixture is created when two or more substances are mixed in a certain ratio.

### Types of mixtures

#### 1. Simple mixture


A simple mixture is formed by the mixture of two or more different substances.

Ex. Water and Wine mixture

#### 2. Compound mixture

Compound mixture is formed by the mixture of two or more simple mixtures.

## Cracku Tip 3 – Mixtures and Alligations



If  $M_1$  and  $M_2$  are the values,  $Q_1$  and  $Q_2$  are the quantities of item 1 and item 2 respectively and  $M_A$  is the weighted average of the two items, then

$$\frac{Q_1}{Q_2} = \frac{M_2 - M_A}{M_A - M_1}$$

Weighted average  $M_A$  can be calculated by,  $M_A = \frac{Q_1 M_1 + Q_2 M_2}{Q_1 + Q_2}$

## Cracku Tip 4 – Mixtures and Alligations

The allegation rule can be applied when cheaper substance is mixed with expensive substance

$$\frac{\text{Quantity of cheaper}}{\text{Quantity of dearer}} = \frac{\text{Price of dearer} - \text{Mean price}}{\text{Mean price} - \text{Price of cheaper}}$$

Personal guidance for CAT by IIM alumni

<https://www.facebook.com/groups/CrackuCATExam/> -

## Cracku Tip 5 – Mixtures and Alligations

If we have two mixtures  $M_1$  and  $M_2$  and each mixture having two substances  $S_1$  and  $S_2$  in the ratio  $a:b$  and  $p:q$  are mixed and a final solution is obtained, then in the final solution

$$\frac{\text{Quantity of } S_1}{\text{Quantity of } S_2} = \frac{M_1 \left[ \frac{a}{a+b} \right] + M_2 \left[ \frac{p}{p+q} \right]}{M_1 \left[ \frac{b}{a+b} \right] + M_2 \left[ \frac{q}{p+q} \right]}$$

## Cracku Tip 6 – Mixtures and Alligations

If there is a container with 'a' liters of liquid A and if 'b' liters are withdrawn and equal amount is replaced by another liquid B and if the operation is repeated for 'n' times

After nth operation,

- Liquid A in the container =  $\left[\frac{a-b}{a}\right]^n \times \text{Initial quantity of A in the container}$
- $\frac{\text{Liquid A after nth operation}}{\text{Liquid B after nth operation}} = \frac{\left[\frac{a-b}{a}\right]^n}{1 - \left[\frac{a-b}{a}\right]^n}$

**For a free guidance for CAT  
from IIM alumni**

**WhatsApp 'CAT' to +91 7661025559**



CRACKU.IN is an online learning platform for CAT, XAT, Bank exams, RRB, SSC and many other exams

Personal guidance for CAT by IIM alumni  
<https://www.facebook.com/groups/CrackuCATExam/> -

Daily free Live session for CAT <https://cracku.in/cat/classroom>

For free CAT mocks visit <https://cracku.in/cat>