

Tips, Formulae and shortcuts for Logarithms, surds and indices

By

CRACKU.IN

Cracku Tip 1 – Logarithms, Surds and Indices

- “Logarithms, Surds and Indices” is one of the easiest topics in the quantitative section of the CAT exam.
- Although the number of formulae is high, the basic concepts are very simple to understand and apply.
- The scope of the questions that can be asked from this topic is very limited.
- The accuracy of answering questions from this section is very high and well prepared aspirants tend to score very well here.

For free CAT mocks visit <https://cracku.in/cat>

Cracku Tip 2 – Logarithms, Surds and Indices

If $X, Y > 0$ and m, n are rational numbers then

- $X^m \times X^n = X^{m+n}$
- $X^0 = 1$
- $\frac{X^m}{X^n} = X^{m-n}$
- $(X^m)^n = X^{mn}$
- $X^m \times Y^m = (X \times Y)^m$
- $\frac{X^m}{Y^m} = (X/Y)^m$
- $X^{-m} = \frac{1}{X^m}$

For free CAT mocks visit <https://cracku.in/cat>

Cracku Tip 3 – Logarithms, Surds and Indices

If X and Y are positive real numbers and a,b are rational numbers.

- $(X/Y)^{-a} = (Y/X)^a$
- $X^{1/a} = \sqrt[a]{X}$
- $X^{a/b} = \sqrt[b]{X^a}$
- $\sqrt[a]{X} \times \sqrt[a]{Y} = \sqrt[a]{XY}$
- $\sqrt[a]{X} / \sqrt[a]{Y} = \sqrt[a]{X/Y}$
- $\frac{1}{\sqrt{N+1}-\sqrt{N}} = \sqrt{N+1} + \sqrt{N}$

For free CAT mocks visit <https://cracku.in/cat>

Cracku Tip 4 – Logarithms, Surds and Indices

- Surd is an irrational number involving a root ex : $\sqrt{5}$, $\sqrt[3]{7}$, $\sqrt[5]{2}$
- Like surds are two surds having same number under radical sign.
- Like surds can be added or subtracted. $6\sqrt{2} + 3\sqrt{2} = 9\sqrt{2}$

Cracku Tip 5 – Logarithms, Surds and Indices

- If $a + \sqrt{b} = c + \sqrt{d}$, then $a = c$ and $b = d$
- The conjugate of $a + \sqrt{b}$ is $a - \sqrt{b}$
- $\sqrt{a\sqrt{a\sqrt{a\ldots\infty}}} = a$
- $\sqrt{a\sqrt{a\sqrt{a\ldots x \text{ times}}}} = a^{1 - [1/(2^x)]}$
- To find $\sqrt{\sqrt{x} + \sqrt{y}}$, $\sqrt{x} + \sqrt{y}$ should be written in the form of $m+n+2\sqrt{mn}$ where $x = m+n$ and $4mn = y$ and $\sqrt{\sqrt{x} + \sqrt{y}} = \pm (\sqrt{m} + \sqrt{n})$

Cracku Tip 6 – Logarithms, Surds and Indices

If $N = a^x$ then, x is defined as the logarithm of N to base a
or $x = \log_a N$

Logarithm of a negative number or zero is not defined.

- $\log_a 1 = 0$
- $\log_a xy = \log_a x + \log_a y$
- $\log_a b^c = c \log_a b$
- $\log_a a = 1$
- $x^{\log_b y} = y^{\log_b x}$

Cracku Tip 7 – Logarithms, Surds and Indices

- $\log_a \sqrt[n]{b} = \frac{\log_a b}{n}$
- $\log_a x = \frac{1}{\log_x a}$
- $b^{\log_b x} = x$
- $\log_a b = \frac{\log_c b}{\log_c a}$
- $\log_a b * \log_b a = 1$
- $\log_a (X/Y) = \log_a X - \log_a Y$

Cracku Tip 8 – Logarithms, Surds and Indices

- If $0 < a < 1$, then $\log_a x < \log_a y$ (if $x > y$)
- If $a > 1$ then $\log_a x > \log_a y$ (if $x > y$)

**For a free guidance for CAT
from IIM alumni**

WhatsApp 'CAT' to +91 7661025559

CRACKU.IN is an online learning platform for CAT, XAT, Bank exams, RRB, SSC and many other exams

Personal guidance for CAT by IIM alumni
<https://www.facebook.com/groups/CrackuCATExam/> -

Daily free Live session for CAT <https://cracku.in/cat/classroom>

For free CAT mocks visit <https://cracku.in/cat>