

Scorecard (procview.jsp?sid=aaaN5tjtX0b7WgArBjowySun Jan 08 23:39:44 IST
2023&qsetId=Qkl/zCiPcGE=&qsetName=)

Accuracy (AccSelectGraph.jsp?sid=aaaN5tjtX0b7WgArBjowySun Jan 08 23:39:44 IST
2023&qsetId=Qkl/zCiPcGE=&qsetName=)

Qs Analysis (QsAnalysis.jsp?sid=aaaN5tjtX0b7WgArBjowySun Jan 08 23:39:44 IST
2023&qsetId=Qkl/zCiPcGE=&qsetName=)

Video Attempt / Solution (VideoAnalysis.jsp?sid=aaaN5tjtX0b7WgArBjowySun Jan 08 23:39:44 IST
2023&qsetId=Qkl/zCiPcGE=&qsetName=)

Solutions (Solution.jsp?sid=aaaN5tjtX0b7WgArBjowySun Jan 08 23:39:44 IST
2023&qsetId=Qkl/zCiPcGE=&qsetName=)

Bookmarks (Bookmarks.jsp?sid=aaaN5tjtX0b7WgArBjowySun Jan 08 23:39:44 IST
2023&qsetId=Qkl/zCiPcGE=&qsetName=)

Section-1

Sec 1

Q.1 [11831809]

What is the least sum of a set of 5 distinct prime numbers whose average is also a prime number?

Solution:

Correct Answer : 55

The sequence of the numbers is 3, 5, 7, 11, 29.

Bookmark

FeedBack

 Answer key/Solution

Q.2 [11831809]

The mean and median of Ashok's 7 scores is 74. If his lowest 2 scores are not considered, the mean increased by 1 whereas the median remained the same. What is the highest score he could have got, if all his scores were whole numbers and no number except 74 appears more than once?

1 ☐ 78

2 ☐ 79

3 ☐ 80

4 ☐ 81

Solution:

Correct Answer : 3

For the given conditions the scores could have been in the increasing order A, B, C, 74, 74, D, E. This is because the median scores with 7, 5 were in both cases 74.

So the 4th and 5th scores have to both be 74

We need to maximize E. So A, B, C, D must be as small as possible but satisfying all the conditions.

Let A, B be $74 - x$ and $74 - y$ respectively. Since when these numbers are removed, the average of the remaining numbers is going up by 1, we can say that $x + y$ will be equal to 5.

So the possible values of A, B, C, D are 71, 72, 73, 74 respectively.

In this case $E = 80$. This is the largest value of E that is possible.

Bookmark

FeedBack

 Answer key/Solution

Q.3 [11831809]

A big cube is completely sliced into 64 identical cubes. If 12 litres of paint was used to paint the original cube, how many more litres of paint will be required to paint the smaller cubes on all the faces?

1 ☐ 18

2 ☐ 24

3 ☐ 36

4 ☐ 48

Solution:

Correct Answer : 3

There are six faces in big cube. Each face takes 2 L paint. When it is cut into 64 identical and smaller cubes (by making 3 cuts along each dimension), each cut gives two new faces. So, 9 cut gives 18 new faces. Hence, extra amount of paint required is 36 L.

Bookmark

FeedBack

 Answer key/Solution

Q.4 [11831809]

Anuj can complete a piece of work in 8 days, Bhanu in 12 days and Chetan in 16 days. All of them started the work together but Chetan left when he realised that the remaining work will be completed without him in the next three days, and Bhanu left when he realised that the remaining work will be completed without him in the next two days. Find the total number of days taken to complete the work.

1 ☐ 4

2 ☐ 5

3 ☐ 6

4 ☐ 10

Solution:

Correct Answer : 2

 Answer key/Solution

Let number of units of work = LCM (8, 12, 16) = 48.

So, Anuj can do 6 units per day, Bhanu can do 4 units per day, Chetan can do 3 units per day and they together can do 13 units per day.

Let the required number of days be n .

$$6n + 4(n - 2) + 3(n - 3) = 48$$

$$\Rightarrow 13n - 8 - 9 = 48$$

$$\Rightarrow 13n = 65$$

$$\Rightarrow n = 5 \text{ days.}$$

Bookmark

FeedBack

Q.5 [11831809]

The interest accrued in the first two years and that in the first three years on a sum invested at a certain rate of interest, compounded annually, were Rs. 1,100 and Rs. 1,820 respectively. What was the rate of interest?

1 ☐ 10%

2 ☐ 24%

3 ☐ 30%

4 ☐ 20%

Solution:

Correct Answer : 4

Let the rate of interest be $r\%$.

The interest received in the first year = $\frac{Pr}{100} = k$ (say)

Total interest received in the first two years

$$= k + k\left(1 + \frac{r}{100}\right) = k\left(2 + \frac{r}{100}\right) = 1100 \quad \dots (i)$$

Interest received in the third year

$$= k\left(1 + \frac{r}{100}\right)^2 = 1820 - 1100 = 720 \quad \dots (ii)$$

Dividing (i) by (ii), we get

$$\frac{k\left(2 + \frac{r}{100}\right)}{k\left(1 + \frac{r}{100}\right)^2} = \frac{1100}{720} \Rightarrow \frac{\left(2 + \frac{r}{100}\right)}{\left(1 + \frac{r}{100}\right)^2} = \frac{55}{36} \Rightarrow r = 20\%.$$

Bookmark

FeedBack

Q.6 [11831809]

At the end of the year 2022, Madan had Rs. 500 and Birju had Rs. 50,000. At the end of every year, Madan will have twice as much money as he did at the end of the previous year and Birju will have 60% of the money that he did at the end of the previous year. At the end of which year will Madan have more money than Birju for the first time?

1 ☐ 2025

2 ☐ 2026

3 ☐ 2027

4 ☐ 2028

Solution:

Correct Answer : 2

Let this happen k years after 2022.

Then, $500 \times (2^k) > (3/5)^k \times 50000$

Or, $(10/3)^k > 100$

So $k = 4$.

Hence, this would be the case in the year 2026.

Bookmark

FeedBack

 Answer key/Solution

 Answer key/Solution

Q.7 [11831809]

Let $f(x) = x^2 + ax + b$ be a quadratic, where a and b are integers with $5 \leq a, b \leq 11$. For how many distinct ordered pairs (a, b) will $f(x)$ not have real solutions ?

Solution:

Correct Answer : 7

If the graph does not cut the X-axis, then the roots must be imaginary.

$$a^2 - 4b < 0:$$

If $a = 5$, there are 5 cases: $b = 7$ to $b = 11$

If $a = 6$, there are 2 cases: $b = 10, 11$

Hence, 7 ordered pairs (a, b) are there.

Bookmark

FeedBack

 Answer key/Solution

Q.8 [11831809]

In how many ways can 10 be written as the sum of two or more natural numbers? Consider $(1 + 9)$ to be different from $(9 + 1)$.

1 ☐ 1024

2 ☐ 512

3 ☐ 511

4 ☐ 1023

Solution:

Correct Answer : 3

 Answer key/Solution

The number of solutions of $x_1 + x_2 + x_3 + \dots + x_r = n$, where $x_1, x_2, x_3, \dots, x_r$ are natural numbers, is ${}^{n-1}C_{r-1}$.

According to the question, the value of n is 10 and the values that ' r ' can have are 2, 3, 4, 5, ..., 10.

Hence, the required number of ways = ${}^9C_1 + {}^9C_2 + {}^9C_3 + \dots + {}^9C_9 = 2^9 - {}^9C_0 = 2^9 - 1 = 512 - 1 = 511$.

Bookmark

FeedBack

Q.9 [11831809]

Find the number of solutions of the equation $\log_3 (x + 5) = 6 - x$.

Solution:

Correct Answer : 1

[Answer key/Solution](#)

$$\log_3 (x + 5) = 6 - x$$

$$\Rightarrow x + 5 = 3^{6-x}$$


x will always be an even number.

(Power of 3 is always an even number. In left hand side if x is even, then $(x + 5)$ will be odd)

Possible values of $x = 0, 2, 4, 6, \dots$

Only $x = 4$ satisfies the condition.

Alternative method:




Only one intersection point.

Bookmark

FeedBack

Q.10 [11831809]

In the figure given below, $PS : SR = 1 : 2$, $QL : LS = 1 : 2$ and $LM = MR$. Find the ratio of the area of triangle PQR to that of LMN.



1 ☐ 9 : 1

2 ☐ 6 : 1


3 ☐ 24 : 5

4 ☐ 3 : 1

Solution:

Correct Answer : 2

 Answer key/Solution



It can be observed that $\triangle RMO$ and $\triangle RLS$ are similar.

$$\therefore \frac{RM}{RL} = \frac{RO}{RS} = \frac{MO}{LS} = \frac{1}{2} \quad (\text{As } RM : ML = 1 : 1 \text{ or } RM = ML)$$

$RO : OS = 1 : 1$ and $PS : SR = 1 : 2$.

$$\therefore RO = OS = SP$$

It can also be observed that $\triangle PNS$ and $\triangle PMO$ are also similar.

$$\therefore \frac{PS}{PO} = \frac{NS}{MO} = \frac{1}{2} \quad (\because OS = SP)$$

$$\Rightarrow NS = \frac{MO}{2} \text{ and } MO = \frac{LS}{2} \text{ or } NS = \frac{LS}{4}$$

$$\Rightarrow LN : NS = 3 : 1$$

Let the area of $\triangle PQR = x$ sq. units.

QS divides PR in the ratio 1 : 2 at S.

$$\text{Area of } \triangle QSR = \frac{2}{3}x$$

$$\therefore \text{Area of } \triangle QLR = \frac{1}{3} \left(\frac{2}{3}x \right) = \frac{2}{9}x$$

$$\therefore \text{Area of } \triangle RLS = \frac{2}{3}x - \frac{2}{9}x = \frac{4}{9}x$$

$$\text{Also, the area of } \triangle QLP = \frac{1}{3} \left(\frac{1}{3}x \right) = \frac{1}{9}x$$

$$\therefore \text{Area of } \triangle PLS = \frac{1}{3}x - \frac{1}{9}x = \frac{2}{9}x$$

$$\therefore \text{Area of } \triangle PLN = \frac{3}{4} \left(\frac{2}{9}x \right) = \frac{1}{6}x \quad (\because LN : NS = 3 : 1)$$

$$\therefore \text{Area of } \triangle PLR = \text{Area of } \triangle PLS + \text{Area of } \triangle RLS$$

$$= \frac{4}{9}x + \frac{2}{9}x = \frac{2}{3}x$$

$$\therefore \text{Area of } \triangle PLM = \frac{1}{2} \left(\frac{2}{3}x \right) = \frac{1}{3}x \quad (\because LM = MR)$$

$$\text{Area of } \triangle LNM = \text{Area of } \triangle PLM - \text{Area of } \triangle PLN$$

$$= \frac{1}{3}x - \frac{1}{6}x = \frac{1}{6}x \text{ sq. units}$$

Hence, the required ratio = 6 : 1.

Alternate solution:

If the area of triangle PQR is 'x' sq. units.

Since $PS : SR = 1 : 2$, therefore, area of triangle QSR = $\frac{2x}{3}$ sq. units

Since $QL : LS = 1 : 2$, therefore, area of triangle $LSR = 2x/3 \times 2/3$ sq. units

Since $LM = MR$ and $LN = 3NS$ (Approximately, from the diagram)

So area of triangle LMN will be between $2x/3 \times 2/3 \times 1/4$ and $2x/3 \times 2/3 \times 1/3$

i.e., between $x/9$ and $x/6.75$

Hence, only option (2) i.e., $6 : 1$ satisfies the condition.

Bookmark

FeedBack

Q.11 [11831809]

The number of factors of the number of balls in four boxes – P, Q, R and S – is 16, 28, 40 and 49 respectively. If Ravi wants to pick a box, out of the four boxes, which box should he pick to be sure of picking up of a box in which the number of balls is the cube of a natural number?

1 ☐ P

2 ☐ Q

3 ☐ R

4 ☐ S

Solution:

Correct Answer : 4

The number of factors of a number $(N) = p^a \times q^b \times r^c \dots$, where a, b, c, \dots are natural numbers and p, q, r, \dots are prime numbers, is $(a + 1) \times (b + 1) \times (c + 1)$.

If N is a perfect cube, a, b, c, \dots have to be multiples of 3.

If the number of factors is 16 $\dots(a, b, c, \dots)$, these can be (1, 7) or (3, 3) or (1, 1, 3) and their ordered sets. In all cases we do not see that the number is a perfect cube.

Similarly, for 28 the set should be (1, 13); (3, 6); (1, 1, 6), \dots and their ordered sets.

For 40 it is (1, 19); (3, 9); (4, 7), \dots , and their ordered sets.

For 49 it is (6, 6) only. This is the only number that can be a perfect cube only.

Bookmark

FeedBack

 Answer key/Solution

Q.12 [11831809]

A square shaped floor of dimensions 35×35 is fully covered with square shaped tiles of dimensions 1×1 , 2×2 and 3×3 . The number of tiles used of dimensions 1×1 to cover the floor cannot be less than

1 ☐ 1

2 ☐ 2

3 ☐ 4


4 ☐ 8

Solution:

Correct Answer : 3

[Answer key/Solution](#)

3×3 tiles may be used to cover 33×33 units part of the floor.



The left out part, as shown in the above figure, may be covered by using the 2×2 tiles.

The shaded part shows the part that cannot be filled even by using 2×2 tiles and have to be filled by 1×1 tiles.

Hence, the minimum number of tiles of dimension 1×1 is required is 4.

Bookmark

FeedBack

Q.13 [11831809]

Vishal went from A to B at a certain speed and came back to A from B at a speed that was twice the speed at which he went from A to B. The entire journey took 5 hours. Had Vishal returned at a speed that was 25% more than the speed at which he actually returned, how long would it have taken him for the entire journey?

1 ☐ 4 hours 40 minutes

2 ☐ 4 hours 30 minutes


3 ☐ 5 hours 15 minutes

4 ☐ 3 hours 40 minutes

Solution:

Correct Answer : 1

[Answer key/Solution](#)



Let d be the distance between A and B and x km/h be the speed at which Vishal travels.

Time taken for the onward and return journey = $d/x + d/2x = 5$... (i)

$$\Rightarrow d/x = 10/3$$

When the return speed is increased by 25%, the time taken is = $d/x + 2d/5x = 7d/5x$

$$\Rightarrow 7d/5x = 7/5 \times 10/3 = 14/3 \text{ hours} = 4 \text{ hours } 40 \text{ minutes.}$$

Bookmark

FeedBack

Q.14 [11831809]

A total of ' x ' litres of pure milk was drawn from a vessel containing 256 litres of pure milk and was replaced with water. The resulting mixture was thoroughly mixed to obtain a homogenous solution and then again ' x ' litres of solution was drawn, and replaced with water. The resulting mixture was thoroughly mixed. This procedure was performed four times and the vessel now contained 81 litres of pure milk. Find the value of ' x '.

Solution:

Correct Answer : 64

[Answer key/Solution](#)

$$\text{Final quantity of the component} = \text{Initial quantity of the component} \left(1 - \frac{\text{replaced quantity in each turn}}{\text{total quantity of the mixture}} \right)^n$$

$$\Rightarrow 81 = 256 \left(1 - \frac{x}{256} \right)^4$$

$$\Rightarrow \left(\frac{81}{256} \right)^{\frac{1}{4}} = 1 - \frac{x}{256}$$

$$\Rightarrow 1 - \frac{x}{256} = \frac{3}{4}$$

$$\Rightarrow x = 64.$$

Bookmark

FeedBack

Q.15 [11831809]

For non-negative real numbers x and y , $3x + 4y \leq 16$ and $5x + 2y \leq 10$. What is the maximum possible value of $(4x + 10y)$?

1 ☐ 40

2 ☐ 32

3 ☐ 48

4 ☐ 52

Solution:

Correct Answer : 1

If we want to maximise $4x + 10y$, y should be maximum as coefficient of y is 10.
From given inequalities maximum value of y is 4 and then the value of x will be zero.

So maximum of $4x + 10y = 4(0) + 10(4) = 40$.

Bookmark

FeedBack

 Answer key/Solution

Q.16 [11831809]

Find the value of the following expression:

$$\log_{0.001} \left[\frac{1}{2} \right] + \log_{0.001} \left[\frac{2}{3} \right] + \log_{0.001} \left[\frac{3}{4} \right] + \dots + \log_{0.001} \left[\frac{99}{100} \right].$$

1 ☐ $-3/2$

2 ☐ $2/3$

3 ☐ $3/2$

4 ☐ $-2/3$

Solution:

Correct Answer : 2

 Answer key/Solution

$$\log_{(0.001)} \left[\frac{1}{2} \right] + \log_{(0.001)} \left[\frac{2}{3} \right] + \log_{(0.001)} \left[\frac{3}{4} \right] + \dots + \log_{(0.001)} \left[\frac{99}{100} \right]$$

$$= \log_{(0.001)} \left[\frac{1}{2} \right] \times \left[\frac{2}{3} \right] \times \left[\frac{3}{4} \right] \times \dots \times \left[\frac{99}{100} \right]$$

$$= \log_{(0.001)} \left[\frac{1}{100} \right] = \log_{(0.1)^3} \left[\frac{1}{10} \right]^2 = \frac{2}{3}.$$

Bookmark

FeedBack

Q.17 [11831809]

Ramu gave away return gifts on the occasion of his daughter's birthday. Every gift was a pack of 1 Toy, 2 Chocolates and 1 Book. The Toys were available in boxes of 30 Toys each, the Chocolates in boxes of 40 Chocolates each and the Books came in boxes of 15 books each. Ramu made no incomplete gift and there were no unused Toys, Chocolates or Books. What is the minimum number of boxes purchased for the party?

Solution:

Correct Answer : 9

There must be N number of Toys, 2N of Chocolates and N of Books.

LCM of (30, 60, 30) is 60. So the number of Toys must be 60, the number of

Chocolates must be 120 and the number of Books must be 60. This will make sure that no box is left unused.

And there are no incomplete gifts.

So the minimum number of boxes is $2 + 3 + 4 = 9$.

Bookmark

FeedBack

[Answer key/Solution](#)

Q.18 [11831809]

If P is a point on $x^2 + y^2 + 18 = 2(5x + 3y)$ and Q is a point on $x^2 + y^2 + 240 = 2(16x + 3y)$, the distance (in units) between P and Q cannot be more than

1 ☐ 18

2 ☐ 20

3 ☐ 21

4 ☐ 23

Solution:

Correct Answer : 2

[Answer key/Solution](#)


The given equations can be rewritten as:

$$(x - 5)^2 + (y - 3)^2 = 4^2 \quad \dots (i)$$

$$(x - 16)^2 + (y - 3)^2 = 5^2 \quad \dots (ii)$$

From the above equations, it can be noted that the given equations represent circles.

The following graph shows the positions of the two circles.



From the graph, it can be noted that the maximum distance between any two points on the circle, with one point on each, cannot be more than AF i.e., 20 units.

Bookmark

FeedBack

Q.19 [11831809]

Ramlal, a fruit vendor, sells only apples and oranges. One day in the morning, he bought the two types of fruits in the ratio 3 : 5 respectively. Later on, he found that 8% of the apples and 12% of the oranges were rotten. At the end of the day, he was left with 24 fruits, which accounted for 4% of the total number of fruits that he had bought in the morning. On the given day, if Ramlal sold only those fresh fruits that he had bought in the morning and disposed of all the rotten fruits during the course of the day, the number of fruits sold by Ramlal was

Solution:

Correct Answer : 513

 Answer key/Solution

Let the total number of fruit bought by Ramlal be 'x'.

$\therefore 4\% \text{ of } x = 24 \Rightarrow x = 600$

Number of apples bought by Ramlal $= \frac{3}{8} \times 600 = 225$

Number of pineapples bought by Ramlal $= \frac{5}{8} \times 600 = 375$

Hence, the required number of fruits $= 600 - (8\% \text{ of } 225 + 12\% \text{ of } 375 + 24) = 513$.

Bookmark

FeedBack

Q.20 [11831809]

If $f(x) = |x - 4|$ and $g(x) = x^2$, find the number of values of x for which $f(g(x)) = g(f(x))$.

Solution:

Correct Answer : 1

 Answer key/Solution

$f(g(x)) = |x^2 - 4|$; $g(f(x)) = |x - 4|^2$

Both sides have to be positive because $|x - 4|^2$ will always be positive.

Therefore, solution of $f(g(x)) = g(f(x))$ is same as solution of $x^2 - 4 = x^2 - 8x + 16$; $x = 2.5$ (this is consistent with our original assumption)

Hence, the number of values of x for which $f(g(x)) = g(f(x))$ is 1, that is., $x = 2.5$.

Bookmark

FeedBack

Q.21 [11831809]

The percentage volumes of liquid 'A' in three solutions – I, II and III – form a geometric progression in that order. If we mix I, II and III solutions in the ratio 2 : 3 : 4, we obtain a solution containing 32% of A. If we mix them in the ratio 3 : 2 : 1, we obtain a solution containing 22% of A. What is the percentage of liquid A in II?

1 ☐ 18%

2 ☐ 36%

3 ☐ 24%

4 ☐ 48%

Solution:

Correct Answer : 3

[🔍 Answer key/Solution](#)

Let percentage of liquid A in I, II and III vessels be a , ar and ar^2 , respectively.

$$\therefore \frac{2a + 3ar + 4ar^2}{900} = \frac{32}{100}$$

$$\text{and } \frac{3a + 2ar + ar^2}{600} = \frac{22}{100}$$

$$2a + 3ar + 4ar^2 = 288 \quad \dots (i)$$

$$3a + 2ar + ar^2 = 132 \quad \dots (ii)$$

Dividing (i) by (ii), we get

$$132r^2 + 99r + 66 = 72r^2 + 144r + 216$$

$$\Rightarrow (4r + 5) \times (r - 2) = 0$$

$$\Rightarrow r = 2 \text{ (As } r \text{ is an integer)}$$

$$\therefore a = 12.$$

Hence, the percentage of liquid A in II is 24%.

Bookmark

FeedBack

Q.22 [11831809]

The marks scored by Pankaj in three subjects are in the ratio 4 : 5 : 6. Pankaj scored an overall aggregate of 62.5% in the exam. If the maximum marks in each subject are in the ratio 3 : 4 : 5, in how many subjects did Pankaj score more than 65% marks?

Solution:

Correct Answer : 1

[🔍 Answer key/Solution](#)

Let the maximum marks in each subject be $3k$, $4k$ and $5k$.

$$\therefore \text{Pankaj's total score} = 62.5\% \text{ of } (12k) = 7.5k$$

Let Pankaj have scored $4x$, $5x$ and $6x$ in the three subjects.

$$\therefore 4x + 5x + 6x = 15x = 7.5k \Rightarrow x/k = 1/2$$

If maximum marks in each subject is 60, 80 and 100, then his score in the respective subjects are 40, 50 and 60.

We know that $40/60 = 66.67\%$ whereas the other two are less than 65%.

Hence, he has scored more than 65% in exactly one subject.

Bookmark

FeedBack