



45 mm	85 mm  FeXentrix 60 Fexofenadine Hydrochloride Tablets USP 60mg Antihistamin Allergy Relief 100 Film-Coated Tablets For Hay Fever & Allergy Rhinitis 36 mm	36 mm  FeXentrix 60
	Composition : Each film coated tablet contains: Fexofenadine Hydrochloride USP 60 mg Excipients Q.S. Colour: Titanium Dioxide BP Dosage : As directed by the Physician. Storage : Store at a temperature not exceeding 30°C and protect from light & moisture. KEEP THE MEDICINE OUT OF REACH OF CHILDREN. Swallow the tablet with a drink of water. Do not chew or crush the tablets Caution: To be sold by retail on the prescription of a Registered Medical Practitioner only.  8 906168 426718	
45 mm  FeXentrix 60 36 mm	 FeXentrix 60 Fexofenadine Hydrochloride Tablets USP 60mg Antihistamin Allergy Relief 100 Film-Coated Tablets For Hay Fever & Allergy Rhinitis Manufactured for :  WHITE LOTUS HEALTHCARE (WHO-GMP & ISO CERTIFIED COMPANY) Surat, Gujarat, India. Contact to: +91 94290 20306 N. Code No. : GUJ/DRUGS/G/25/2631 Batch No. : Mfg. Date : Exp. Date :	



210 mm

148.00 mm

Fexentrix 60

Fexofenadine Hydrochloride Tablets USP 60mg

Composition:

Each film coated tablet contains:
Fexofenadine Hydrochloride USP 60 mg
Excipients Q.S.
Colour: Titanium Dioxide BP

Read this entire leaflet carefully before you start taking this medicine, because it contains important information for you.

- Keep this leaflet. You may need to read it again.
- If you have any further questions, ask your doctor or pharmacist.
- This medicine has been prescribed for you only. Do not pass it on to others. It may harm them, even if their signs of illness are the same as yours.
- If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet.

What is in this leaflet

1. What FEXENTRIX 60 is and what it is used for
2. What you need to know before you take FEXENTRIX 60
3. How to take FEXENTRIX 60
4. Possible side effects
5. How to store
6. Contents of the pack and other information

1. What FEXENTRIX 60 is and what it is used for

FEXENTRIX 60 contains fexofenadine hydrochloride, which is an antihistamine. Fexofenadine Hydrochloride Tablets is used in adults and adolescents of 12 years and older to relieve the symptoms that occur with long term allergic skin reactions (chronic idiopathic urticaria) such as itching, swelling and rashes.

2. What you need to know before you take FEXENTRIX 60

Do not take FEXENTRIX 60 if:

If you are allergic to fexofenadine or any of the other ingredients of this medicine.

Warnings and precautions

Talk to your doctor or pharmacist before taking Fexofenadine Hydrochloride Tablets if

- You have problems with your liver or kidneys
- You have or ever had heart disease, since this kind of medicine may lead to a fast or irregular heartbeat
- You are elderly.

If any of these apply to you, or if you are not sure, tell your doctor before taking Fexofenadine Hydrochloride Tablets.

Other medicines and Fexofenadine Hydrochloride Tablets

Tell your doctor or pharmacist if you are taking, have recently taken or might take any other medicines.

If you are taking apalutamide (a medicine to treat prostate cancer), as the effect of fexofenadine may be decreased.

Indigestion remedies containing aluminium and magnesium may affect the action of Fexofenadine Hydrochloride Tablets by lowering the amount of medicinal product absorbed.

It is recommended that you leave about 2 hours between the time that you take Fexofenadine Hydrochloride Tablets and your indigestion remedy.

Pregnancy and breast-feeding

Ask your doctor or pharmacist for advice before taking any medicine.

Do not take Fexofenadine Hydrochloride Tablets if you are pregnant, unless necessary.

Fexofenadine Hydrochloride Tablets is not recommended during breast-feeding.

Driving and using machines

Fexofenadine Hydrochloride Tablets is unlikely to affect your ability to drive or operate machinery. However, you should check that these tablets do not make you feel sleepy or dizzy before driving or operating machinery.

3. How to take FEXENTRIX 60

Always take this medicine exactly as your doctor has told you. Check with your doctor or pharmacist if you are not sure.

For adults and children aged 12 years and over

Seasonal Allergic Rhinitis Adults and Children 12 Years and Older

The recommended dose of Fexofenadine Hydrochloride Tablets is 60 mg twice daily, or 180 mg once daily with water. A dose of 60 mg once daily is recommended as the starting dose in patients with decreased renal function.

Children 6 to 11 Years

The recommended dose of Fexofenadine Hydrochloride Tablets is 30 mg twice daily with water. A dose of 30 mg once daily is recommended as the starting dose in paediatric patients with decreased renal function.

Chronic Idiopathic Urticaria Adults and Children 12 Years and Older The recommended dose of Fexofenadine Hydrochloride Tablets is 60 mg twice daily or 180 mg once daily with water. A dose of 60 mg once daily is recommended as the starting dose in patients with decreased renal function.

Children 6 to 11 Years The recommended dose of Fexofenadine Hydrochloride Tablets is 30 mg

twice daily with water. A dose of 30 mg once daily is recommended as the starting dose in pediatric patients with decreased renal function.

If you take more Fexofenadine Hydrochloride Tablets than you should

If you take too many tablets, contact your doctor or the nearest hospital emergency department immediately.

Symptoms of an overdose in adults are dizziness, drowsiness, fatigue and dry mouth.

If you forget to take Fexofenadine Hydrochloride Tablets

Do not take a double dose to make up for a forgotten tablet.

Take the next dose at the usual time as prescribed by your doctor.

If you stop taking Fexofenadine Hydrochloride Tablets

Tell your doctor if you want to stop taking Fexofenadine Hydrochloride Tablets before you have finished your course of treatment.

If you stop taking Fexofenadine Hydrochloride Tablets earlier than planned, your symptoms may return.

If you have any further questions on the use of this medicine, ask your doctor or pharmacist.

4. Possible side effects

Like all medicines, this medicine can cause side effects, although not everybody gets them.

Tell your doctor immediately and stop taking Fexofenadine Hydrochloride Tablets if you experience:

swelling of the face, lips, tongue or throat and difficulty breathing, as these may be signs of a serious allergic reaction.

Common side effects (may affect up to 1 in 10 people):

- headache
- drowsiness
- Feeling sick (nausea)
- Dizziness.

Uncommon side effects (may affect up to 1 in 100 people):

- Tiredness
- Sleepiness.

Additional side effects (frequency not known: cannot be estimated from the available data) which may occur are:

- Difficulty sleeping (insomnia)
- Sleeping disorders
- Bad dreams
- Nervousness
- Fast or irregular heartbeat
- Diarrhoea
- Skin rash and itching
- Hives
- Blurred vision
- Serious allergic reactions which can cause swelling of the face, lips, tongue or throat, flushing, chest tightness, and difficulty breathing.

Reporting of side effects

If you get any side effects, talk to your doctor. This includes any possible side effects not listed in this leaflet. You can also report side effects directly.

5. How to store

Keep this medicine out of the sight and reach of children.

Do not use this medicine after the expiry date which is stated on the label and blister pack after EXP. Store at a temperature not exceeding 30°C and protect from light & moisture.

Do not throw away medicines via wastewater or household waste. Ask your pharmacist how to throw away medicines you no longer use. These measures will help protect the environment.

6. Contents of the pack and other information

What FEXENTRIX 60 contains

- The active substance is fexofenadine hydrochloride.

Packing:

10 tablets packed in blister, such 10 blister packed into carton along with insert.

Manufactured for :


WHITE LOTUS
HEALTHCARE
(WHO-GMP & ISO CERTIFIED COMPANY)
Surat, Gujarat, India. Contact to: +91 94290 20306

