BISO
JICA



PROCESS FOR CULTIVATING ONIONS AND VEGETABLES

FOR A 1-HECTARE PLOT

BISO JICA

INGREDIENTS FOR PREPARATION

DAICHI NO TOMO 500

Enzyme inhibition organic fertilizer

DAICHI NO TOMO

Enzymatic restoration arable land

3000 METALS

Enzyme for flower bud differentiation

SP 10-15-20

Organic enzymes

1000 METALS

Enzymatic therapy

STEPS TO EXECUTE

Phase 1: Soil Preparation and Seed Treatment

Seed Soaking

Daichi No Tomo	-10ml	=> Dissolve in 10-20 liters of purified water and immerse for 12-24 hours.
Daichi No Tomo500	-100ml	
Mitaka 1000	- 10ml	

Prepare the terrain.

- Select loose, well-drained soil with a pH level of 5.5 to 6.5 and access to a water source.
- Apply soil spray:
  Daichi No Tomo - 1L
  Biso antagonistic fungus - 2kg. => For every hectare
- Plow the field and allow the soil to dry for 7 to 10 days, then construct raised
- beds that are 0.8 to 1.2 meters wide. Apply the spray uniformly across the field:

Daichi No Tomo Biso - 1L
Antagonistic Mushroom - 2 kilograms => Dissolve in clean water at a concentration of 2,000 L/ha.

- Apply a foundational fertilizer composed of vermicompost, manure, and well-decomposed organic matter (composted in accordance with the Biso process).

BISO
JICA

Planting:

- Select disease-free seeds and ensure the field is not waterlogged. Water and apply spray uniformly across the field:
-

Daichi No Tomo - 500ml
Daichi No Tomo 500 - 3L
1000 bags - 500ml

Dissolve in clean water at a concentration of 2,000 L/ha.

- Following planting, water gently to maintain soil moisture, and offer shade if the sunlight is excessively intense.

BISO
JICA

Phase 2: Growth Stage Management

- Utilize Daichi No Tomo 500 + SP 10-15-20 + Mitaka 3000
- Mixing ratio:

Daichi No Tomo 500 - 2 Liters
SP 10-15-20 - 500ml
Meters 3000 - 500ml

Dissolve in 1000 L of purified water - once every five days.

- Utilize Daichi No Tomo + Mitaka 1000.
- Mixing ratio:

Daichi No Tomo - 500ml
Mitaka 1000 - 700ml

Dissolve in 1000 liters of clean water once every 15 days.