

BISO JICA

RICE CULTIVATION PROCESS

BISO
JICA

INGREDIENTS TO PREPARE

Biso organic enzyme:

- Daichi No Tomo
- Daichi No Tomo 500
- Mikata 1000
- NPK 10-15-20
- Mitaka 3000

BISO
JICA

STEPS TO EXECUTE

Phase 1: Preparing straw and soil prior to sowing

- Soak the seeds in Daichi No Tomo 500 at a ratio of 2 parts cold water to 1 part hot water, maintaining a temperature of 30-35 degrees Celsius for 24 hours, followed by incubation prior to sowing.
- Activate 2 kg of Biso microbial inoculum with 500 ml of Daichi No Tomo, dissolving it in 100-200 liters of water, and allow it to activate for 6 hours. Reduce the water level in the field by 5 cm.
- Evenly spray the microbial inoculum onto the straw and rice roots in the afternoon. After 5-7 days, plow the land and flood the field with water.
- Soak the field for 7 to 15 days, depending on the extent of straw decomposition in each field, and then drain the water.

Phase 1: Pre-sowing of rice

- Combine 6 liters of Daichi 500 with 1 liter of Daichi No Tomo and mix with 100 liters of water until the desired volume is achieved, depending on the intended application.
- Spray directly onto the field prior to sowing.
- When applying, incorporate additional water to ensure the appropriate dosage and proceed with immediate application in the field.
- Subsequently, sow the seeds.

Stage 3: Rice aged 7-10 days

- Combine SP 10-15-20 with 100 liters of water, adding sufficient water to fully immerse the ingredients. Incorporate Mitaka 1000 and stir until completely dissolved. Apply in the early morning.

Stage 4: Rice aged 17-20 days

- Daichi No Tomo 500 + Daichi No Tomo with 100 liters of water, or adjust the water quantity as necessary. Stir thoroughly until fully dissolved. Apply in the early morning.

Stage 5: Rice aged 27-30 days

- Mix SP 10-15-20+ Mitaka 1000 with 100 liters of water, stir thoroughly, and apply as a spray in the early morning or after 4 PM in the afternoon.

Stage 6: Rice in the process of flowering or already in bloom (37-40 days)

- Combine 15 liters of Mitaka 3000 with 100 to 1,000 liters of water, mix thoroughly, and apply as a spray in the early morning.

Stage 7: Rice 47-50 days

- Combine SP 10-15-20 with 100 liters of water, adjusting the water quantity as
- necessary. Incorporate Mitaka 1000, stirring thoroughly until fully dissolved, and apply in the early morning.

Stage 8: Rice 57-60 days

- Combine 15 liters of Mitaka 3000 with 100 to 1,000 liters of water, mix thoroughly, and apply as a spray in the early morning.

Stage 9: Rice, 70 to 75 days

- Combine 1 liter of Mitaka 1000 with 100 to 1,000 liters of water, mix thoroughly, and apply as a spray in the early morning.

BISO
JICA

JOIN THE COMMUNITY

This rice cultivation process is generally applicable. However, depending on the specific conditions of each field and the growth state of the rice plants, the dosage must be adjusted flexibly. The products may be mixed for simultaneous use, if necessary.