BISO
JICA

# PROCEDURE FOR APPLYING BISO ENZYME ON YELLOW APRICOT TREES OLDER THAN ONE YEAR

BISO
JICA

Reducing chemical pesticides fosters a transition to safe organic farming, safeguarding the health of growers while promoting the natural and healthy growth of plants. Concurrently, it encourages the uniform and balanced development of shoots, leaves, and flowers, thereby enhancing resistance to fungal diseases and adverse weather conditions. It is advisable to integrate this approach with composting based on the Japanese method (Biso Jica) to reduce costs and enhance efficiency for farmers.



# INGREDIENTS FOR PREPARATION

Biso organic enzyme

- Daichi No Tomo
- Daichi No Tomo 500
- Mikata 1000
- SP 10-15-20
- Mikata 3000

BISO
JICA

# STEPS TO ADHERE TO

## Phase 1: Preparing the fermentation starter post-Tet - Revitalizing and enhancing the canopy (January - March in the Lunar Calendar)

- Objective: To assist the plant in its recovery following flowering and sprouting.
  The shoots are robust and emerging.
- Daichi No Tomo + Daichi No Tomo 500 Root Watering Solution.
- Mikata 1000 spray can be enhanced with Daichi No. Tomo (for the treatment of fungal diseases) on foliage.
- Frequency: Every ten days (first month) following dilation. Gradually, every 20 to 30 days.
- Combination: Incorporate SP 10-15-20 into the watering session. Monday or Tuesday.

This product effectively eliminates and inhibits nematodes. During hot weather, plants require supplementary potassium.

## Phase 2: Nurturing Branches - Cultivating a Full Canopy (April – June in the Lunar Calendar)

- Objective: To improve nutrient delivery to the branches and foliage while restricting premature flowering.
- Apply Daichi No Tomo and Daichi No Tomo 500 to the roots → Adjust Daichi No Tomo 500 based on the plant's condition.
- Chemical fertilizers (NPK 16.16.8 or 30.10.13) may be blended according to the plant's condition to enhance growth (dissolve them in conjunction with Daichi No Tomo 500 to convert them into a form that is readily absorbable by the plant) → transforming raw food into cooked food.
  Mikata 1000 foliar spray combined with Daichi No
- Tomo.
- Frequency: Every 15 to 20 days.

Limit phosphorus and potassium consumption during this phase to avert premature flower bud differentiation; effective water management is the most critical factor during this time, particularly to prevent water stress.

BISO
JICA

## Phase 3: Flower Bud Formation - Encouraging Flower Bud Differentiation (July - September in the Lunar Calendar)

- Objective: To promote bud differentiation in plants. The flowers exhibit uniformity and vitality.
- Water or apply a spray of organic NPK 10-15-20 enzyme + Daichi No Tomo, in conjunction with foliar application. 1000 meters

Frequency: Every 20 to 30 days.

Root watering: Utilize Biso enzyme to enrich the roots and facilitate nutrient transport. Trim branches and modify watering practices in accordance with the specific garden requirements.

## Phase 4: Bud Nurturing - Preparation for Flowering (October – December in the Lunar Calendar)

- Objective: To foster the development of large, uniform, and healthy buds, ensuring they are adequately prepared for flowering in time for Tet (Vietnamese Lunar New Year).
- Watering with Daichi No Tomo - Daichi No Tomo 500: modify the quantity of SP 10-15-20 based on weather conditions and nutrient levels.
  Frequency: Every 20 to 30 days.
-

Limit protein consumption.

## Phase 5: Inducing flowering – ensuring that blooms occur in time for Tet (Vietnamese Lunar New Year).

- Objective: To offer nutritional support that aids plants in boosting bud and flower production, enhancing color, and prolonging the lifespan of flowers, ensuring timely blooming for Tet (Vietnamese Lunar New Year) with the use of Mikata 3000.
- 20-30 days prior to the Lunar New Year (leaf-picking stage): Irrigate with Mikata 3000 (flower bud differentiation) every 10 days → crucial.

BISO
JICA

## JOIN THE COMMUNITY

Mikata 1000 can be integrated with bio-based products (seasoning oil, nano copper, nano silver, abamectin) to enhance its efficacy in pest management.

The process is contingent upon individual experience, regional weather conditions, and the techniques employed by each gardener. It is crucial to consult with reputable gardeners to adapt to the current circumstances and determine the appropriate volume of water for irrigation.

***THE ABOVE PROCEDURE IS APPLICABLE ONLY TO CENTRAL AND SOUTHERN VIETNAM.**