POTATO AND SWEET POTATO GROWING PROCEDURE AND VEGETABLES GROWN IN SOIL

Phase 1:

Soil preparation & seed selection: seed treatment and soil preparation are crucial.

• Soak the seeds together with Daichi no Tomo + Daichi no Tomo 500 + Mikata 1000 (soaking time) 6 to 12 hours

• Choose loose, well-drained soil with a good water source and pH level between 5.5 and 6.5. Before plowing, spray with

Daichi No Tomo and Biso antagonistic fungus (1 liter of Daichi No Tomo and 2 kg of antagonistic fungus per hectare).

Plow and let the soil dry for 7-10 days, then create raised beds 0.8-1.2 m wide. Spray with

Lai Daichi No Tomo + 2 kg of antagonistic fungi before planting.

Note: Dissolve Daichi No Tomo together with the antagonistic fungus Biso. Mix with 2,000 – 3,000 liters.

Water, spray evenly over the field.

• Apply base fertilizer using:

• Vermicompost, animal manure, and decomposed organic matter (composted using the Biso process): to reduce costs.

* Planting the seeds.

* combined with daichi no tomo 500

• Enzyme Daichi No Tomo 500: 3 lít + Mikata 1000: 500 ml + Daichi No Tomo: 500 ml.

(Spray or water evenly over the field) to activate beneficial microorganisms in the soil.

• Choose disease-free seeds that are not damaged by waterlogging.

• Mix with 2,000 – 3,000 liters of water.

• After planting, water lightly to keep the soil moist, and provide shade if the sun is too strong.

Note: They can be mixed together for spraying.

3. Growth stage care: Regularly every 3-5 days.

• Regularly apply Daichi no tomo500 + Daichi no tomo enzyme solution.

• Spraying Mikata 1000 + Daichi No Tomo helps prevent pests and fungal

diseases; 10-15-20 organic fertilizer can be added.

• Weeding – lightly tilling twice:

• First dose: after 10–15 days.

• Second application: after 30 days, combined with hilling around the base

of the plant.

• First top dressing (after 15–20 days): depending on the type of crop.

• Second topdressing (after 35–40 days):

• Supplement with organic potassium (banana fertilizer, organic K).

Pest and disease control – 100% biological

• Use biological products:

• Control leaf-eating insects and stem borers.

• Neem oil: spray regularly to prevent insect infestations.

• Spray Mikata 3000 (for flowering or fruit setting), for example: strawberries, tomatoes, and various flowers.

---

5. Harvesting

• Sweet potatoes: 90–120 days after planting.

• Potatoes: 80–100 days after planting.

• 15–20 days before harvest:

• Stop all fertilizers and microbial products – only water the plants.

• Clean up organic waste and weeds.

• Dig by hand or machine, avoiding crushing the bark.

II. BENEFITS OF USING MICROBIAL ENZYMES IN THE PROCESS OF STRONG ROOT DEVELOPMENT