BISO JICA JAPAN - VIETNAM CO., LTD.
Address: No. 16, D4 Street, Cat Lai Ward, Thu Duc City, Ho Chi Minh City, Vietnam. Phone: 0928899939

BASIC STANDARD TCCS 02:2024/BISOJICA BIOLOGICAL PRODUCT PRODUCT NAME: DAICHI NO TOMO

PREFACE

Standard publisher: BISO JICA JAPAN - VIETNAM CO., LTD. Address: No. 16, D4 Street, Cat Lai Ward, Thu Duc City, Ho Chi Minh City, Vietnam. Phone: 0928899939. The company developed the TCCS 02:2024/BISOJICA standard for the product: DAICHI NO TOMO.
This standard is published together with Decision No.: 01/QD/BISOJICA

1. Scope of application

- This standard specifies requirements for quality, labeling, instructions for use, storage, and transportation. Applicable to the product: DAICHI NO TOMO as announced by the Company.

2. Document

2.1 References

- Law on Product Quality No. 05/2007/QH12 dated November 21, 2007, of the 12th National Assembly.
- Law on Technical Standards and Regulations No. 68/2006/QH11 dated June 29, 2006, of the 11th National Assembly.
- Circular No. 21/2007.TT-BKHCN issued on September 28, 2007, providing guidance on the development and application of standards.

2.2 Referenced Documents

3. Technical Requirements

3.1 Sensory Requirements

- The sensory evaluation criteria for the product are specified in Table 1.

Table 1. Sensory evaluation criteria for the product.

STT	*Indicator name*	*Liquid form*	*Test method*
1	External shape	***required.***	Sensory perception
2	Color	Light yellowish-brown to dark yellowish-brown	
3	Smell	It has the product's characteristic smell.	
4	Sharp-edged foreign object	Do not have	

3.2 Quality Requirements

- The product quality specifications are listed in Table 2.

Table 2. Product quality indicators

STT	*Target*	*Unit of measurement*	*Quality level*	*Test method*
1	Energy	Kcal/100g	53,6	
2	Protein (Not less than)	g/100g	0,4	
3	Lipid (Not less than)	g/100g	2,8	

STT	Target	Unitof measurement:	Quality level 6.7	Testing methodology
4	Carbohydrate (Minimum)	g per 100g,	0,3	
5	Ash (Maximum)	g per 100g,	56	
6	Sodium (Minimum)	mg per 100g,	6.000	
7	Protease (Minimum)	Unit per g,	3.000	
8	Lipase (Minimum)	Unit per g,	89,8	
9	Water	g per 100g,	0,1	
10	Sodium equivalent	g per 100g		

3.3 Hygiene and Safety Requirements

- The product safety standards are outlined in Table 3.

Table 3. Standards for Product Safety

STT	Target	Unit of measurement	Quality standard	Testing methodology
Microorganism				
1	E. coli (maximum)	CFU/g	1.1 x 10^3	
2	Salmonella (maximum)	CFU/25g	None	
Heavy metals				
3	Asen (As) (maximum)	mg/kg	10	
4	Cadmium (Cd) (maximum)	mg/kg	5	
5	Lead (Pb) (maximum)	mg/kg	200	
6	Mercury (Hg) (maximum)	mg/kg	2	

4. Ingredients

- Ingredients consist of: Water and enzymatic components derived from plants and animals in their natural habitats.

5. Packaging, labeling, storage, and transportation 5.1 Packaging

- The product is packaged in plastic bottles and containers.
- Packaging specifications: 1L, 2L, 5L, 10L, 20L (or according to customer specifications).

5.2 Labeling

- Product labeling is conducted in compliance with Government Decree 43/2017/ND-CP, dated April 14, 2017, regarding product labeling, as well as Decree 111/2021/ND-CP, which amends and supplements Decree 43/2017/ND-CP.

5.3 Storage

- The product is kept in a clean, cool, and well-ventilated warehouse, shielded from prolonged exposure to direct sunlight.

5.4 Expiration Date

- Two years from the date of manufacture.

5.5 Transportation

- Vehicles utilized for the transportation of products must be clean, dry, covered, devoid of hazardous substances, and free from pathogens that may lead to cross-contamination of the products.

6. Usage Instructions

- For soil and irrigation, dilute 150ml or 250ml per 1,000m² of soil with a minimum of 500 times that volume of water, applying it to the base of the plant to ensure complete contact with the roots. Water the base of the plant sufficiently to reach the roots.
- When the irrigation pump operates for 10 minutes in a 1,000m2 greenhouse, water the plant roots after 5 minutes (50%) of the irrigation duration has passed. During the 10 minutes that the irrigation pump is running in the 1,000m2 greenhouse, dissolve this product in water and apply it using a chemical mixer after the remaining 5 minutes (50%) of the irrigation duration has passed.
  This product does not qualify as a fertilizer.
  The standard frequency is once or twice monthly.
- It can be applied as a mist; when spraying on leaves, ensure that both the upper and lower surfaces of the leaf, as well as the stem and roots, are uniformly coated.
- Refrain from utilizing hot water sprayers at elevated temperatures.
- ***Kindly consult the product label or the technical documentation provided with the product for further details.***

Ho Chi Minh City, March 16, 2024