

Number Theory

Vinesh Ramgi

June 15, 2018

Abstract

What did the number theorist say as he drowned?

Log, log, log, log....

For an up to date version of this pdf, check my GitHub :)

<https://github.com/vrvinny/number-theory>

Contents

1	Introduction/Review	4
1.1	Introduction	4
1.2	Review	4
1.2.1	Congruences	4
1.2.2	Solving Linear Congruences	5
1.3	Chinese Remainder Theorem	6
1.4	Prime numbers	8
1.5	Fermat's Little Theorem	8
1.5.1	General method to solve $x^a \equiv b \pmod{p}$	9
1.6	Fundamental Theorem of Arithmetic	9
1.6.1	Euclid's Lemma	9
1.6.2	Checking whether a number is prime	10
2	Elementary Number Theory	11
2.1	Euler Totient Function	11
2.2	Euler's Theorem	13
2.2.1	Solving equations of the form $x^a \equiv b \pmod{n}$	14
2.3	Primitive roots	15
2.4	Roots of unity and Cyclotomic Polynomials	16
2.4.1	How to calculate $\Phi_n(x)$	17
2.4.2	Gauss' Theorem	18
2.5	Quadratic reciprocity (Quadratic equations modulo prime numbers)	19
2.5.1	Quadratic Reciprocity Law	20
2.5.2	First Nebensatz	20
2.5.3	Second Nebensatz	21
2.6	Uniqueness Lemma	23
2.6.1	General Uniqueness Lemma	23
2.6.2	Uniqueness Lemma for $\mathbb{Z}[\zeta_p]$	27
3	P-adic Number theory	29
3.1	Hensel's Lemma	30
3.2	Quadratic Congruences	33
3.3	P-adic congruence	35
3.4	Power Series Trick	37
3.4.1	P-adic log & exp	40
3.5	Teichmüller Lifts	43
3.6	Fractional Powers	48
3.7	P-adic integers	49
4	Quadratic rings	52
4.0.1	Properties of conjugates	53
4.0.2	Formula for norms	54
4.0.3	Properties of norms	55

1 Introduction/Review

1.1 Introduction

Number Theory is the theory of the ring \mathbb{Z} and other related rings. A ring (in this course) is a set R with two binary operations $+$ and $*$ such that:

- $(R, +)$ is an abelian group
- $*$ is associative, commutative and has an identity element 1
- $x(y + z) = xy + xz \quad \forall x, y, z \in R$

Examples of rings:

- \mathbb{Z} is a ring
- Every field is a ring, (e.g. $\mathbb{R}, \mathbb{C}, \mathbb{Q}$)
- \mathbb{Z}/n \mathbb{Z} modulo $n = \{0, \dots, n-1\}$
- $\mathbb{F}[X] = \{ \text{polynomials } f(x) \text{ with coefficients in } \mathbb{F} \}$

1.2 Review

1.2.1 Congruences

Let n be a positive integer. Given $x, y \in \mathbb{Z}$, we say x is congruent to y modulo n if $x - y$ is a multiple of n .

$$x \equiv y(n) \quad \text{or} \quad x \equiv y \pmod{n}$$

E.g
$$\begin{aligned} 2 &\equiv 12 \pmod{10} \\ &\equiv -8 \pmod{10} \end{aligned}$$

We write \mathbb{Z}/n for the ring of congruency classes modulo n , i.e. the elements are integer, with two of them regarded as the same if they are congruent modulo n .

Since every integer is congruent to a unique integer in the set $\{0, \dots, n-1\}$, we have $\mathbb{Z}/n = \{0, \dots, n-1\}$.

An element x of \mathbb{Z}/n is called "invertible" or a "unit" if $\exists y \in \mathbb{Z}/n$ such that $xy \equiv 1(n)$.

Theorem 1.1. x is invertible modulo n iff x and n are coprime

Recall Two numbers are coprime if their highest common factor is 1.

Here's how we find the inverse of x in \mathbb{Z}/n . Since X and n are coprime we can find $h, k \in \mathbb{Z}$ such that $hx + kn = 1 \implies hx \equiv 1 \pmod{n}$. So h is the inverse of x modulo n .

E.g We'll find the inberse of 7 modulo 25 using Euclid's algorithm

$$25 = 3 \times 7 + 4$$

$$7 = 1 \times 4 + 3$$

$$4 = 1 \times 3 + 1$$

$$1 = 4 - 1(3)$$

$$1 = 4 - 1(7 - 1(4)) = 2(4) - 1(7)$$

$$1 = 2(25 - 3(7)) - 1(7) = 2(25) - 7(7)$$

$$2(25) - 7(7) = 1$$

$$- 7(7) = 1 \pmod{25}$$

$$(7^{-1}) = -7 = 18 \pmod{25}$$

$$7 \times 18 = 126 = 1 \pmod{25}$$

We'll write $(\mathbb{Z}/n)^\times$ for the invertible elements in \mathbb{Z}/n

E.g

$$(\mathbb{Z}/3)^\times = \{ \emptyset, 1, 2 \}$$

$$(\mathbb{Z}/6)^\times = \{ \emptyset, 1, \cancel{2}, \cancel{3}, \cancel{4}, 5 \}$$

Theorem 1.2. $(\mathbb{Z}/n)^\times$ is a group with the operation of multiplicity.

1.2.2 Solving Linear Congruences

Suppose we want to solve $ax \equiv b \pmod{n}$ (given a, b and n).

Case 1: If a is coprime to n then we can find a^{-1} modulo n by Euclid's algorithm,
 $x \equiv a^{-1}b \pmod{n}$

Case 2: If a is a factor of n , then there are two possibilities:

2a) if a is also a factor of b then $ax \equiv b \pmod{n}$ is equivalent to $x = \frac{b}{a} \pmod{\frac{n}{a}}$

2b) if a is not a factor of b then there are no solutions

E.g. Solve $5x = 11 \pmod{13}$

This is case 1 because 5 and 13 are coprime

$$13 = 2 \times 5 + 3$$

$$5 = 1 \times 3 + 2$$

$$3 = 1 \times 2 + 1$$

$$1 = (3) - 1(2)$$

$$1 = (3) - 1(5 - 1(3)) = 2(3) - (5)$$

$$1 = 2(13 - 2(5)) - (5) = 2(13) - 5(5)$$

$$1 \equiv -5(5) \pmod{13}$$

$$5^{-1} \equiv -5 \equiv 8 \pmod{13}$$

$$5x \equiv 11 \pmod{13}$$

$$x \equiv 8 \times 11 \equiv 88 \pmod{13}$$

$$x \equiv 10 \pmod{13}$$

E.g. Solve $7x \equiv 84 \pmod{490}$

7 is a factor of 490 so case 2)

7 is a factor of 84 so case 2a)

$$7x \equiv 84 \pmod{490}$$

$$x \equiv 12 \pmod{70}$$

E.g. Solve $7x \equiv 85 \pmod{490}$

This is case 2b (7 is a factor of 490 but not of 85) \therefore No solutions

$$7x \equiv 85 \pmod{490}$$

$$\implies 7x = 85 + 490y \text{ for some } y \in \mathbb{Z}$$

$$\implies 0 \equiv 1 \pmod{7}$$

E.g. Solve $6x \equiv 3 \pmod{21}$

This is neither case 1 nor case 2 but we can rewrite as:

$$3(2x) \equiv 3 \pmod{21}$$

$$\text{By case 2 we can solve for } 2x \equiv 1 \pmod{7}$$

but now 2 is invertible modulo 7 so now solve by case 1

$$\therefore x \equiv 4 \pmod{7}$$

1.3 Chinese Remainder Theorem

Suppose we know the congruency class of x modulo 10. Then we can work out its congruency class mod 2 and mod 5.

E.g. if $x \equiv 7 \pmod{10}$, then $x \equiv 1 \pmod{2}$ and $x \equiv 2 \pmod{5}$

Then the Chinese Remainder Theorem allows us to do the opposite, i.e. if we know x modulo 2 and modulo 5, then we can work out the value of x modulo 10.

Suppose n & m are coprime positive integers, let $a \in (\mathbb{Z}/n)$ and $b \in (\mathbb{Z}/m)$ then there is a unique

$$x \in (\mathbb{Z}/nm) \text{ such that } \begin{aligned} x &\equiv a \pmod{n} \\ x &\equiv b \pmod{m} \end{aligned}$$

Proof of existence part:

Since n & m are coprime, we can find $h, k \in \mathbb{Z}$ such that $hn + km = 1$.

Let $x = hnb + kma$

Check that this a solution to both congruences:

$$\begin{aligned} x &\equiv kma \pmod{n} \\ x &\equiv (1 - hn)a \pmod{n} \\ x &\equiv (1)a \pmod{n} \\ x &\equiv a \pmod{n} \end{aligned}$$

Similarly, this holds for $x \equiv b \pmod{m}$.

E.g. Solve the simultaneous congruence:

$$\begin{aligned} x &\equiv 3 \pmod{8} \\ x &\equiv 4 \pmod{5} \end{aligned}$$

By the Chinese Remainder Theorem, there is unique solution modulo 40. To find the solution we let $x = hnb + kma$.

First find h, k by Euclid's algorithm.

$$\begin{aligned} 8 &= 1 \times 5 + 3 & 1 &= (3) - 1(2) \\ 5 &= 1 \times 3 + 2 & 1 &= (3) - 1(5 - 1(3)) = 2(3) - (5) \\ 3 &= 1 \times 2 + 1 & 1 &= 2(8 - 2(5)) - (5) = 2(8) - 5(5) \end{aligned}$$

$$\begin{aligned} \therefore x &= (2 * 8 * 4) - (3 * 5 * 3) \\ x &= 64 - 45 \\ \implies x &\equiv 19 \pmod{40} \end{aligned}$$

Remark: We can use the Chinese Remainder Theorem to solve a congruence modulo nm , by first solving mod n and then mod m and then combining the results.

E.g. Solve $x^2 \equiv 2 \pmod{119}$. Note $119 = 7 * 17$.

By CRT this is equivalent to:

$$\begin{aligned} x^2 &\equiv 2 \pmod{7} & \implies x &\equiv \pm 3 \pmod{7} \\ x^2 &\equiv 2 \pmod{17} & \implies x &\equiv \pm 6 \pmod{17} \end{aligned}$$

Now we combine the solutions:

$$\begin{aligned} 17 &= 2 * 7 + 3 & 1 &= (7) - 2(3) \\ 7 &= 2 * 3 + 1 & 1 &= (7) - 2(17 - 2(7)) \\ & & 1 &= 5(7) - 2(17) \end{aligned}$$

Since

$$\begin{array}{ll} x \equiv \pm 3 \pmod{7} & \text{We get } x \equiv 5 * 7 * (\pm 6) - 2 * 17 * (\pm 3) \\ x \equiv \pm 6 \pmod{17} & x \equiv \pm 11 \text{ or } \pm 45 \pmod{119} \end{array}$$

1.4 Prime numbers

Defintion 1.3. An integer $p \geq 2$ is a prime number if the only factors of p are $\pm 1, \pm p$

We'll write \mathbb{F}_p for \mathbb{Z}/p . This is because:

Theorem 1.4. If p is prime, then \mathbb{F}_p is a field

Proof. Need to check that the non-zero elements of \mathbb{F}_p all have inverses.

Let $x \in \mathbb{F}_p$ with $x \not\equiv 0 \pmod{p}$ i.e. x is not a multiple of p

$$\therefore \text{hcf}(x, p) = 1$$

$\therefore x$ & p coprime □

1.5 Fermat's Little Theorem

Theorem 1.5. Let p be a prime number. If x is not a multiple of p then $x^{p-1} \equiv 1 \pmod{p}$

Proof. $x \in \mathbb{F}_p^\times = \{1, 2, \dots, p-1\}$ a group with $p-1$ elements.

Let n be the order of x in this group.

(order of x is smallest $n > 0$ such that $x^n \equiv 1 \pmod{p}$)

By corollary to Lagrange's Theorem, $p-1$ is a multiple of n

$$\begin{aligned} x^n &\equiv 1 \pmod{p} \\ x^{p-1} &\equiv 1 \pmod{p} \end{aligned} \quad \square$$

Theorem 1.6. Lagrange's Theorem: If H is a subgroup of a finite group G , then $|H|$ is a factor of $|G|$.

Corollary 1.7. Order of an element is a factor of $|G|$

We can use Fermat's Little Theorem to do calculations.

E.g. Calculate 10^{100} modulo 19

By Fermat's Little Theorem: $10^{18} \equiv 1 \pmod{19}$

$$\begin{aligned} 10^{100} &\equiv (10^{18})^5 * 10^{10} \pmod{19} \\ &\equiv 100^5 \pmod{19} \\ &\equiv 5^5 \pmod{19} \\ &\equiv 25 * 125 \equiv 6 * 11 \equiv 9 \pmod{19} \end{aligned}$$

Also using Fermat's Little Theorem we can solve congruence of the form $x^a \equiv b \pmod{p}$ as long as p prime and a invertible modulo $p-1$

1.5.1 General method to solve $x^a \equiv b \pmod{p}$

Let

$$\begin{aligned}c &= a^{-1} \pmod{p-1} \\ac &= 1 + (p-1)r\end{aligned}$$

Raise both sides of the congruence to power c :

$$\begin{aligned}\therefore x^{ac} &\equiv b^c \pmod{p} \\x^{1+(p-1)r} &\equiv b^c \pmod{p} \\x &\equiv b^c\end{aligned}$$

So the solution is $x \equiv b^c \pmod{p}$

E.g. Solve $x^5 \equiv 2 \pmod{19}$

19 is prime and 5 is coprime to 18.

Find $c = 5^{-1} \pmod{18}$

$$\begin{array}{ll}18 = 3 * 5 + 3 & 1 = 2 * 3 - 5 \\5 = 2 * 3 - 1 & 1 = 2(18 - 3 * 5) - 5 \\& 1 = 2 * 18 - 7 * 5\end{array}$$

$$\begin{aligned}\therefore 5^{-1} &\equiv -7 \pmod{18} \\&\equiv 11 \pmod{18}\end{aligned}$$

$$\begin{aligned}\therefore x &\equiv 2^{11} \pmod{19} \\&\equiv 2048 \pmod{19} \\&\equiv 15 \pmod{19}\end{aligned}$$

1.6 Fundamental Theorem of Arithmetic

If n is a positive integer then there is a unique factorisation, $n = p_1 p_2 \dots p_r$ with p_i prime. "Unique" means up to reordering the primes p_1, \dots, p_r . Showing that a factorisation exists is easy. For the uniqueness part we use:

1.6.1 Euclid's Lemma

Lemma 1.8. Suppose p prime, and $p|ab$. Then $p|a$ or $p|b$.

To prove Euclid's lemma we use Bezout's lemma.

Proof. Assume $p|ab$ but $p \nmid a$. Then $\text{hcf}(a, p) = 1$

By Bezout's lemma, $\exists h, k$ such that:

$$1 = ha + kp$$

$$b = hab + kpb$$

Both hab and kpb are multiples of p .

$\therefore p|b$

□

If n is composite then the smallest factor of n is (apart from 1) is a prime number $p \leq \sqrt{n}$, i.e. to show that n is prime, we just need to show that none of the primes up to \sqrt{n} are factors of n .

$$\sqrt{199} < 15 \text{ since } 15^2 = 225$$

The primes up to 15 are ~~2~~, ~~3~~, ~~5~~, ~~7~~, ~~11~~, ~~13~~

$199 \equiv 3 \pmod{2}$ (7)

$199 \equiv 4 \pmod{3}$ (13)

$\therefore 199$ is prime

Proof. Suppose p_1, \dots, p_n are all the primes.

$\therefore N$ has no prime factors \nmid

Proof. Assume there are only finitely many primes, call them p_1, p_2, \dots, p_r . All other primes are either 3 or are congruent to 1 mod 3.

$\therefore N \equiv 1 \pmod{3} \implies$ because clearly $N \equiv 2 \pmod{3}$

2 Elementary Number Theory

2.1 Euler Totient Function

Recall $(\mathbb{Z}/n)^\times$ is the group of invertible elements in \mathbb{Z}/n .

E.g. $(\mathbb{Z}/6)^\times = \{1, 5\}$

$(\mathbb{Z}/8)^\times = \{1, 3, 5, 7\}$

These are groups with the multiplication operation, $*$. The multiplication table for $(\mathbb{Z}/8)^\times$ is given below.

$*$	1	3	5	7
1	1	3	5	7
3	3	1	7	5
5	5	7	1	3
7	7	5	3	1

Definition 2.1. The Euler Totient function is $\phi(n) = |(\mathbb{Z}/n)^\times|$

E.g. $\phi(6) = 2$

$\phi(8) = 4$

If p prime then $(\mathbb{Z}/p)^\times = \{1, \dots, p-1\}$ so $\phi(p) = p-1$

Theorem 2.2. Euler's Theorem- Let $x \in (\mathbb{Z}/n)^\times$ then $x^{\phi(n)} \equiv 1 \pmod{n}$

In the case $n = p$ is prime, this is just Fermat's Little Theorem.

Proof. Let d be the order of x , i.e. $x^d \equiv 1 \pmod{n}$. By a corollary to Lagrange's Theorem, d is a factor of $\phi(n) \implies x^{\phi(n)} \equiv 1 \pmod{n}$ \square

We can use Euler's theorem to solve congruences and calculate powers mod n . To use the theorem, we need a quick way of calculating $\phi(n)$.

Lemma 2.3. Let $n = p^a$ where p is prime $a > 0$. Then $\phi(n) = (p-1)p^{a-1}$

E.g. $\phi(8) = \phi(2^3) = (2-1)2^{3-1} = 4$

Proof. An integer is coprime to p^a as long as it's not a multiple of p .

\therefore The elements of \mathbb{Z}/p^a which are not invertible are the multiples of p . $0, p, 2p, \dots, p^a - p$.

There are $p^a - 1$ of these:

$$\therefore |(\mathbb{Z}/p^a)^\times| = p^a - p^{a-1} = (p-1)p^{a-1} \quad \square$$

Theorem 2.4. Let n and m be coprime. Then there is an isomorphism:

$$(\mathbb{Z}/nm)^\times \cong (\mathbb{Z}/n)^\times * (\mathbb{Z}/m)^\times$$

We'll use the theorem before we prove it.

Remark: If G and H are groups, $G \times H = \{(x, y) : x \in G, y \in H\}$, then $G \times H$ is a group with the operation $(x, y)(x', y') = (xx', yy')$ and $G \times H$ is the "direct product" of G and H

Corollary 2.5. *If n and m are coprime then $\phi(nm) = \phi(n)\phi(m)$*

Proof.

$$\begin{aligned}\phi(nm) &= |(\mathbb{Z}/nm)^\times| = |(\mathbb{Z}/n)^\times * (\mathbb{Z}/m)^\times| \\ &= |(\mathbb{Z}/n)^\times| * |(\mathbb{Z}/m)^\times| \\ &= \phi(n)\phi(m)\end{aligned}$$

□

Corollary 2.6. *(Corollary of the corollary): Suppose $n = p_1^{a_1} \dots p_r^{a_r}$ with p_1, \dots, p_r distinct primes and $a_i > 0$. Then*

$$\phi(n) = (p_1 - 1)p_1^{a_1-1} * \dots * (p_r - 1)p_r^{a_r-1}$$

Proof. Since $p_1^{a_1}, \dots, p_r^{a_r}$ are coprime,

$$\begin{aligned}\phi(n) &= \phi(p_1^{a_1}) \dots \phi(p_r^{a_r}) && \text{by the corollary} \\ &= (p_1 - 1)p_1^{a_1-1} \dots (p_r - 1)p_r^{a_r-1} && \text{by the lemma}\end{aligned}$$

□

E.g. Calculate $\phi(200)$

$$\begin{aligned}\phi(200) &= \phi(2^3 * 5^2) \\ &= (2 - 1)2^{3-1} * (5 - 1)5^{2-1} \\ &= 4 * 4 * 5 \\ &= 80\end{aligned}$$

Theorem 2.7. *Suppose n and m are coprime, then $(\mathbb{Z}/nm)^\times \cong (\mathbb{Z}/n)^\times * (\mathbb{Z}/m)^\times$. The isomorphism is the map $x \mapsto (x \bmod n, x \bmod m)$*

E.g. $n = 4, m = 5$

$$\begin{aligned}(\mathbb{Z}/4)^\times &= \{1, 3\} \\ (\mathbb{Z}/5)^\times &= \{1, 2, 3, 4\} \\ \therefore (\mathbb{Z}/4)^\times * (\mathbb{Z}/5)^\times &= \{(1, 1), (1, 2), (1, 3), (1, 4), \\ &\quad (3, 1), (3, 2), (3, 3), (3, 4)\} \\ (\mathbb{Z}/20)^\times &= \{1, 3, 7, 9, 11, 13, 17, 19\}\end{aligned}$$

The isomorphism is:

$$\begin{array}{ll} 1 \mapsto (1, 1) & 11 \mapsto (3, 1) \\ 3 \mapsto (3, 3) & 13 \mapsto (1, 3) \\ 7 \mapsto (3, 2) & 17 \mapsto (1, 2) \\ 9 \mapsto (1, 4) & 19 \mapsto (3, 4) \end{array}$$

Proof. Let $\Phi : \mathbb{Z}/nm \mapsto \mathbb{Z}/n * \mathbb{Z}/m$

$$\Phi(x) = (x \bmod n, x \bmod m)$$

This is a bijection by the Chinese Remainder Theorem.

We'll next show that x is invertible mod $nm \iff x$ is invertible mod n and mod m

(\implies) Suppose x is invertible mod nm

$$\text{Let } xy \equiv 1 \pmod{nm}$$

$$\therefore xy \equiv 1 \pmod{n}$$

$$xy \equiv 1 \pmod{m}$$

$$\therefore x \text{ invertible mod } n \text{ and } m$$

(\impliedby) Suppose x invertible mod n and m

$$xa \equiv 1 \pmod{n}$$

$$xb \equiv 1 \pmod{m}$$

By the Chinese Remainder Theorem, $\exists y$ such that $y \equiv a \pmod{n}$

$$y \equiv b \pmod{m}$$

$$\left. \begin{array}{l} \therefore xy \equiv xa \equiv 1 \pmod{n} \\ \equiv xb \equiv 1 \pmod{m} \end{array} \right\} \implies xy \equiv 1 \pmod{nm} \text{ by the Chinese Remainder Theorem}$$

We've shown that Φ gives a bijection between $(\mathbb{Z}/nm)^\times$ and $(\mathbb{Z}/n)^\times * (\mathbb{Z}/m)^\times$. We'll next check that $\Phi(xy) = \Phi(x)\Phi(y)$.

$$\begin{aligned} \Phi(xy) &= (xy \bmod n, xy \bmod m) \\ &= (x \bmod n, x \bmod m) * (y \bmod n, y \bmod m) \\ &= \Phi(x)\Phi(y) \end{aligned}$$

□

2.2 Euler's Theorem

If $x \in (\mathbb{Z}/n)^\times$ then $x^{\phi(n)} \equiv 1 \pmod{n}$ and $\phi(p_1^{a_1} \dots p_r^{a_r}) = (p_1 - 1)p_1^{a_1-1} \dots (p_r - 1)p_r^{a_r-1}$

E.g. Calculate $7^{135246872002} \bmod 10000$

$$7 \text{ coprime to } 10000 \text{ so } 7^{\phi(10000)} \equiv 1 \pmod{10000}$$

$$10000 = 2^4 * 5^4$$

$$\therefore \phi(10000) = (2-1)2^3 * (5-1) * 5^3 = 8 * 500$$

$$7^{4000} \equiv 1 \pmod{10000} \implies 7^n \text{ depends only on } n \bmod 4000$$

$$135246872002 \equiv 2 \pmod{4000}$$

$$\therefore 7^{135246872002} \equiv 7^2 \equiv 49 \pmod{10000}$$

We can also use Euler's THEorem to solve congruence with powers

2.2.1 Solving equations of the form $x^a \equiv b \pmod{n}$

Suppose we want to solve $x^a \equiv b \pmod{n}$ where b is coprime to n and a is coprime to $\phi(n)$.

Clearly any solution x must be coprime to n by Euler's Theorem $x^{\phi(n)} \equiv 1 \pmod{n}$.

\therefore The congruency class of $x^y \pmod{n}$ depends only $y \pmod{\phi(n)}$

Let

$$c = a^{-1} \pmod{\phi(n)}$$

Raise both sides of the congruence to power c :

$$x^{ac} \equiv x^1 \equiv b^c \pmod{n}$$

\therefore The solution is $x \equiv b^c \pmod{n}$

E.g. $x^7 \equiv 3 \pmod{50}$

3 is coprime to 50,

$$\begin{aligned} 50 &= 2 * 5^2 \\ \implies \phi(50) &= 1 * 4 * 5 = 20 \end{aligned}$$

7 is coprime to $\phi(50)$. To solve, we need to find

$$\begin{aligned} c &\equiv 7^{-1} \pmod{\phi(50)} \\ &\equiv 3 \pmod{20} \end{aligned}$$

$$x \equiv 3^3 \equiv 27 \pmod{50}$$

E.g. $x^{27} \equiv 5 \pmod{123}$

5 is coprime to 123,

$$\begin{aligned} 123 &= 3 * 41 \\ \implies \phi(123) &= 2 * 40 = 80 \end{aligned}$$

27 is coprime to 80

To solve, we find $27^{-1} \pmod{80}$

$$\begin{aligned} 80 &= 3 * 27 - 1 \\ \implies 1 &= 3 * 27 - 80 \end{aligned}$$

$$27^{-1} = 3$$

$$\begin{aligned} x &= 5^3 \\ x &= 125 \equiv 2 \pmod{123} \end{aligned}$$

2.3 Primitive roots

Recall, let G be a finite group. G is called a cyclic group if $\exists x \in G$ such that, every element in G has the form x^n for some $n \in \mathbb{Z}$, i.e. $G = \{1, x, x^2, \dots, x^{n-1}\}$ where n is the order of x , equivalentl the order of x is $|G|$. The element x is called a generator of G .

Theorem 2.8. (Gauss' Theorem), For ever prime number p , the group \mathbb{F}_p^\times is cyclic

Defintion 2.9. A generator of \mathbb{F}_p^\times is called a primitive root. Equivalently, this is an element of order $p - 1$

E.g. $p = 7, x = 3$ We'll see that 3 is a primitive root modulo 7

$$\begin{array}{llll} \text{Powers of 3 in } F_7^\times : & 3^0 = 1 & 3^3 \equiv 6 \pmod{7} & 3^6 \equiv 1 \pmod{7} \\ & 3^1 = 3 & 3^4 \equiv 4 \pmod{7} & \\ & 3^2 \equiv 2 \pmod{7} & 3^5 \equiv 1 \pmod{7} & \end{array}$$

so 3 is a primitive root modulo 7. There is a quicker way to check whether x is a primitive root.

Proposition 2.10. Let $x \in \mathbb{F}_p^\times$, then x is a primitive root modulo p if and only if for every prime factor q of $p - 1$:

$$x^{\frac{p-1}{q}} \not\equiv 1 \pmod{p}$$

Proof. Assume the second statement is false, so \exists prime factor q of $p - 1$ such that:

$$\begin{array}{ll} x^{\frac{p-1}{q}} \equiv 1 \pmod{p} & \therefore \text{order of } x \leq \frac{p-1}{q} < p-1 \\ & \therefore x \text{ is not a primitive root} \end{array}$$

Conversely, assume x is not a primitive root, so x doe not have order $p - 1$. But the order of x is a factor of $p - 1$.

Suppose the order of x is $\frac{p-1}{d}$, $d > 1$.

Let q be a prime factor of $d \implies q|p-1$

$$\frac{p-1}{q} \text{ is a multiple of } \frac{p-1}{d} \text{ but } x^{\frac{p-1}{q}} \equiv 1 \pmod{p} \implies x^{\frac{p-1}{d}} \equiv 1 \pmod{p}$$

□

E.g. $p = 29$

By the proposition x is a primitive root mod 29 $\iff x^{28/2} \not\equiv 1 \pmod{29}$ and $x^{28/7} \not\equiv 1 \pmod{29}$

$$\iff x^{14} \not\equiv 1 \pmod{29} \text{ and } x^4 \not\equiv 1 \pmod{29}$$

$$\begin{array}{ll} \text{Try } x = 2 : & 2^4 \equiv 16 \not\equiv 1 \pmod{29} \\ & 2^{14} \equiv 128^2 \equiv 12^2 \equiv 144 \equiv -1 \pmod{29} \end{array}$$

$\therefore 2$ is a primitive root mod 29

Another trick to speed up the calculation:

\mathbb{F}_p is a field \therefore every polynomial of d has no more than d in \mathbb{F} (proved in 2201).

\therefore if $x^2 \equiv 1 \pmod{p}$ then $x \equiv \pm 1 \pmod{p}$

This means that checking whether $x^{14} \equiv 1 \pmod{29}$ is equivalent to checking whether $x^7 \equiv \pm 1 \pmod{29}$.

E.g 3 is also a primitive root modulo 29

$$3^2 \equiv 9 \not\equiv \pm 1 \pmod{29}$$

$$3^4 \equiv 1 \pmod{29}$$

$$3^7 \equiv 27^2 * 3 \pmod{29}$$

$$\equiv (-2)^2 * 3 \equiv 12 \pmod{29}$$

$$\equiv \pm 1 \pmod{29}$$

$$\therefore 3^{14} \not\equiv 1 \pmod{29}$$

2.4 Roots of unity and Cyclotomic Polynomials

A complex number ζ is called an n^{th} root of unity if $\zeta^n = 1$. The n^{th} roots of unity are $e^{2\pi i \frac{a}{n}}$ for $a = \{0, 1, \dots, n-1\}$

We call ζ a primitive n^{th} root of unity if n is the smallest power than ζ^n is equal to 1, i.e. ζ has order n in \mathbb{C}^\times if ζ is not a primitive n^{th} root of unity $\zeta = e^{2\pi i \frac{b}{d}}$ where $b = \{0, \dots, d-1\}$ for $d < n$

$$\therefore \frac{a}{n} = \frac{b}{d}$$

The cancellation happens when a is not coprime to n . This shows that the primitive n^{th} roots of unity are $e^{2\pi i \frac{a}{n}}$, $a \in (\mathbb{Z}/n)^\times$.

Corollary 2.11. *There are exactly $\phi(n)$ primitive n^{th} roots of unity*

We'll actually prove a more precise version of Gauss' Theorem.

Theorem 2.12. *For every factor d of $p-1$ there are $\phi(d)$ elements in \mathbb{F}_p^\times of order d .*

Definition 2.13. *The n^{th} cyclotomic polynomial is:*

$$\Phi_n(x) = \prod_{\substack{\text{primitive} \\ n^{th} \text{ roots} \\ \text{of unity } \zeta}} (X - \zeta)$$

i.e. $\zeta^n = 1$ and no smaller power of ζ is 1, $\zeta = e^{2\pi i \frac{a}{n}}$, $a \in (\mathbb{Z}/n)^\times$

This has degree $\phi(n)$.

E.g. $n=4$

Primitive 4^{th} roots of unity are $i, -i$:

$$\begin{aligned}\Phi_4(x) &= (x - i)(x - (-i)) \\ &= x^2 + 1\end{aligned}$$

Lemma 2.14. For every $n > 0$:

$$x^n - 1 = \prod_{\substack{d \text{ factors} \\ d \text{ of } n}} \Phi_d(x)$$

E.g. Calculate $\Phi_6(x)$

$$\begin{aligned}\text{By the lemma} \quad x^6 - 1 &= \Phi_1 \Phi_2 \Phi_3 \Phi_6 & x^6 - 1 &= (x^3 - 1) \Phi_2 \Phi_6 \\ x^3 - 1 &= \Phi_1 \Phi_3\end{aligned}$$

$$\therefore \Phi_6 = \frac{x^6 - 1}{(x^3 - 1)(x + 1)} = \frac{x^3 + 1}{x + 1} = x^2 - x + 1$$

Let p be a prime number. A primitive root mod p is an $x \in \mathbb{F}_p^\times$, such that x generates \mathbb{F}_p^\times .
Equivalently order = $p - 1$

2.4.1 How to calculate $\Phi_n(x)$

Lemma 2.15. $x^n - 1 = \prod_{d|n} \Phi_d(x)$

E.g. $n = 4$

$$\begin{aligned}x^4 - 1 &= \Phi_1 \Phi_2 \Phi_4 & \Phi_1 &= x - 1 \\ & & \Phi_2 &= (x - (-1)) = x + 1 \\ & & \Phi_4 &= (x - i)(x - (-i)) = x^2 + 1 \\ &= (x - 1)(x + 1)(x^2 + 1)\end{aligned}$$

Proof.

$$x^n - 1 = \prod_{\substack{\zeta \text{ is an} \\ n^{th} \text{ root of} \\ \text{unity}}} (x - \zeta)$$

but every n^{th} root of unity is a primitive d^{th} root of unity for some $d|n$.

$$x^n = \prod_{d|n} (\prod_{\substack{\text{primitive} \\ d^{th} \text{ roots} \\ \text{of unity}}} (x - \zeta)) = \prod_{d|n} \Phi_d(x)$$

□

E.g. Calculate $\Phi_5(x)$

$$\begin{aligned} x^5 - 1 &= \Phi_1(x)\Phi_5(x) \\ &= (x - 1)\Phi_5(x) \end{aligned}$$

$$\Phi_5(x) = \frac{x^5 - 1}{x - 1} = 1 + x + x^2 + x^3 + x^4$$

More generally if p prime then $x^p - 1 = (x - 1)\Phi_p(x) \implies \Phi_p(x) = 1 + x + \dots + x^{p-1}$

E.g. Calculate $\Phi_8(x)$

$$x^8 - 1 = \Phi_1(x)\Phi_2(x)\Phi_4(x)\Phi_8(x)$$

$$x^4 - 1 = \Phi_1(x)\Phi_2(x)\Phi_4(x) \implies \Phi_8(x) = \frac{x^8 - 1}{x^4 - 1} = x^4 + 1$$

Corollary 2.16. $\Phi_n(x)$ has coefficients in \mathbb{Z}

$$\text{Proof. } \Phi_n(x) = \frac{x^n - 1}{\prod_{\substack{d|n \\ d \neq n}} \Phi_d(x)}$$

We'll prove the corollary by induction on n , clearly true when $n = 1$. Assume Φ_d has integer coefficients $\forall d < n$.

It is proved in Algebra 3 (MATH2201) that, if $f, g \in \mathbb{Z}[X]$ and g monic then $f = qg + r$ where $\deg(r) < \deg(g)$ and $g, r \in \mathbb{Z}[x]$.

Using this, we get that the denominator $\prod_{\substack{d|n \\ d \neq n}} \Phi_d(x)$ is a monic polynomial with coefficients in $\mathbb{Z} \implies \Phi_n \in \mathbb{Z}[X]$. □

2.4.2 Gauss' Theorem

Theorem 2.17. Let n be a factor of $p - 1$, where p is prime. Then there are exactly $\phi(n)$ elements of order n in \mathbb{F}_p^\times . These are the roots of Φ in \mathbb{F}_p^\times . In particular there are $\phi(p - 1)$ primitive roots.

Proof. Let $f(x) = x^{p-1} - 1$

By Fermat's Little theorem, $f(x) = 0 \pmod{p}$ for $x = 1, \dots, p - 1$ for $(x \neq 0)$

$$\begin{aligned} \therefore f(x) &= (x - 1)(x - 2) \dots (x - (p - 1)) \\ &= \prod_{n|p-1} \Phi_n(x) \end{aligned}$$

This implies that:

- Each Φ_n (for $n|p - 1$) factorises completely into linear factors with no repeated roots $\therefore \Phi_n$ has $\phi(n)$ roots in \mathbb{F}_p
- Every element of \mathbb{F}_p^\times is a root of exactly one of the polynomials Φ_n with $n|p - 1$

It remains to show that the roots of $\Phi_n(x)$ in \mathbb{F}_p has order of exactly n .
 Suppose $\Phi_n(x) \equiv 0 \pmod{p}$

By the lemma $\Phi_n(x)$ is a factor $x^n - 1$
 $\therefore x^n - 1 \equiv 0 \pmod{p}$
 $\therefore x^n \equiv 1 \pmod{p}$

Suppose $x^m \equiv 1 \pmod{p}$ for some $m|n, m < n$
 $\implies x^m - 1 \equiv 0 \pmod{p}$

By the lemma $\Pi_{d|m} \Phi_d(x) \equiv 0 \pmod{p}$
 $\implies \Phi_d(x) \equiv 0 \pmod{p}$ for some $d \nmid n$

We already know that x is only a root of 1 of the cyclotomic polynomials, therefore x has order n . \square

2.5 Quadratic reciprocity (Quadratic equations modulo prime numbers)

Recall we can solve $x^a \equiv b \pmod{p}$ as long as a is coprime to $p - 1$. This won't work if $a = 2$ because a will not be invertible mod $p - 1$. An easier question to ask is, which quadratic equations have solutions modulo p ?

E.g. Does $x^2 \equiv 37 \pmod{149}$ have solutions?

Notation: We always let p be an odd prime (i.e. $p \neq 2$)

An element $a \in \mathbb{F}_p^\times$ is a quadratic residue if $x^2 \equiv a \pmod{p}$ has solutions.

An element $a \in \mathbb{F}_p^\times$ is a quadratic non-residue if there are no solutions.

The quadratic residue symbol is defined for $a \in \mathbb{F}_p^\times$ by

$$\left(\frac{a}{p}\right) = \begin{cases} 1 & \text{a quadratic residue} \\ -1 & \text{a quadratic non-residue} \end{cases}$$

Lemma 2.18. *Let g be a primitive root modulo p (p odd prime). Then g^r is a quadratic residue iff r even.*

Proof.

(\Leftarrow) Assume r even

Clearly g^r is a square in \mathbb{F}_p^\times

So g^r is a quadratic residue

(\Rightarrow) Assume $g^r \equiv x^2 \pmod{p}$

$x \equiv g^s \pmod{p}$ ($s \in \mathbb{Z}$) since g primitive roots

$\therefore g^r \equiv g^{2s} \pmod{p}$

$g^{r-2s} \equiv 1 \pmod{p}$

g has order $p - 1$, so $r - 2s$ is a multiple of $p - 1$

p odd $\implies p - 1$ is even $\implies r$ is even

\square

E.g. $p = 7$

x	$x^2 \pmod{7}$		a	$\left(\frac{a}{7}\right)$
± 1	1	\implies	1	1
± 2	4		2	1
± 3	2		3	-1
			4	1
			5	-1
			6	-1

So 1,2,4 are quadratic residues; 3,4,6 are quadratic non-residues

Corollary 2.19. *There are exactly $\frac{p-1}{2}$ quadratic residues and $\frac{p-1}{2}$ quadratic non-residues mod p*

Defintion 2.20. *Euler's criterion: Let p be an odd prime and $a \in \mathbb{F}_p^\times \implies \left(\frac{a}{p}\right) \equiv a^{\frac{p-1}{2}} \pmod{p}$*
Also $\left(\frac{ab}{p}\right) = \left(\frac{a}{p}\right)\left(\frac{b}{p}\right)$

Proof. $(a^{\frac{p-1}{2}})^2 \equiv 1 \pmod{p}$ by Fermat's Little theorem.

$$\therefore a^{\frac{p-1}{2}} \equiv \pm 1 \pmod{p}$$

Let $a = g^r$ where g is a primitive root $\implies a^{\frac{p-1}{2}} \equiv g^{(p-1)\frac{r}{2}}$

$$\begin{aligned} a \text{ is a quadratic residue} &\iff r \text{ is even} \\ &\iff (p-1)\frac{r}{2} \text{ is a multiple of } p-1 \\ &\iff g^{(p-1)\frac{r}{2}} \equiv 1 \pmod{p} \\ &\iff a^{\frac{p-1}{2}} \equiv 1 \pmod{p} \end{aligned}$$

□

To calculate $\left(\frac{a}{p}\right)$, we'll use three theorems:

2.5.1 Quadratic Reciprocity Law

Let p, q be distinct odd prime numbers. Then $\left(\frac{p}{q}\right) = (-1)^{\frac{(p-1)(q-1)}{4}}$

$$\text{i.e. } \left(\frac{p}{q}\right) = \begin{cases} \left(\frac{q}{p}\right) & \text{if } p \equiv 1 \pmod{4} \text{ or } q \equiv 1 \pmod{4} \\ -\left(\frac{q}{p}\right) & \text{if } p \equiv q \equiv -1 \pmod{4} \end{cases}$$

2.5.2 First Nebensatz

If p is an odd prime, then $\left(\frac{-1}{p}\right) = (-1)^{\frac{p-1}{2}}$

$$\text{i.e. } \left(\frac{-1}{p}\right) = \begin{cases} 1 & p \equiv 1 \pmod{4} \\ -1 & p \equiv -1 \pmod{4} \end{cases}$$

2.5.3 Second Nebensatz

Let p be an odd prime, then $(\frac{2}{p}) = (-1)^{\frac{p^2-1}{8}}$

$$\text{i.e. } (\frac{2}{p}) = \begin{cases} 1 & p \equiv \pm 1 \pmod{8} \\ -1 & p \equiv \pm 3 \pmod{8} \end{cases}$$

We'll prove the theorems later.

E.g. Does the congruence $x^2 \equiv 37 \pmod{199}$ have solutions?

$$\begin{aligned} 199 \text{ is an odd prime } (\frac{37}{199}) &= +(\frac{199}{37}) && \text{by quadratic reciprocity} \\ &\equiv (\frac{14}{37}) && \text{because } 199 \equiv 14 \pmod{37} \\ &\equiv (\frac{2}{37})(\frac{7}{37}) && \text{by the corollary} \\ &\equiv (-1)(\frac{7}{37}) && \text{by the 2}^{nd} \text{ Nebensatz} \\ &\equiv (-1)(+1)(\frac{37}{7}) && \text{by the quadratic reciprocity law} \\ &\equiv -(\frac{2}{7}) && \text{because } 37 \equiv 2 \pmod{7} \\ &\equiv -(+1) && \text{by the 2}^{nd} \text{ Nebensatz} \\ &\equiv -1 && \therefore x^2 \equiv 37 \pmod{199} \text{ has no solutions} \end{aligned}$$

E.g. $x^2 \equiv 47 \pmod{53}$ have solutions?

$$(\frac{47}{53}) = +(\frac{53}{47}) = (\frac{6}{47}) = (\frac{2}{47})(\frac{3}{47}) = (+1)(-1)(\frac{47}{3}) = -(-\frac{1}{3}) = -(-1) = +1$$

This shows that 47 is a quadratic residue mod 53, so $x^2 \equiv 47 \pmod{53}$ does have solutions. ($x = 10$)

We can speed up the test for primitive roots using quadratic reciprocity,

$$x \text{ is a primitive root mod } p \iff \forall q|p-1, q \text{ prime } x^{\frac{p-1}{q}} \not\equiv 1 \pmod{p}$$

This means we need to calculate $x^{\frac{p-1}{q}} \pmod{p}$ for primes $q|p-1$, the biggest power of x to calculate is $x^{\frac{p-1}{2}}$. But we can calculate this, because it is $(\frac{x}{p})$ by Euler's criterion.

E.g. Is 35 a primitive root modulo 83?

The primes q dividing 82 are 2, 41, need to check $35^2, 35^{41}$
 $35^2 \not\equiv 1 \pmod{83}$ because $35 \not\equiv \pm 1 \pmod{83}$, a quadratic equation cannot have more than 2 roots.
 $35^{41} \equiv (\frac{35}{83}) \pmod{83} = (\frac{5}{83})(\frac{7}{83}) = (\frac{83}{5})(-1)(\frac{83}{7}) = (\frac{3}{5})(-1)(\frac{-1}{7}) = (\frac{5}{3})(-1)(-1) = (\frac{2}{3})$
 $= -1 \not\equiv 1 \pmod{83}$

So 35 is a primitive root modulo 83.

Proof. First Nebensatz:

By Euler's criterion, $\left(\frac{-1}{p}\right) \equiv (-1)^{\frac{p-1}{2}} \pmod{p}$.

Both sides are ± 1 , and $+1 \not\equiv -1 \pmod{p}$ because $p \geq 3 \implies$ they are equal. \square

E.g. Find the first primitive root modulo 41

$$40 = 2^3 * 5$$

$$x \in \mathbb{F}_{41}^\times \text{ is a primitive root} \iff \begin{cases} x^{\frac{40}{2}} \not\equiv 1 \pmod{41} \\ x^{\frac{40}{5}} \not\equiv 1 \pmod{41} \end{cases}$$

$$\text{We can then simplify the conditions to: } \begin{cases} \frac{x}{41} = -1 \\ x^4 \not\equiv \pm 1 \pmod{41} \end{cases}$$

$$\text{Try } x = 2 : \left(\frac{2}{41}\right) = 1 \implies \text{not a primitive root}$$

$$\text{Try } x = 3 : \left(\frac{3}{41}\right) = \left(\frac{41}{3}\right) = \left(\frac{2}{3}\right) = -1 \quad \text{and } 3^4 = 81 \equiv -1 \pmod{41} \implies \text{not a primitive root}$$

$$\text{Try } x = 4 : \implies \text{not a primitive root}$$

$$\text{Try } x = 5 : \left(\frac{5}{41}\right) = \left(\frac{41}{5}\right) = \left(\frac{1}{5}\right) = 1 \implies \text{not a primitive root}$$

$$\begin{aligned} \text{Try } x = 6 : \left(\frac{6}{41}\right) &= \left(\frac{2}{41}\right)\left(\frac{3}{41}\right) = 1 * -1 = -1 \\ 2^4 * 3^4 &= -2^4 \equiv 16 \pmod{41} \not\equiv \pm 1 \implies \text{so 6 is a primitive root} \end{aligned}$$

E.g. For which primes p does the congruence $x^2 \equiv -3 \pmod{p}$ have solutions?

Notice $x = 1$ is a solution mod 2,

$x = 2$ is a solution mod 3.

For primes $p \neq 2, 3$ it depends on $\left(\frac{-3}{p}\right)$

$$\begin{aligned} \text{We'll calculate } \left(\frac{-3}{p}\right) &= \left(\frac{-1}{p}\right)\left(\frac{3}{p}\right) \\ &= (-1)^{\frac{p-1}{2}} \left(\frac{3}{p}\right) \\ &= (-1)^{\frac{(3-1)(p-1)}{4}} \left(\frac{p}{3}\right) \\ &= \left(\frac{p}{3}\right) \end{aligned}$$

List the squares mod 3, $1^2 = 1 \pmod{3}, 2^2 = 1 \pmod{3}$

$$\therefore \left(\frac{p}{3}\right) = \begin{cases} 1 & \text{if } p \equiv 1 \pmod{3} \\ -1 & \text{if } p \equiv 2 \pmod{3} \end{cases}$$

We've shown that $x^2 \equiv -3 \pmod{p}$ has solutions iff $p \neq 2$ or $p \equiv 1 \pmod{3}$.

Corollary 2.21. *There are infinitely many primes $p \equiv 1 \pmod{3}$*

Proof. Assume there are only finitely many, and call them p_1, p_2, \dots, p_r
Let $N = n^2 + 3$ where $n = 2p_1 \dots p_r$
Take a prime factor q of N

$$N \equiv 0 \pmod{q}$$

$$n^2 + 3 \equiv 0 \pmod{q}$$

$$n^2 \equiv -3 \pmod{q}$$

We've just shown that this implies $q = 2$ or 3 or $q \equiv 1 \pmod{3}$ but $q \neq 2, 3, q \not\equiv 1 \pmod{3}$ \nmid □

Before we prove the 2^{nd} Nebensatz, we need to know about a new ring.

Let $\zeta = e^{\frac{2\pi i}{8}}$, a primitive 8^{th} root of unity.

We'll use the ring $\mathbb{Z}[\zeta] = \{f(\zeta) : f \in \mathbb{Z}\} = \{a_0 + a_1\zeta + a_2\zeta^2 + \dots + a_n\zeta^n : a_i \in \mathbb{Z}\}$

This is clearly a ring (closed under $+, *$).

2.6 Uniqueness Lemma

Every $A \in \mathbb{Z}[\zeta]$ can be written uniquely as $A = W + x\zeta + y\zeta^2 + z\zeta^3$ with $w, x, y, z \in \mathbb{Z}$.

We'll use congruence modulo p in the ring $\mathbb{Z}[\zeta]$ to prove the 2^{nd} Nebensatz.

Defintion 2.22. Let $A, B \in \mathbb{Z}[\zeta]$

We'll say $A \equiv B \pmod{p\mathbb{Z}[\zeta]}$ if $A - B = pC$ for some $C \in \mathbb{Z}[\zeta]$

$$\text{Suppose } A = a_0 + a_1\zeta + a_2\zeta^2 + a_3\zeta^3$$

$$B = b_0 + b_1\zeta + b_2\zeta^2 + b_3\zeta^3$$

$$C = c_0 + c_1\zeta + c_2\zeta^2 + c_3\zeta^3$$

The equation $A - B = pC$ is equivalent (by uniqueness lemma) to:

$$a_0 - b_0 = pC_0,$$

$$a_1 - b_1 = pC_1,$$

$$a_2 - b_2 = pC_2,$$

$$a_3 - b_3 = pC_3,$$

This implies that the congruence $A \equiv B \pmod{p\mathbb{Z}[\zeta]}$ is equivalent to $a_i \equiv b_i \pmod{p}$ for $i = 0, 1, 2, 3$

Corollary 2.23. $1 \not\equiv -1 \pmod{p\mathbb{Z}[\zeta]}$ if p is an odd prime.

This means that to calculate $\left(\frac{2}{p}\right)$ it is enough to calculate its congruency class mod $(p\mathbb{Z}[\zeta])$

The uniqueness lemma is implied by a more general result:

2.6.1 General Uniqueness Lemma

Let $m \in \mathbb{Z}[X]$ be monic and irreducible over \mathbb{Q} of degree d . If $\alpha \in \mathbb{C}$ is a root of m , then every element of $\mathbb{Z}[\alpha]$ can be written uniquely as $a_0 + a_1\alpha + \dots + a_{d-1}\alpha^{d-1}$ with $a_i \in \mathbb{Z}$.

The uniqueness lemma for $\mathbb{Z}[\zeta]$ follows because ζ is a root of $m(x) = \Phi_8(x) = x^4 + 1$. It is proved in (7202 Groups & Rings) that $x^4 + 1$ is irreducible over \mathbb{Q} .

Proof. (General Uniqueness Lemma)

Let $A \in \mathbb{Z}[\alpha]$ and $m(\alpha) = 0$

Existence: $A = f(\alpha)$ for some $f \in \mathbb{Z}[X]$

divide f by m with remainder, $f = q * m + r$ $\deg(r) < \deg(m) < d$

$$\therefore f(\alpha) = q(\alpha)m(\alpha) + r(\alpha)$$

$$\therefore A = r(\alpha)$$

Uniqueness: Suppose $A = f(\alpha) = g(\alpha)$ ($f \neq g$) where f & g both have degree $< d$

$$\therefore h(\alpha) = 0 \text{ where } h = f - g \text{ } (\neq 0)$$

m is irreducible over \mathbb{Q} and has a bigger degree than h

$$\therefore m \nmid h \text{ in } \mathbb{Q}[x], \text{ so } m \text{ and } h \text{ are coprime in } \mathbb{Q}[x]$$

$\exists a, b \in \mathbb{Q}[x]$ such that :

$$1 = am + bh = a(\alpha)m(\alpha) + b(\alpha)h(\alpha) = 0$$

$$m(\alpha) = 0 \quad h(\alpha) = 0$$

$$\implies 1 = 0$$

$$\implies f = g$$

□

Lemma 2.24. *In any ring R with any prime p*

$$(x + y)^p \equiv x^p + y^p \text{ } (pR) \text{ for any } x, y \in R$$

Proof. Sufficient to show that each binomial coefficient:

$$c = \frac{p!}{i!(p-i)!}$$

$i = 1, 2, \dots, p-1$ is a multiple of p

$$i!(p-i)! \not\equiv 0 \text{ } (p) \implies \in \mathbb{F}_p^\times$$

□

Proof. 2nd Nebensatz

Let p be an odd prime and let $G = \zeta + \zeta^{-1} = \sqrt{2}$. We'll calculate $G^p \bmod (p\mathbb{Z}[\zeta])$ in two ways.

First Calculation:

$$\begin{aligned} G^p &= (\zeta + \zeta^{-1})^p \\ &= \zeta^p + \zeta^{-p} \bmod (p\mathbb{Z}[\zeta]) \text{ by the lemma} \end{aligned}$$

Since $\zeta^8 = 1$ this only depends p modulo 8 if $p \equiv \pm 1(8)$ then,

$$G^p = \zeta + \zeta^{-1} \equiv G \bmod (p\mathbb{Z}[\zeta])$$

If $p \equiv \pm 3(8)$ then,

$$G^p \equiv \zeta^3 + \zeta^{-3} \equiv -G \bmod (p\mathbb{Z}[\zeta])$$

So in summary,

$$G^p \equiv (-1)^{\frac{p^2-1}{8}} G \bmod (p\mathbb{Z}[\zeta])$$

Second Calculation:

Since $G^2 = 2$,

$$\begin{aligned} G^p &= G * 2^{\frac{p^2-1}{2}} \\ &= G * \left(\frac{2}{p}\right) \bmod (p\mathbb{Z}[\zeta]) \text{ by Euler's criterion} \end{aligned}$$

Comparing the results of these two calculations we get:

$$\left(\frac{2}{p}\right)G = (-1)^{\frac{p^2-1}{8}} G \bmod (p\mathbb{Z}[\zeta])$$

Note $G^2 * \frac{p+1}{2} \equiv 1 \bmod (p\mathbb{Z}[\zeta])$, i.e. G is invertible modulo $p\mathbb{Z}[\zeta]$ with inverse $G * \frac{p+1}{2}$

$$\implies \left(\frac{2}{p}\right) \equiv (-1)^{\frac{p^2-1}{8}} \bmod (p\mathbb{Z}[\zeta])$$

Since $1 \equiv -1 \bmod (p\mathbb{Z}[\zeta])$,

$$\left(\frac{2}{p}\right) = (-1)^{\frac{p^2-1}{8}}$$

□

The proof of the 2nd Nebensatz worked because $\sqrt{2} \in \mathbb{Z}[\zeta]$
To prove the quadratic reciprocity law, we'll show that $\sqrt{\pm p}$ is in another cyclotomic ring

Let $\zeta_p = e^{\frac{2\pi i}{p}}$, a primitive p^{th} root of unity. We'll work in the ring modulo $q\mathbb{Z}[\zeta]$.

Defintion 2.25. The p^{th} Gauss sum (where p is an odd prime):

$$G(p) = \sum_{a=1}^{p-1} \left(\frac{a}{p}\right) \zeta_p^a \in \mathbb{Z}[\zeta_p]$$

Lemma 2.26. $G(p)^2 = (-1)^{\frac{p-1}{2}}$

Proof.

$$\begin{aligned} G(p)^2 &= \left(\sum_{a=1}^{p-1} \left(\frac{a}{p}\right) \zeta_p^a \right) \left(\sum_{b=1}^{p-1} \left(\frac{b}{p}\right) \zeta_p^b \right) \\ &= \sum_{a,b \in \mathbb{F}_p^\times} \left(\frac{a}{p}\right) \left(\frac{b}{p}\right) \zeta_p^a \zeta_p^b \\ &= \sum_{a,b \in \mathbb{F}_p^\times} \left(\frac{ab}{p}\right) \zeta_p^{a+b} \end{aligned}$$

Let $c \equiv a^{-1}b \pmod{p}$, as b runs through \mathbb{F}_p^\times , so does c

$$\begin{aligned} &= \sum_{a,c \in \mathbb{F}_p^\times} \left(\frac{a^2}{p}\right) \zeta_p^{a+ac} \\ &= \sum_{c \in \mathbb{F}_p^\times} \left(\frac{c}{p}\right) \left(\sum_{a=1}^{p-1} (\zeta_p^{1+c})^a \right) \end{aligned}$$

Note the second summation is a geometric progression. Recall that,

$$\sum_{i=1}^{p-1} r^i = \begin{cases} \frac{r^p - 1}{r - 1} & r \neq 1 \\ p - 1 & r = 1 \end{cases}$$

Summing the geometric progression:

$$\begin{aligned} \sum_{a=1}^{p-1} (\zeta_p^{1+c})^a &= \begin{cases} \frac{(\zeta_p^{1+c})^p - \zeta_p^{1+c}}{\zeta_p^{1+c} - 1} & \text{if } c \not\equiv 1 \pmod{p} \\ p - 1 & \text{if } c \equiv 1 \pmod{p} \end{cases} \\ &= \begin{cases} -1 & c \not\equiv -1 \pmod{p} \\ p - 1 & c \equiv -1 \pmod{p} \end{cases} \end{aligned}$$

$$\therefore G(p)^2 = \sum_{c \in \mathbb{F}_p^\times} \left(\frac{c}{p}\right)(-1) + p\left(\frac{-1}{p}\right)$$

$\sum_{c \in \mathbb{F}_p^\times} \left(\frac{c}{p}\right)(-1) = 0$ since there are $\frac{p-1}{2}$ quadratic residues and quadratic non-residues.

$$\begin{aligned} &= p\left(\frac{-1}{p}\right) \\ &= (-1)^{\frac{p-1}{2}} p \end{aligned} \quad \text{by the 1}^{st} \text{ Nebensatz}$$

□

2.6.2 Uniqueness Lemma for $\mathbb{Z}[\zeta_p]$

Every element $A \in \mathbb{Z}[\zeta_p]$ can be written uniquely as:

$$A = a_0 + a_1\zeta + \cdots + a_{p-2}\zeta^{p-2} \quad \text{with } a_i \in \mathbb{Z}$$

This is because ζ_p is a root of $m(x) = \Phi_p(x) = 1 + x + \cdots + x^{p-1}$. It's proved in 7202 that Φ_p is irreducible over \mathbb{Q} .

Proof. Quadratic Reciprocity law

We'll calculate $G(p)^q$ ($q\mathbb{Z}[\zeta_p]$) in two ways.

First Calculation:

$$\begin{aligned} G(p)^q &= \left(\sum_{a=1}^{p-1} \left(\frac{a}{p}\right) \zeta_p^a \right)^q \\ &= \sum_{a=1}^{p-1} \left(\left(\frac{a}{p}\right) \zeta_p^a \right)^q \quad (p\mathbb{Z}[\zeta]) \end{aligned}$$

Since q is odd, $\left(\frac{a}{p}\right)^q = \left(\frac{a}{p}\right)$

$$G(p)^q \equiv \sum_{a \in \mathbb{F}_p^\times} \left(\frac{a}{p}\right) \zeta_p^{aq}$$

Let $b \equiv aq \pmod{p}$, and as a runs through \mathbb{F}_p^\times so does b

$$\begin{aligned} G(p)^q &\equiv \sum_{b \in \mathbb{F}_p^\times} \left(\frac{bq^{-1}}{p}\right) \zeta_p^b \\ &= \left(\frac{q^{-1}}{p}\right) \sum_{b \in \mathbb{F}_p^\times} \left(\frac{b}{p}\right) \zeta_p^b \end{aligned}$$

Note that $G(p) = \sum_{b \in \mathbb{F}_p^\times} \left(\frac{b}{p}\right) \zeta_p^b$ which implies,

$$\begin{aligned} G(p)^q &\equiv \left(\frac{q^{-1}}{p}\right) G(p) \pmod{q\mathbb{Z}[\zeta_p]} \\ &\equiv \left(\frac{q}{p}\right) G(p) \pmod{q\mathbb{Z}[\zeta_p]} \end{aligned}$$

Second Calculation:

Since $G(p)^2 = (-1)^{\frac{p-1}{2}} p$,

$$\begin{aligned} G(p)^q &= G(p) \left((-1)^{\frac{p-1}{2}} p \right)^{\frac{q-1}{2}} \\ &= G(p) (-1)^{\frac{(p-1)(q-1)}{4}} p^{\frac{q-1}{2}} \\ \therefore G(p)^q &\equiv G(p) (-1)^{\frac{(p-1)(q-1)}{4}} \left(\frac{p}{q}\right) \pmod{q\mathbb{Z}[\zeta_p]} \quad \text{by Euler's criterion} \end{aligned}$$

Comparing the two results we get:

$$\left(\frac{q}{p}\right) G(p) \equiv (-1)^{\frac{(p-1)(q-1)}{4}} \left(\frac{p}{q}\right) G(p) \pmod{q\mathbb{Z}[\zeta_p]}$$

We need to check that $G(p)$ is invertible modulo $q\mathbb{Z}[\zeta_p]$,

$G(p)^2 = \pm p$, which is invertible modulo q

$G(p)$ has inverse $G(p) * (\pm p)^{-1} \pmod{q\mathbb{Z}[\zeta_p]}$

$$\therefore \left(\frac{p}{q}\right) \equiv (-1)^{\frac{(p-1)(q-1)}{4}} \left(\frac{p}{q}\right) \pmod{q\mathbb{Z}[\zeta_p]}$$

Since $1 \equiv -1 \pmod{q\mathbb{Z}[\zeta_p]}$, it follows that $\left(\frac{q}{p}\right) = (-1)^{\frac{(p-1)(q-1)}{4}} \left(\frac{p}{q}\right)$ □

3 P-adic Number theory

This means methods for congruences modulo p^n , p prime and n large.

If we want to solve $f(x) = 0$, $x \in \mathbb{R}$ we can use the Newton-Raphson method:

- Begin with an "approximate solution" a_0
- Define a sequence recursively $a_{n+1} = a_n - \frac{f(a_n)}{f'(a_n)}$

Very often a_n converge to a limit a and $f(a) = 0$.

We can use the same method in number theory for solving congruences. Suppose $f(x)$ is a polynomial with coefficients in \mathbb{Z} and we want to solve $f(x) \equiv 0 \pmod{p^N}$ (p prime, n large)

We can try this:

- Find a solution a_0 to $f(a_0) \equiv 0 \pmod{p^r}$ where r is small
- Define a recursive sequence $a_{n+1} = a_n - \frac{f(a_n)}{f'(a_n)}$

If n is large enough, then often $f(a_n) \equiv 0 \pmod{p^N}$

E.g. Let $f(x) = x^2 + 2$, $p = 3$

Suppose we want to solve $x^2 + 2 \equiv 0 \pmod{3^N}$

Let $a_0 = 1$: $f(a_0) = 1^2 + 2 = 3 \equiv 0 \pmod{3}$

Define the sequence a_n by $a_{n+1} = a_n - \frac{f(a_n)}{f'(a_n)} = a_n - \frac{a_n^2 + 2}{2a_n} = \frac{a_n}{2} - \frac{1}{a_n}$

$$\begin{aligned} a_0 &= 1 \\ a_1 &= \frac{1}{2} - 1 = \frac{-1}{2} \\ a_2 &= \frac{-1}{4} + 2 = \frac{7}{4} \end{aligned}$$

It turns out that $\frac{-1}{2}$ is a solution mod 9 $\implies -1 * 2^{-1} \pmod{9}$
 $\frac{7}{4}$ is a solution mod 81 $\implies 7 * 4^{-1} \pmod{81}$

$$2^{-1} \equiv 5 \pmod{9} \implies a_1 \equiv 4 \pmod{9}$$

$$4^{-1} \equiv -20 \pmod{81} \implies a_2 = \frac{7}{4} \equiv -140 \equiv 22 \pmod{81}$$

a_3 would be a solution mod 3^8 .

In this example, we're reducing rational numbers mod p^n not just integers. If $\frac{a}{b}$ is a rational number then we can reduce this modulo p^n as long as b is invertible mod p^n , i.e. when b is not a multiple of p . We'll write:

$$\mathbb{Z}_{(p)} = \left\{ \frac{a}{b} : a, b \in \mathbb{Z}, p \nmid b \right\}$$

$\mathbb{Z}_{(p)}$ is closed under $+$, $*$, so $\mathbb{Z}_{(p)}$ is a ring contained in \mathbb{Q} containing \mathbb{Z} . This is called the "local ring of p " and is the set of rational number which can be reduced modulo p^n ($\forall n$)

Defintion 3.1. If p is a prime number and $n \in \mathbb{Z}$, then the valuation of n , at p is:

$$V_p(n) = \begin{cases} \max\{a : p^a | n\} & n \neq 0 \\ \infty & n = 0 \end{cases}$$

A simple statement that can be made is, $V_p(nm) = V_p(n) + V_p(m)$. We can also extend V_p to a function on \mathbb{Q} , $V_p(\frac{n}{m}) = V_p(n) - V_p(m)$.

With this notation:

$$Z_{(p)} = \{x \in \mathbb{Q} : V_p(x) \geq 0\}$$

$$x \equiv y \pmod{p^a} \iff V_p(x - y) \geq a$$

E.g

$$V_2(\frac{7}{12}) = -2 \quad V_2(\frac{7}{12}) = -1 \quad V_5(\frac{7}{12}) = 0 \quad V_7(\frac{-7}{12}) = +1$$

3.1 Hensel's Lemma

Let p be a prime number. Let $f \in \mathbb{Z}_{(p)}[x]$ and $a_0 \in \mathbb{Z}_{(p)}$ such that $f(a_0) \equiv 0 \pmod{p^{2c+1}}$ where $c = V_p(f'(a_0))$.

Then if we define $a_{n+1} = a_n - \frac{f(a_n)}{f'(a_n)}$ then $a_n \in \mathbb{Z}_{(p)}$ and $f(a_n) \equiv 0 \pmod{p^{2c+2^n}}$

Proof. We'll prove the following by induction on n

1. $a_n \in \mathbb{Z}_{(p)}$ and $a_n \equiv a_0 \pmod{p^{c+1}}$
2. $V_p(f'(a_n)) = c$
3. $f(a_n) \equiv 0 \pmod{p^{2c+2^n}}$

If $n = 0$ then the statements 1,2,3 are all true for a by assumption. Now assume 1,2,3 for a_n , we'll prove them for a_{n+1}

Let $a_{n+1} = a_n - \delta$ where $\delta = \frac{f(a_n)}{f'(a_n)}$

1:

$$\begin{aligned} V_p(\delta) &= V_p(f(a_n)) - V_p(f'(a_n)) \\ &= c \end{aligned}$$

by **2:**

$$\geq 2c + 2^n$$

by **3:**

$$V_p(\delta) \geq 2c + 2^n - c$$

$$V_p(\delta) \geq c + 2^n$$

(*)

By (*)

$$V_p(\delta) \geq 0 \implies \delta \in \mathbb{Z}_{(p)}$$

$$\therefore a_{n+1} = a_n - \delta \in \mathbb{Z}_{(p)}$$

By (*)

$$V_p \geq c + 1 \implies \delta \equiv 0 \pmod{p^{c+1}}$$

$$a_{n+1} \equiv a_n \pmod{p^{c+1}}$$

$$\equiv a_0 \pmod{p^{c+1}}$$

by 1

2: We've shown that $a_{n+1} \equiv a_0 \pmod{p^{c+1}}$

$$\therefore f'(a_{n+1}) \equiv f'(a_0) \pmod{p^{c+1}}$$

$$\not\equiv 0$$

$$\text{because } V_p(f'(a_0)) = c$$

$$\text{also } f'(a_{n+1}) \equiv f'(a_0) \pmod{p^c}$$

$$\equiv 0 \pmod{p^c}$$

$$\text{because } V_p(f'(a_0)) = c \pmod{p^c}$$

$$\therefore V_p(f'(a_{n+1})) = c$$

3: Must show that $f(a_{n+1}) \equiv 0 \pmod{p^{2c+2^{n+1}}}$

$$a_{n+1} = a_n - \delta$$

$$a_{n+1}^r = (a_n - \delta)^r$$

$$= a_n^r - r a_n^{r-1} \delta + \text{multiples of } \delta^2$$

By (*):

$$V_p(\delta) \geq c + 2^n$$

$$\therefore V_p(\delta^2) \geq 2c + 2^{n+1}$$

$$\therefore \delta^2 \equiv 0 \pmod{p^{2c+2^{n+1}}}$$

This implies $a_{n+1}^r \equiv a_n^r - r a_n^{r-1} \delta \pmod{p^{2c+2^{n+1}}}$

Suppose $f(x) = \sum c_r * x$. Substituting a_{n+1} , we get:

$$\begin{aligned} f(a_{n+1}) &= \sum c_r (a_n^r - r a_n^{r-1} \delta) \pmod{p^{2c+2^{n+1}}} \\ &= \sum c_r a_n^r - \left(\sum r c_r a_n^{r-1} \right) \delta \pmod{p^{2c+2^{n+1}}} \\ &= f(a_n) - f'(a_n) * \frac{f(a_n)}{f'(a_n)} \equiv 0 \pmod{p^{2c+2^{n+1}}} \end{aligned}$$

□

E.g. $f(x) = x^3 + x + 1$, $p = 3$

Find a root of $f \bmod 81$

Note that $f'(x) = 3x^2 + 1$ and $f(1) = 3 \equiv 0$

Try $a_0 = 1$

$$\begin{aligned}c &= V_3(f'(a_0)) \\&= V_3(4) \\&= 0\end{aligned}$$

$3^{2c+1} = 3$ and a_0 is a root of f modulo 3

$\therefore a_0 = 1$ satisfies the conditions of Hensel's lemma.

$$\begin{aligned}a_1 &= 1 - \frac{a_0}{f'(a_0)} \\&= 1 - \frac{3}{4}\end{aligned}$$

It is sufficient to work out $a_1 \bmod 9$

$$4^{-1} \equiv 1 \pmod{3} \qquad \frac{3}{4} \equiv 3 * 1 \pmod{9} \qquad a_1 \equiv -2 \pmod{9}$$

Check

$$\begin{aligned}f(a_1) &\equiv (-2)^3 + (-2) + 1 \\f(2) &= -9 \equiv 0 \pmod{9}\end{aligned}$$

$$\begin{aligned}a_2 &= -2 - \frac{f(-2)}{f'(-2)} \\&= -2 - \frac{-9}{13}\end{aligned}$$

This should be a root of f modulo 81.

$$13^{-1} \equiv -2 \pmod{9}$$

$$\implies \frac{9}{13} \equiv -18$$

$$a_2 \equiv -2 - 18 \equiv -20$$

$$\begin{aligned}\text{Check } f(a_2) &= (-20)^3 - 20 + 1 \equiv -8000 - 19 \\&= -8019 \\&= -81 * 99 \\&= 0 \pmod{81}\end{aligned}$$

3.2 Quadratic Congruences

We'll see how to find out whether $x^2 \equiv b \pmod{n}$ has solutions.

Suppose $n = p_1^{a_1} \dots p_r^{a_r}$ (p_i distinct primes). There are solutions modulo $n \iff \forall i$, there are solutions modulo $p_i^{a_i}$ by the Chinese Remainder Theorem.

Proposition 3.2. *Suppose p is an odd prime not dividing b . If $x^2 \equiv b \pmod{p}$ has solutions then $x^2 \equiv b \pmod{p^r}$ has solutions for all r*

Proof. Suppose there is a solution a_0 modulo p , i.e. $a_0^2 \equiv b \pmod{p}$

Let $f(x) = x^2 - b$. We'll check that a_0 satisfies the conditions of Hensel's lemma.

$$\begin{aligned} c &= V_p(f'(a_0)) \\ &= V_p(2a_0) \quad \text{and since } p \neq 2 \\ \implies c &= V_p(a_0) \end{aligned}$$

Also since $p \nmid b$, we know $p \nmid a_0$:

$$\begin{aligned} \therefore c &= 0 \\ \therefore f(a_0) &\equiv 0 \pmod{p^{2c+1}} \implies a_0 \text{ satisfies the conditions of Hensel's lemma} \\ \therefore &\text{ We have roots of } f \text{ modulo all powers of } p \end{aligned}$$

□

Remark

Suppose we want a root of f modulo p^{13}

Choose n so that $2c + 2^n \geq 13$

$$f(a_n) \equiv 0 \pmod{p^{2c+2^n}} \implies f(a_n) \equiv 0 \pmod{p^{13}}$$

The proposition would be false if we allowed $p = 2$

E.g. Let $b = 3$

x	$x^2 \pmod{4}$
0	0
1	1
2	0
3	1

$$x^2 \equiv 3 \pmod{2} \text{ has a solution}$$

$$x^2 \equiv 3 \pmod{4} \text{ has no solutions}$$

if $b = 5$

x	$x^2 \pmod{8}$
0	0
± 1	1
± 2	4
± 3	1
± 4	0

$$x^2 \equiv 5 \pmod{2} \text{ has a solution}$$

$$x^2 \equiv 5 \pmod{4} \text{ has solutions}$$

$$x^2 \equiv 5 \pmod{8} \text{ has no solutions}$$

Proposition 3.3. Suppose b is odd. If $x^2 \equiv b \pmod{8}$ has solutions then $x^2 \equiv b \pmod{2^r}$ has solutions for all r

Proof. Suppose $a_0 \equiv b \pmod{8}$, this implies a_0 is odd.

Let $f(x) = x^2 - b$

$\therefore c = V_2(f'(a_0)) = V_2(2a_0) = 1$ because a_0 is odd

$\therefore 2^{2c+1} = 8$

$\therefore a_0$ is a root of f modulo p^{2c+1}

By Hensel's lemma, there are solutions modulo all powers of 2.

□

E.g. For which n does the congruence $x^2 \equiv 5 \pmod{5}$ have solutions?

First consider the case $n \equiv p^r$ (p prime)

If $p \neq 2, 5$ then by the first proposition, there are solutions $p^n \iff \left(\frac{5}{p}\right) = 1$

$\left(\frac{5}{p}\right) = +\left(\frac{p}{5}\right)$ depends on $p \pmod{5}$

x	s	
1	1	(different x)
2	-1	The congruence $x^2 \equiv 5 \pmod{p}$ has solutions
3	-1	$\iff p \equiv 1, 4 \pmod{5}$ (in the cases $p \neq 2, 5$)
4	1	

For $p = 2$, $x^2 \equiv 5 \pmod{2}$ has a solution, $x = 1$

$x^2 \equiv 5 \pmod{4}$ has a solution, $x = 1$

But the only odd square mod 8 is 1. So $x^2 \equiv 5 \pmod{8}$ has no solutions.

\therefore no solutions mod 2^n if $n \geq 3$

For $p = 5$ $x^2 \equiv 5 \pmod{5}$ has solutions, here's how we check. Assume:

$$x^2 \equiv 5 \pmod{25}$$

$$\therefore x^2 \equiv 0 \pmod{5}$$

$$\text{So } 5|x^2$$

$$\text{So } 5|x$$

$$\therefore x^2 \equiv 0 \pmod{25} \quad \nexists$$

So there are solution modulo n if $n = 2^a * 5^b * \prod p_i^{c_i}$ where $a \leq 2, b \leq 1, p_i \equiv 1 \pmod{5}, c_i \in \mathbb{N}$

E.g. For which n does $x^2 \equiv -7 \pmod{n}$ have solutions?

Assume p is a prime $\neq 2, 7$

$$\begin{aligned}
\left(\frac{-7}{p}\right) &= \left(\frac{-1}{p}\right)\left(\frac{7}{p}\right) \\
&= (-1)^{\frac{p-1}{2}} (-1)^{\frac{(7-1)(p-1)}{4}} \left(\frac{p}{7}\right) \\
&= (-1)^{\frac{p-1}{2} + \frac{3(p-1)}{2}} \left(\frac{p}{7}\right) \\
&= (+1)\left(\frac{p}{7}\right) \text{ depends on } p \bmod 7
\end{aligned}$$

x	$\left(\frac{x}{7}\right)$
1	1
2	1
3	-1
4	1
5	-1
6	-1

$$\begin{aligned}
3^2 &\equiv 9 \equiv 2 \pmod{7} \\
x^2 &\equiv -7 \pmod{p^r} \text{ has solutions} \\
&\implies p \equiv 1, 2, 4 \pmod{7}
\end{aligned}$$

For $p = 2$: $-7 \equiv 1 \pmod{8}$ so -7 is a square modulo 8 by the proposition.
 $x^2 \equiv -7 \pmod{2^r}$ has solutions for all r .

For $p = 7$: $x^2 \equiv -7 \pmod{7}$ has a solution $x = 0$ but $x^2 \equiv -7 \pmod{7^2}$ has no solutions. Suppose

$$\begin{aligned}
x^2 &\equiv -7 \pmod{7^2} \\
\therefore x^2 &\equiv 0 \pmod{7} \\
\therefore 7|x^2 \\
\implies 7|x \\
\implies x^2 &\equiv 0 \pmod{49} \quad \nexists
\end{aligned}$$

So $x^2 \equiv -7 \pmod{n}$ has solutions $\iff n = 7^a * \prod p_i^{b_i}$ where $a \leq 1$, $p_i \equiv 1, 2, 4 \pmod{7}$, $b_i \in \mathbb{N}$

3.3 P-adic congruence

Suppose we have a series $\sum_{n=1}^{\infty} x_n$ for $x_n \in \mathbb{Z}_{(p)}$. We'll say that the series converges **p-adically** if for every a , there are only finitely many terms x_n with $x_n \not\equiv 0 \pmod{p^a}$. We can add up the series in \mathbb{Z}/p^a because only finitely many terms are non zero.

Lemma 3.4. $\sum x_n$ converges *p-adically* $\iff V_p(x_n) \rightarrow \infty$

Proof. If $V_p(x_n) \rightarrow \infty$ then for n significantly large, $V_p(x_n) \geq a$, i.e., $x_n \equiv 0 \pmod{p^a}$

□

E.g. $p=3$

$$(1 + 3x)^{\frac{1}{2}} = 1 + \frac{1}{2}(3x) + \frac{\left(\frac{1}{2}\right)\left(\frac{-1}{2}\right)(3x)^2}{2!} + \frac{\left(\frac{1}{2}\right)\left(\frac{-1}{2}\right)\left(\frac{-3}{2}\right)(3x)^3}{3!}$$

if $x \in \mathbb{Z}_{(3)}$ then this series converge 3-adically.

$$\begin{aligned}
(1 + 3x)^{\frac{1}{2}} &\equiv 1 \quad (3) \\
&\equiv 1 + \frac{3x}{2} \quad (9) \\
&\equiv 1 + \frac{3x}{2} + \frac{9}{8}x^2 \quad (27) \\
&\equiv 1 + \frac{3x}{2} + \frac{9}{8}x^2 + \frac{27}{16}x^3 \quad (27)
\end{aligned}$$

We can write these polynomials with integer coefficients.

$$\begin{aligned}
(1 + 3x)^{\frac{1}{2}} &\equiv 1 + 15x + 9x^2 \quad (27) \\
&\equiv 1 + 42x + 9x^2 + 27x^3 \quad (81)
\end{aligned}$$

Important point; these polynomials play the same role in number theory $(1 + 3x)^{\frac{1}{2}}$ does in analysis $\sqrt{1 + 3x}$

E.g.

$$\begin{aligned}
(1 + 15x + 9x^2)^2 &= 1 + (30)x + (18 + 15^2)x^2 + (2 * 9 * 15)x^3 + (81)x^4 \\
&\equiv 1 + 3x \quad (27)
\end{aligned}$$

E.g. Find a square root of 7 in $\mathbb{Z}/81$

$$\begin{aligned}
7^{\frac{1}{2}} &= (1 + 3 * 2)^{\frac{1}{2}} \\
&\equiv 1 + 42 * 2 + 9 * 2^2 + 27 * 2^3 \quad (81) \\
&\equiv 1 + 84 + 36 - 27 \quad (81) \\
&\equiv 13 \quad (81)
\end{aligned}$$

Check $13^2 = 169 \equiv 7 \quad (81)$

This works because of a result called the power series trick.

Notation We'll write $\mathbb{Z}_{(p)}[[x]]$ for the set of power series in x with coefficient in $\mathbb{Z}_{(p)}$.

$\mathbb{Z}_{(p)}[[x]]$ is a ring with addition and multiplication of power series as operations. We can

often compose two power series $f, g \in \mathbb{Z}_{(p)}[[x]]$ to get a new power series $f \circ g$.

$(f \circ g)(x) = f(g(x))$.

We can define $f \circ g$ as long as either f is a polynomial or g has zero constant term. Suppose

$$\begin{aligned}
f(x) &= \sum_{n=0}^{\infty} a_n x^n \\
g(x) &= \sum_{n=0}^{\infty} b_n x^n
\end{aligned}$$

We'll see that $f \circ g$ is a power series

$$\begin{aligned} f(g(x)) &= \sum_{n=0}^{\infty} a_n \left(\sum_{m=1}^{\infty} b_m x^m \right)^n \\ &= \sum_{n=0}^{\infty} a_n \sum_{m_1=0}^{\infty} \sum_{m_2=0}^{\infty} \cdots \sum_{m_n=0}^{\infty} b_{m_1} \cdots b_{m_n} x^{m_1 + \cdots + m_n} \end{aligned}$$

so $f(g(x)) = \sum c_d x^d$ where

$$c_d = \underbrace{\sum_{m_1, \dots, m_n=1}^{\infty} a_n b_{m_1} \cdots b_{m_n}}_{\text{finite sum in } \mathbb{Z}_{(p)}}$$

Note $f \circ g$ is not defined otherwise.

E.g

$$\begin{aligned} f(x) &= 1 + x + x^2 + \dots \\ g(x) &= 1 + x \end{aligned}$$

$$\implies f(g(x)) = 1 + (1 + x) + (1 + x)^2$$

This has constant term $1 + 1 + 1 + 1 + \dots$, so $f \circ g$ is not defined.

3.4 Power Series Trick

Suppose f, g, h are power series with coefficients in $\mathbb{Z}_{(p)}$. Assume either f is a polynomial or g has no constant term. Also assume:

- For small real numbers x , $f(x), g(x), h(x)$ converge and $f(g(x)) = h(x)$
- For all $x \in \mathbb{Z}_{(p)}$, $f(x), g(x)$ and $h(x)$ converge p-adically

Then for all $x \in \mathbb{Z}_{(p)}$, $f(g(x)) \equiv h(x) \pmod{p^n}$

In the example $f(x) = x^2, g(x) = (1 + 3x)^{\frac{1}{2}}, h(x) = 1 + 3x$.

For small real x , $f(g(x)) = h(x)$, so as long as we know that $g(x)$ converges 3-adically ($\forall x \in \mathbb{Z}_{(3)}$) the power series trick implies $g(x)^2 \equiv 1 + 3x \pmod{3^n}$

How do we check for p-adic convergence?

Lemma 3.5. $\sum x_n$ converge p-adically if and only if $V_p(x_n) \rightarrow \infty$

We need a way of calculating valuations of n^t term of a square.

Proposition 3.6. $V_p(n!) = \lfloor \frac{n}{p} \rfloor + \lfloor \frac{n}{p^2} \rfloor + \lfloor \frac{n}{p^3} \rfloor + \cdots \leq \frac{n}{p-1}$

We'll prove this later, first use the properties to show that $(1 + 3x)^{\frac{1}{2}}$ converge 3-adically for all $x \in \mathbb{Z}_{(3)}$

$$(1 + 3x)^{\frac{1}{2}} = 1 + \frac{1}{2}(3x) + \frac{(\frac{1}{2})(\frac{-1}{2})(3x)}{2!} + \dots$$

$$n^{th} \text{ term} = \frac{(\frac{1}{2})(\frac{1}{2} - 1)(\frac{1}{2} - 2) \dots (\frac{1}{2} - n + 1)}{n!} (3x)^n$$

$$\begin{aligned} V_3(n^{th} \text{ term}) &= V_3\left(\frac{1}{2}(\frac{1}{2} - 1) \dots (\frac{1}{2} - n + 1)\right) - V_3(n!) + V_3((3x)^n) \\ &\geq 0 - \frac{n}{3-1} + n \\ &\geq \frac{n}{2} \rightarrow \infty \text{ as } n \rightarrow \infty \\ &\implies \text{series converges 3-adically} \end{aligned}$$

E.g. Assume p is an odd prime

Let $\exp(px) = 1 + px + \frac{(px)^2}{2!} + \frac{(px)^3}{3!} + \dots$

We'll see that this converges for all $x \in \mathbb{Z}_{(p)}$

$$n^{th} \text{ term} = \frac{(px)^n}{n!}$$

$$\begin{aligned} V_p(n^{th} \text{ term}) &= V_p((px)^n) - V_p(n!) \\ &= (n * V_p(p)) + (n * V_p(x)) - V_p(n!) \\ &\quad 1 \qquad \qquad \geq 0 \qquad \leq \frac{n}{p-1} \\ &\geq n - \frac{n}{p-1} \\ &\geq \left(\frac{p-2}{p-1}\right)n \rightarrow \infty \text{ for } p \neq 2 \end{aligned}$$

E.g. $\log(1 + px)$ converges p-adically for all $x \in \mathbb{Z}_{(p)}$

$$\begin{aligned} V_p\left(\pm \frac{(px)^n}{n}\right) &= V_p((px)^n) - V_p(n) \\ &\quad nV_p(px) < V_p(n!) \\ &\geq n - \frac{n}{p-1} \\ &\geq \left(\frac{p-2}{p-1}\right)n \rightarrow \infty \text{ if } p \neq 2 \end{aligned}$$

Remark A quick way to remember the series for $\log(1 + px)$

$$\frac{1}{1-x} = 1 + x + x^2 + \dots \quad \text{geometric series}$$

$$\frac{1}{1+x} = 1 - x + x^2 - x^3 + \dots$$

$$\log(1+x) = x - \frac{x^2}{2} + \frac{x^3}{3} - \frac{x^4}{4} + \dots$$

$$\log(1+px) = x - \frac{(px)^2}{2} + \frac{(px)^3}{3} - \frac{(px)^4}{4} + \dots$$

Proof. Calculating $V_p(n!)$

$$n! = 1 * 2 * \dots * n$$

$$V_p(n) = \sum_{i=1}^n V_p(i) \quad (*)$$

The number of i between 1 & n which are multiples of p is $\lfloor \frac{n}{p} \rfloor$.

There are $\frac{n}{p^2}$ values of i which are multiples of p^2 , etc.

$\lfloor \frac{n}{p} \rfloor - \lfloor \frac{n}{p^2} \rfloor$ values of i are multiples of p , but not of p^2 , i.e. $V_p(i) = 1$

So $\lfloor \frac{n}{p} \rfloor - \lfloor \frac{n}{p^2} \rfloor$ terms in the sum $(*)$ are equal to 1.

Similarly $\lfloor \frac{n}{p^2} \rfloor - \lfloor \frac{n}{p^3} \rfloor$ terms in the sum $(*)$ are equal to 2.

In general there are exactly $\lfloor \frac{n}{p^a} \rfloor - \lfloor \frac{n}{p^{a+1}} \rfloor$ terms in $(*)$ which are equal to a

$$\therefore V_p(n!) = 1 * \text{no of terms equal to 1} + 2 * \text{number of terms equal to 2} + \dots$$

$$\begin{aligned} V_p(n!) &= 1 * (\lfloor \frac{n}{p} \rfloor - \lfloor \frac{n}{p^2} \rfloor) \\ &\quad + 2 * (\lfloor \frac{n}{p^2} \rfloor - \lfloor \frac{n}{p^3} \rfloor) \\ &\quad + 3 * (\lfloor \frac{n}{p^3} \rfloor - \lfloor \frac{n}{p^4} \rfloor) \\ &\quad + \dots \\ &= \lfloor \frac{n}{p} \rfloor + (2-1)\lfloor \frac{n}{p^2} \rfloor + (3-2)\lfloor \frac{n}{p^3} \rfloor + \dots \\ &= \lfloor \frac{n}{p} \rfloor + \lfloor \frac{n}{p^2} \rfloor + \lfloor \frac{n}{p^3} \rfloor + \dots \end{aligned}$$

Using this we can prove the upper bound.

$$\begin{aligned}
V_p(n!) &\leq \frac{n}{p} + \frac{n}{p^2} + \dots \\
&\leq \frac{n}{p} \underbrace{\left(1 + \frac{1}{p} + \frac{1}{p^2} + \dots\right)}_{\text{geometric series } \frac{1}{1-\frac{1}{p}}} \\
&\leq \frac{n}{p-1}
\end{aligned}$$

□

3.4.1 P-adic log & exp

Let p be an odd prime. Use the notation

$$p\mathbb{Z}/p^n = \{px : x \in p\mathbb{Z}/p^n\}$$

E.g.

$$3\mathbb{Z}/27 = \{0, 3, 6, 9, 12, 15, 18, 21, 24\}$$

$p\mathbb{Z}/p^n$ is closed under $+$, so it is a subgroup of $(\mathbb{Z}/p^n)^\times$

$$1 + p\mathbb{Z}/p^n = \{1 + px : x \in p\mathbb{Z}/p^n\}$$

E.g.

$$1 + 3\mathbb{Z}/27 = \{1, 4, 7, 10, 13, 16, 19, 22, 25\}$$

$1 + p\mathbb{Z}/p^n$ is closed under $*$, so $1 + p\mathbb{Z}/p^n$ is a subgroup $(\mathbb{Z}/p^n)^\times$. Both subgroups have p^{n-1} elements, but one is additive and the other is multiplicative. But actually there are isomorphic. This isomorphism is exp & log.

Theorem 3.7. *Let p be an odd prime. Then there is an isomorphism:*

$$p\mathbb{Z}/p^n \xleftrightarrow[\exp]{\log} 1 + p\mathbb{Z}/p^n$$

$$px \longmapsto \exp(px)$$

$$1 + px \longmapsto \log(1 + px)$$

E.g. $\mathbb{Z}/27$ ($p = 3$) We'll find the isomorphisms in this case.

$$\exp(3x) \equiv 1 + 3x + \frac{3^2 x^2}{2!} + \frac{3^3 x^3}{3!} \quad (27)$$

$$\equiv 1 + 3x + 18x^2 + 18x^3 \quad (27)$$

$$\log(1 + 3x) \equiv 3x - \frac{(3x)^2}{2} + \frac{(3x)^3}{3} \quad (27)$$

$$\equiv 3x + 9x^2 + 9x^3 \quad (27)$$

Check:

$$\begin{aligned}
 \log(\exp(3x)) &\equiv \log(1 + 3(x + 6x^2 + 6x^3)) \\
 &\equiv 3(x + 6x^2 + 6x^3) + 9(x + 6x^2 + 6x^3)^2 + 9(x + 6x^2 + 6x^3) \\
 &\equiv 3x + 18x^2 + 18x^3 + 9x^2 + 9x^3 \\
 &\equiv 3x
 \end{aligned}$$

Similarly $\exp(\log(1 + 3x)) \equiv 1 + 3x$ (27)

We can use the theorem to solve congruences.

E.g. Solve $7^x \equiv 13$ (27)

7 and 13 are in $1 + 3\mathbb{Z}/27$, so we can take their logarithms.

$$x \log(7) \equiv \log(13)$$

Using the formula for $\log(1 + 3x)$, we get:

$$\begin{aligned}
 \log(7) &= \log(1 + 6) \\
 &\equiv 6 - \frac{6^2}{2} + \frac{6^3}{3} - \frac{6^4}{4} \\
 &\equiv 6 - 18 + 72 \\
 &\equiv 6 \quad (27)
 \end{aligned}$$

$$\begin{aligned}
 \log(13) &\equiv \log(1 + 12) \\
 &\equiv 12 - \frac{12^2}{2} + \frac{12^3}{3} \quad (27) \\
 &\equiv 12 - 72 + 3^2 * 4^3 \quad (27) \\
 &\equiv 12 - 72 + 9 \quad (27) \\
 &\equiv 3 \quad (27)
 \end{aligned}$$

So $7^x \equiv 13$ (27) reduces to:

$$\begin{aligned}
 7^x &\equiv 13 \quad (27) \\
 \implies 6x &\equiv 3 \quad (27) \\
 \implies 2x &\equiv 1 \quad (9) \\
 \implies x &\equiv 5 \quad (9)
 \end{aligned}$$

Proof.

We'll use the power series trick. We've shown that $\log(1 + px), \exp(px)$ converge p-adically for $x \in \mathbb{Z}_{(p)}$ and they converge for small real numbers and for small real x

$$\begin{aligned}\log(\exp(px)) &= px \\ \exp(\log(1 + px)) &= 1 + px\end{aligned}$$

By the power series trick:

$$\begin{aligned}\exp(\log(1 + px)) &\equiv 1 + px \pmod{p^n} \\ \log(\exp(px)) &\equiv px \pmod{p^n}\end{aligned}$$

$\therefore \log$ and \exp are inverse functions, so they are bijective.

Remains to show that $\exp(px + py) \equiv \exp(px) * \exp(py) \pmod{p^n}$

For any $a \in \mathbb{N}$: $\exp(pax) = \exp(px)^a$ for small real x

By the power series trick with:

$$\begin{aligned}f(x) &= x^a \\ g(x) &= \exp(px) \\ h(x) &= \exp(pax) \\ \exp(pax) &\equiv (\exp(px))^a \pmod{p^a}\end{aligned}$$

Take $x = 1$

$$\begin{aligned}\exp(pa) &\equiv \exp(p)^a \pmod{p^a} \\ \therefore \exp(pa + pb) &\equiv \exp(p)^{a+b} \pmod{p^a} \\ &\equiv \exp(p)^a * \exp(p)^b \\ &\equiv \exp(pa) * \exp(pb) \pmod{p^n}\end{aligned}$$

We've proved this when a & b are positive integers, but every element of \mathbb{Z}/p^n can be written as a positive integer. □

3.5 Teichmüller Lifts

Let p be an odd prime. We saw that $(\mathbb{Z}/p^n)^\times$ has a big subgroup $1 + p\mathbb{Z}/p^n$ and we can easily do calculations in the subgroup. Teichmüller lifts is another subgroup.

$$(\mathbb{Z}/p^n)^\times = \text{Teichmüller lifts} * (1 + p\mathbb{Z}/p^n)$$

Let $x \in \mathbb{Z}_{(p)}$ and assume $x \not\equiv 0 \pmod{p}$:

$$x, x^p, x^{p^2}, x^{p^3}, \dots$$

All these terms are constant mod p :

$$\begin{aligned} x^{p-1} &\equiv 1 \pmod{p} \\ x^p &\equiv x \pmod{p} \end{aligned}$$

The sequence is constant mod p^2 , but all terms after the 2^{nd} are constant mod p^2 .

E.g. $p = 3, x = 2$

We'll look at the sequence mod 9:

$$\begin{aligned} 2^3 &\equiv 8 \pmod{9} \\ 2^9 &\equiv 8^3 \equiv 8 \pmod{9} \\ 2^{27} &\equiv 8 \pmod{9} \quad \text{etc} \end{aligned}$$

The sequence is eventually constant modulo p^n

Defintion 3.8. *The Teichüller lift of x modulo p^n is:*

$$T(x) \equiv x^{p^{n-1}} \pmod{p^n}$$

To calculate Teichmüller lifts, we use:

Lemma 3.9. *Suppose $x \equiv y \pmod{p^n}$ then $x^p \equiv y^p \pmod{p^{n+1}}$*

Proof. Let $x \equiv y + p^n \implies$

$$x^p \equiv (y + p^n)^p$$

$$x^p \equiv y^p + py^{p-1}p^n + \text{multiples of } p^{2n}$$

$$x^p \equiv y^p \pmod{p^{n+1}}$$

□

E.g. Calculate $T(12) \pmod{125}$

Definition is $12^{25} \pmod{125}$. Using the lemma:

$$12 \equiv 2 \pmod{5}$$

$$\begin{aligned} 12^5 &\equiv 2^5 \pmod{5^2} \\ &\equiv 32 \equiv 7 \pmod{25} \end{aligned}$$

$$12^5 \equiv 7 \pmod{25}$$

$$12^{25} \equiv 7^5 \pmod{5^3}$$

$$\begin{aligned} T(12) &\equiv (2 + 5)^5 \pmod{125} \\ &\equiv 2^5 + 5(2^4) * 5 + 10 * 2^3 * 5^2 + \text{multiples of } 125 \\ &\equiv 2^5 + 5^2 * 2^4 \pmod{125} \\ &\equiv 2^5 + 25 * 16 \pmod{125} && \text{note } 16 \equiv 1 \pmod{5} \\ &\equiv 2^5 + 25 * 1 \pmod{125} \\ &\equiv 32 + 25 \pmod{125} \\ &\equiv 57 \pmod{125} \end{aligned}$$

x	$T(x) \pmod{125}$
1	1
2	57
3	$T(-1) * T(2) = -1 * 57 \equiv 68 \pmod{125}$
4	$(-1)^{25} \equiv -1$

Theorem 3.10.

1. If $r > n - 1$ then $x^{p^r} \equiv T(x) \pmod{p^n}$
2. $T(x)^{p-1} \equiv 1 \pmod{p^n}$
3. $T(x)$ depends only on $x \pmod{p}$ and $T(x) \equiv x \pmod{p}$
4. $T: \mathbb{F}_p^\times \mapsto (\mathbb{Z}/p^n)^\times$ is an injective homomorphism

Proof.

By Euler's theorem, $\phi(p^n) = (p-1)p^{n-1}$

$$\begin{aligned} &\implies \underbrace{x^{(p-1)p^{n-1}}}_{T(x)^{p-1}} \equiv 1 \pmod{p^n} \\ &\implies T(x)^{p-1} \equiv 1 \pmod{p^n} \end{aligned}$$

This proves **2**.

$$\therefore T(x)^p \equiv T(x) \pmod{p^n}$$

Doing this several times we get $T(x) \equiv T(x)^p \equiv T(x)^{p^2} \equiv \dots \pmod{p^n}$

This proves **1**.

Suppose:

$$\begin{aligned} x &\equiv y \pmod{p} \\ x^p &\equiv y^p \pmod{p^2} && \text{by the lemma} \\ x^{p^2} &\equiv y^{p^2} \pmod{p^3} && \text{by the lemma} \\ &\vdots \\ T(x) &\equiv T(y) \pmod{p^n} && \text{by Fermat's Little Theorem} \\ x &\equiv x^p \equiv x^{p^2} \equiv \dots \equiv T(x) \pmod{p} \end{aligned}$$

This proves **3**.

$$\begin{aligned} T(xy) &\equiv (xy)^{p^{n-1}} \equiv x^{p^{n-1}} y^{p^{n-1}} \\ &\equiv T(x)T(y) \pmod{p} \end{aligned}$$

So T is a homomorphism, suppose:

$$\begin{aligned} T(x) &\equiv T(y) \pmod{p^n} \\ \therefore T(x) &\equiv T(y) \pmod{p} \\ x &\equiv y \pmod{p} && \text{by 3.} \end{aligned}$$

$$\therefore T: \mathbb{F}_p^\times \mapsto (\mathbb{Z}/p^n)^\times \text{ is injective}$$

□

Corollary 3.11. *Let p be an odd prime, every element in $(\mathbb{Z}/p^n)^\times$ can be written uniquely in the form:*

$$\begin{aligned} &T(x) * \exp(py) \text{ with } x \in \mathbb{F}_p^\times \\ &py \in p\mathbb{Z}/p^n \end{aligned}$$

E.g $22 \in (\mathbb{Z}/125)^\times$

$$22 = T(x) \exp(5y) \pmod{125}$$

$$\equiv 2 \pmod{5}$$

$$\equiv x \pmod{5}$$

$$\implies x \equiv 2 \pmod{5}$$

$$22 = T(2) \exp(5y) \pmod{125}$$

$$22 * T(2^{-1}) \equiv \exp(5y) \pmod{125}$$

from the table $T(3) = 68$

$$\exp(5y) \equiv 22 * 68 \pmod{125}$$

$$\equiv 121 \pmod{125}$$

$$\equiv -4 \pmod{125}$$

$$\therefore 5y \equiv \log(-4) \pmod{125}$$

$$\equiv \log(1 - 5) \pmod{125}$$

$$\equiv -5 - \frac{25}{2} - \frac{125}{3} + \dots$$

$$\equiv -5 - 75$$

$$\equiv 45 \pmod{125}$$

$$\therefore 22 = T(2) \exp(45) \pmod{125}$$

E.g Calculate $22^{37} \pmod{125}$

$$22^{37} \equiv (T(2) \exp(45))^{37}$$

$$\equiv T(2^{37}) * \underbrace{\exp(45 * 37)}_{\equiv 40} \pmod{125}$$

$$\equiv 2 \pmod{5}$$

$$\therefore 22^{37} \equiv \underbrace{T(2)}_{\equiv 57} * \underbrace{\exp(40)}_{\equiv 841}$$

$$\equiv 57 * 91 \pmod{125}$$

$$\equiv 62 \pmod{125}$$

E.g. Calculate $T(23) \pmod{7^3}$

$$\begin{aligned} 23 &\equiv 2 \pmod{7} \\ 23^7 &\equiv 2^7 \equiv 128 \equiv 30 \pmod{7^2} \\ 23^{7^2} &\equiv 30^7 \pmod{7^3} \end{aligned}$$

Using the binomial theorem:

$$\begin{aligned} 23^{7^2} &\equiv (2 + 4 \cdot 7)^7 \pmod{7^3} \\ &\equiv 2^7 + 7 \cdot 2^6 \cdot 4 + \text{multiples of } 7^3 \pmod{7^3} \end{aligned}$$

Since $2^6 \cdot 4 \equiv 4 \pmod{7}$, it follows that $(7^2 \cdot 2^6 \cdot 4) \equiv (49 \cdot 4) \equiv 196 \pmod{7^3}$. This shows that $T(23) \equiv 128 + 196 \equiv 324 \pmod{7^3}$

The following corollary was stated but not proved:

Corollary 3.12. *Let p be an odd prime number. Every element of $(\mathbb{Z}/p^n)^\times$ can be written uniquely in the form $T(x) \cdot \exp(py)$ where $x \in \mathbb{F}_p^\times$ and $py \in p\mathbb{Z}/p^n$. This is an isomorphism of groups:*

$$(\mathbb{Z}/p^n)^\times \cong \mathbb{F}_p^\times * p\mathbb{Z}/p^n$$

Proof. Take any $a \in (\mathbb{Z}/p^n)^\times$ and let $x \equiv a \pmod{p}$. We have $a \equiv x \equiv T(x) \pmod{p}$. Therefore $aT(x)^{-1} \equiv 1 \pmod{p}$. This implies that $\log(\frac{a}{T(x)})$ converges p -adically. Let $py = \log(aT(x)^{-1})$. Then obviously $a = T(x) \exp(py)$.

For uniqueness, suppose $T(x) \cdot \exp(py) \equiv T(x') \cdot \exp(py') \pmod{p^n}$. Since the image of \exp is congruent to 1 \pmod{p} , we have $T(x) \equiv T(x') \pmod{p}$.

This implies $x \equiv x' \pmod{p}$. Therefore $T(x) \equiv T(x') \pmod{p^n}$. From this we get $\exp(py) \equiv \exp(py') \pmod{p^n}$. Taking logs we get $py \equiv py' \pmod{p^n}$

□

This corollary can be used to solve the following types of equations:

E.g. $x^{21} \equiv 71 \pmod{81}$

Note 21 is not coprime to $\phi(81) = 54$, so previous methods cannot be used to solve this equation. Also $(1 + 70)^{\frac{1}{21}}$ does not converge 3-adically. Start with $71 \equiv 2 \pmod{3}$:

$$\implies 71 \equiv T(2) \exp(3y) \pmod{81}$$

$$T(2) \equiv -1 \pmod{81},$$

$$\implies \exp(3y) \equiv -71$$

$$\implies 3y \equiv \log(1 + 9) = 9 - \frac{81}{2} + \dots \equiv 9 \pmod{81}$$

$$\implies 71 = T(2) \exp(9) \pmod{81}$$

Suppose we also decompose $x = T(u) \exp(3v)$. Then

$$x^{21} = T(u^{21}) \exp(3 * 21v) = T(2) \exp(9)$$

Since such a representation is unique, this gives us two simultaneous equations:

$$u^{21} \equiv 2 \pmod{3} \implies u \equiv -1 \equiv 2 \pmod{3}$$

$$\begin{aligned} 63v &\equiv 9 \pmod{81} \implies 7v \equiv 1 \pmod{9} \\ v &\equiv 4 \pmod{9} \\ 3v &\equiv 12 \pmod{27} \end{aligned}$$

$$x \equiv T(2) \exp(12) \pmod{27}$$

$$\begin{aligned} \exp(12) &\equiv 1 + 12 + \frac{12^2}{2} + \frac{12^3}{6} + \frac{12^4}{4!} + \dots \\ &\equiv 1 + 12 + 72 + 288 \\ &\equiv 1 + 12 + 18 + 18 \\ &\equiv 22 \pmod{27} \end{aligned}$$

$$T(2) \equiv -1 \pmod{27}$$

So $x \equiv 5 \pmod{27} \implies x \equiv 5, 32, 59 \pmod{81}$

3.6 Fractional Powers

If p is an odd prime n and $a \equiv 1 \pmod{p}$ and $b \in \mathbb{Z}_{(p)}$ then a^b modulo p^n is:

$$a^b \equiv \exp(b \log(a)) \pmod{p^n}$$

The usual rules hold for powers:

- $(ab)^c \equiv a^c b^c \pmod{p^n}$
- $a^{b+c} \equiv a^b a^c \pmod{p^n}$
- $a^{bc} \equiv (a^b)^c \pmod{p^n}$

E.g $4^{\frac{1}{2}} \pmod{27}$ First find $\log(4) \pmod{27}$

$$\begin{aligned}\log(4) &\equiv (1 + 3) \pmod{27} \\ &\equiv 3 - \frac{9}{2} + \frac{27}{3} \pmod{27} \\ &\equiv 3 + 9 + 9 \pmod{27} \\ &\equiv 6 \pmod{27}\end{aligned}$$

So

$$\begin{aligned}4^b &\equiv \exp(-6b) \pmod{27} \\ 1 - 6b + \frac{36b^2}{2} - \frac{6^3b^3}{6} &\pmod{27} \\ 1 - 6b + 18b^2 - 36b^3 &\pmod{27} \\ 1 - 6b - 9b^2 - 9b^3 &\pmod{27}\end{aligned}$$

$$\begin{aligned}4^{\frac{1}{2}} &\equiv 1 - 3 - 9\left(\frac{1}{4} + \frac{1}{8}\right) \\ &\equiv 1 - 3\frac{27}{8} \\ &\equiv 1 - 3 \\ &\equiv -2 \pmod{27}\end{aligned}$$

3.7 P-adic integers

This section is slightly more highbrow way of looking at the results of the previous lectures. We've defined several congruency classes such as $\exp(px) \pmod{p^n}$, $T(a) \pmod{p^n}$, $\log(1 + px) \pmod{p^n}$. It's a little bit more convenient to be able to write down just $\exp(px)$, $T(a)$, etc ... without needing to write modulo p^n everywhere. The problem is that there is no integer (or even an element of the local ring) which is congruent $T(a) \pmod{p^n}$ for all n . Instead, we work in a bigger ring, the ring \mathbb{Z}_p of p-adic integers. In this ring, the expression $T(a)$, $\exp(px)$ etc all make sense.

Let p be any prime number. By a p-adic integer, we shall mean a p-adically convergent series

$$\sum_{i=1}^{\infty} a_i \qquad a_i \in \mathbb{Z}_{(p)}, V_p(a_i) \longrightarrow \infty$$

Recall that any series represents an element of \mathbb{Z}/p^n for every n . We call two p-adic integers equal if they are congruent modulo p^n for every n . The set of all p-adic integers is denoted \mathbb{Z}_p (without the brackets around the p). Note that we can add, subtract and multiply p-adically convergent series, so in fact \mathbb{Z}_p is a ring.

The advantage of this kind of notation is that we can write (for example) $\log(1 + px)$ to mean a p-adic integer, without having to reduce modulo p^n . This allows us to state many of the recent theorems more simply. If $a \in \mathbb{Z}$ or $\in \mathbb{Z}_{(p)}$, then we can regard a as the series $a = a + 0 + 0 + 0 + \dots$ and so a is a p-adic integer as well. Therefore $\mathbb{Z} \subset \mathbb{Z}_{(p)} \subset \mathbb{Z}$.

However, it turns out that there are many more p-adic integers than there are elements in the local ring $\mathbb{Z}_{(p)}$. For example consider the following 5-adic integer:

$$\begin{aligned} a &= (1 + 5)^{\frac{1}{2}} \\ &= 1 + \frac{1}{2} * 5 + \frac{\frac{1}{2} * \frac{-1}{2}}{2} + 5^2 + \dots \end{aligned}$$

In fact a is a square root of 6. We've shown earlier that $a^2 \equiv 6 \pmod{5^n}$ for all n and therefore $a^2 \equiv 6 \in \mathbb{Z}_5$. However, the local ring $\mathbb{Z}_{(5)}$ has no square roots of 6 since its elements are rational numbers. This shows that a is in \mathbb{Z}_5 but not $\mathbb{Z}_{(5)}$.

Proposition 3.13. *Every p-adic integer can be written uniquely in the form:*

$$\sum_{i=0}^{\infty} a_i p^i$$

with coefficients $a_i \in \{0, 1, \dots, p-1\}$

Proof.

Let x be a p-adic integer, so x is defined modulo p^n for all n . There is a unique choice of a_0 such that $a_0 \equiv x \pmod{p}$.

This means that $x - a_0$ is a multiple of p . There is a unique choice of a_1 such that $a_1 \equiv \frac{x - a_0}{p} \pmod{p}$.

This implies $pa_1 \equiv x - a_0 \pmod{p^2}$, so $x \equiv a_0 + a_1p \pmod{p^2}$. This implies $x - a_0 - a_1p$ is a multiple of p^2 and there is a unique a_2 such that $p^2a_2 \equiv x - a_0 - a_1p \pmod{p^3}$, etc. \square

We've already seen what it means for a series to converge p-adically. We'll now make a corresponding definition for sequences.

Defintion 3.14. *Let a_n be a sequence for elements of $\mathbb{Z}_{(p)}$. We'll say that this sequence converges p-adically if the corresponding series:*

$$a_0 + (a_1 - a_0) + (a_2 - a_1) + \dots$$

If this is the case, then we define the limit of the sequence to be this series, regarded as an element of \mathbb{Z}_p . Note that the partial sums of the series above are exactly the terms of the sequence a_n . In fact, we have already seen many examples of p-adic limits.

Suppose $a_0 \in \mathbb{Z}_{(p)}$ satisfies the conditions of Hensel's lemma for a polynomial $f(x)$, i.e. $f(a_0) \equiv 0 \pmod{p^{2c+1}}$, where $c = V_p(f'(a_0))$. Consider the series:

$$a = a_0 + (a_1 - a_0) + (a_2 - a_1) + (a_3 - a_2) + \dots$$

We'll show that this series converges p-adically. Recall that $a_{n+1} - a_n = \frac{f(a_n)}{f'(a_n)}$

When proving Hensel's lemma, we showed that $f(a_n) \equiv 0 \pmod{p^{2c+2^n}}$, $V_p(f'(a_n)) = c$

Therefore $V_p(a_{n+1} - a_n) \geq 2c + 2^n - c = c + 2^n \rightarrow \infty$

Hence a is a p -adic integer, and is congruent to $a_n \pmod{p^{c+2^n}}$. We can re-interpret Hensel's lemma as saying the following:

Proposition 3.15. *Let a_0 and f satisfy the conditions of Hensel's lemma and let $a \in \mathbb{Z}_p$ be the p -adic integer defined above. Then $f(a) = 0$*

Proof.

We just need to prove that $f(a) \equiv 0 \pmod{\text{any power of } p}$. But we have $f(a) \equiv f(a_n) \equiv 0 \pmod{p^{c+2^n}}$ □

Next consider Teichmüller lifts. For an odd prime p and an element $a \in \mathbb{Z}_{(p)}$ such that $p \nmid a$ let:

$$T(a) = a + (a^p - a) + (a^{p^2} - a^p) + (a^{p^3} - a^{p^2}) + (a^{p^4} - a^{p^3}) + \dots$$

We've shown that $a^{p^n} - a^{p^{n-1}} \equiv 0 \pmod{p^n}$, and therefore the valuation of the n -th term is at least n . This shows that the series converges p -adically, so $T(a) \in \mathbb{Z}_p$. Now the properties of Teichmüller lifts can be restated as follows:

Proposition 3.16. *The p -adic integer $T(a)$ depends only on the congruence class of a modulo p , and the map $T: \mathbb{F}_p^\times \mapsto \mathbb{Z}_p^\times$ is an injective group homomorphism.*

Proof.

Since $T(x) \equiv x \pmod{p}$, it follows that T is injective. For every n , we have $T(xy) \equiv T(x)T(y) \pmod{p^n}$. Therefore $T(xy) = T(x)T(y) \in \mathbb{Z}_p$ □

4 Quadratic rings

An integer d is called square-free if d is not a multiple of a square (apart from 1^2). Let d be a square-free integer with $d \neq 1$. Define a complex number α by:

$$\alpha = \begin{cases} \sqrt{d} & \text{when } d \not\equiv 1 \pmod{4} \\ \frac{1+\sqrt{d}}{2} & \text{when } d \equiv 1 \pmod{4} \end{cases}$$

Consider the set $\{x + y\alpha : x, y \in \mathbb{Z}\}$. This is called a "Quadratic Ring".

Lemma 4.1. *Every quadratic ring is a ring, i.e. closed under $+$, \times .*

Proof. Clearly closed under $+$

$$(x + y\alpha)(r + s\alpha) = xr + (xs + yr)\alpha + ys\alpha^2$$

Sufficient to show that α^2 is in the quadratic ring.

Case 1:

$$\alpha = \sqrt{d} \implies \alpha^2 = d \text{ which is in the quadratic ring}$$

Case 2:

$$\alpha = \frac{1 + \sqrt{d}}{2} \qquad d \equiv 1 \pmod{4}$$

$$\left(\alpha - \frac{1}{2}\right)^2 = \frac{d}{4}$$

$$\alpha^2 - \alpha + \frac{1}{4} = \frac{d}{4}$$

$$\alpha \equiv \alpha + \frac{d-1}{4} \qquad d-1 \equiv 0 \pmod{4} \implies \frac{d-1}{4} \in \mathbb{Z}$$

$\therefore \alpha^2$ is in the quadratic ring. □

We call $\mathbb{Z}[\alpha] = \{x + y\alpha : x, y \in \mathbb{Z}\}$:

- A real quadratic ring if $d > 0$
- A complex quadratic ring if $d < 0$

E.g $d = -1$

$$-1 \equiv 1 \pmod{4}$$

$$\therefore \alpha = \sqrt{-1} = i$$

$\mathbb{Z}[i] = \{x + iy : x, y \in \mathbb{Z}\}$ is the ring of Gaussian integers.

E.g $d = -3$

$$-3 \equiv 1 \pmod{4} \text{ so } \alpha = \frac{1+\sqrt{3}}{2}$$

This is the ring of Eisenstein integers. It is the same as $\mathbb{Z}[\zeta_3] = \mathbb{Z}[e^{\frac{2\pi i}{3}}]$.

Defintion 4.2. Let $\mathbb{Z}[\alpha]$ be a quadratic ring. The elements all have the form $A = x + y\sqrt{d}$ where x, y are rational. The conjugate of such an element $\bar{A} = x - y\sqrt{d}$

4.0.1 Properties of conjugates

1 $\bar{\alpha} =$

$$1. \bar{\alpha} = \begin{cases} -\alpha & d \not\equiv 1 \pmod{4} \\ 1 - \alpha & d \equiv 1 \pmod{4} \end{cases}$$

$$2. \overline{A+B} = \bar{A} + \bar{B} \\ \overline{AB} = \bar{A} \bar{B}$$

$$3. \bar{A} \in \mathbb{Z}[\alpha] \text{ if } A \in \mathbb{Z}[\alpha]$$

$$4. \bar{\bar{A}} = A$$

Proof.

$$1. \text{ If } d \not\equiv 1 \pmod{4} \text{ then } \alpha = \sqrt{d} \\ \implies \bar{\alpha} = -\sqrt{d} = -\alpha$$

$$\text{If } d \equiv 1 \pmod{4} \text{ then } \alpha = \frac{1+\sqrt{d}}{2} \\ \implies \bar{\alpha} = \frac{1-\sqrt{d}}{2} = 1 - \alpha$$

2. Suppose:

$$A = x + y\sqrt{d}$$

$$B = r + s\sqrt{d}$$

Clearly $\overline{A+B} = \bar{A} + \bar{B}$

$$\begin{aligned} \overline{A+B} &= \overline{(x + y\sqrt{d})(r + s\sqrt{d})} \\ &= \overline{(xr + dys) + (xs + yr)\sqrt{d}} \\ &= \overline{(xr + dys) - (xs + yr)\sqrt{d}} \end{aligned}$$

$$\begin{aligned} \bar{A} \cdot \bar{B} &= (x - y\sqrt{d})(r - s\sqrt{d}) \\ &= (xr + dys) + (-xs - yr)\sqrt{d} \end{aligned}$$

3. Let $A = x + y\alpha$ $x, y \in \mathbb{Z}$

by **2.** $\bar{A} = x + y\alpha$

by **1.** $\bar{\alpha} \in \mathbb{Z}[\alpha]$

$\therefore \bar{A} \in \mathbb{Z}[\alpha]$

4. Trivial □

Defintion 4.3. For an element $A \in \mathbb{Z}$ we define $N(A) = A\bar{A}$ - The norm of A

Remark: If $\mathbb{Z}[\alpha]$ is a complex quadratic ring then \bar{A} is the complex conjugate of A . This means $N(A) = |A|^2$

E.g. $d = -1 \implies \mathbb{Z}[\alpha] = \mathbb{Z}[i]$

The elements have the form $x + iy$, $x, y \in \mathbb{Z}$

$$\begin{aligned} N(x + iy) &= (x + iy)(x - iy) \\ &= x^2 + y^2 \end{aligned}$$

E.g. $d = -3 \implies -3 \equiv 1 \pmod{4}$, so $\alpha = \frac{1+\sqrt{-3}}{2}$

$$\begin{aligned} N(x + y\alpha) &= (x + y\alpha)(x + y\bar{\alpha}) & \bar{\alpha} &= 1 - \alpha \\ &= x^2 + xy \underbrace{(\alpha + 1 - \alpha)}_{=1} + y^2 \underbrace{(\alpha(1 - \alpha))}_{=1} \end{aligned}$$

Note that:

$$\begin{aligned} \alpha &= \frac{1 + \sqrt{d}}{2} \\ \implies \left(\alpha - \frac{1}{2}\right)^2 &= \frac{d}{4} \\ \implies \alpha^2 - \alpha + \frac{1}{4} &= \frac{d}{4} \\ \implies \alpha(1 - \alpha) &= \frac{1 - d}{4} \end{aligned}$$

In this case:

$$N(x + y\alpha) = x^2 + xy + y^2$$

4.0.2 Formula for norms

The general formula for norms is given by:

$$N(x + y\alpha) = \begin{cases} x^2 - dy^2 & \text{if } d \not\equiv 1 \pmod{4} \\ x^2 + xy + \frac{1-d}{4}y^2 & \text{if } d \equiv 1 \pmod{4} \end{cases}$$

Case 1: $d \neq 1$ (4)

$$\implies \bar{\alpha} \equiv -\alpha$$

$$N(x + y\alpha) = (x + y\alpha)(x - y\alpha) = x^2 - dy^2 \quad \alpha^2 = d$$

Case 2: $\bar{\alpha} = 1 - \alpha$

$$\implies \alpha = \frac{1+\sqrt{d}}{2}$$

$$\begin{aligned} N(x + y\alpha) &= (x + y\alpha)(x + y(1 - \alpha)) \\ &= x^2 + xy + y^2(\underbrace{\alpha - \alpha^2}_{\frac{1-d}{4}}) \end{aligned}$$

4.0.3 Properties of norms

1. $N(A) \in \mathbb{Z}$
2. $N(AB) = N(A)N(B)$
3. If $N(A) = 0$ then $A = 0$

Proof.

1. Follows from formulas for norms
2. $N(AB) = AB\overline{AB} = AB\bar{A}\bar{B}$
3. If $N(A) = 0$ then $A\bar{A} = 0$
 \therefore either $A = 0$ or $\bar{A} = 0$
 $\therefore \bar{A} = 0$ then $A = \bar{A} = \bar{0} = 0$

□

Recall - A unit in a ring R is an element with an inverse in R .

E.g. $1 + \sqrt{2}$ is a unit in $\mathbb{Z}[\sqrt{2}] \implies (1 + \sqrt{2})(\sqrt{2} - 1) = 1$

Corollary 4.4. *An element $A \in \mathbb{Z}[\alpha]$ is a unit if and only if $N(A) = \pm 1$*

Proof. If $N(A) = \pm 1$ then $A\bar{A} = \pm 1$

$$\implies A^{-1} = \pm A \in \mathbb{Z}[\alpha]. \text{ So } A \text{ is a unit.}$$

□

If A is a unit with inverse B , $AB = 1 \implies N(A)N(B) = N(AB) = N(1) = 1$

Using this proposition, it's easy to find all the units in any complex quadratic ring.

E.g The units in $\mathbb{Z}[i]$ are $\pm 1, \pm i$

Proof. Since $N(x + iy) = x^2 + y^2$, the units correspond to the solutions to $x^2 + y^2 = 1$.

These solutions are $x = \pm 1, y = 0$ and $x = 0, y = \pm 1$

□