

Corollary

50

Let G be a graph with p -vertices. If $\deg v \geq \frac{p-1}{2}$ for every vertex v of G , then G contains a Hamiltonian path.

Proof: If $p=1$, then $G \cong K_1$ and G contains a (trivial) Hamiltonian path.

Suppose $p \geq 2$ and define $H = G + K_1$.

Let v denote the vertex of H that is not in G .

Since H has a vertex $p+1$, it follows that $\deg v \geq p$. Moreover, for every vertex u of G

$$\deg_H u = \deg_G u + 1 \geq \frac{p-1}{2} + 1 = \frac{p+1}{2} = \frac{V(H)}{2}.$$

By Dirac's theorem, H contains a Hamiltonian ~~cycle~~ cycle. By removing the vertex v from C , we obtain a Hamiltonian path in G .

ORE'S THEOREM

If G is a simple graph with $n \geq 3$ number of vertices and if $\deg(u) + \deg(v) \geq n$ for every pair of non-adjacent vertices $u \neq v$, then G is Hamiltonian.

Proof: We use the method of contradiction to prove this theorem. Suppose there exist a non-Hamilton graph with n vertices satisfying

$$\deg(u) + \deg(v) \geq n$$

for ~~for~~ non-adjacent vertices $u \neq v$. Among all such non-Hamiltonian graphs, let G be a non-Hamiltonian graph with maximum number of edges.

Since G is maximal non-Hamiltonian graph, there exists two non-adjacent vertices u and v in G such that addition of any edge joining u and v will result a Hamiltonian graph.

Thus there ~~must~~ exist a Hamiltonian path in G , namely $u = u_1, u_2, \dots, u_n = v$ with u and v are end vertices.

$$\text{Let } A = \{i \mid u_i \text{ is adjacent to } u \text{ in } G\}$$

$$B = \{i \mid v_i \text{ is adjacent to } v \text{ in } G\}$$

$$\text{Now } |A| = \deg(u), |B| = \deg(v).$$

$$\text{Claim: (i) } A \cap B = \emptyset$$

$$(ii) |A \cup B| \leq n-1$$

If possible $A \cap B \neq \emptyset$, then say $i \in A \cap B$

$\Rightarrow (u, u_i)$ and (u_i, v) could be in G ,

and then G could form a Hamiltonian circuit, which is a contradiction. Thus

$$A \cap B = \emptyset$$

$$\text{Now } A \cup B = \{1, 2, \dots, n\}. \text{ Since } u_1 = u \text{ is}$$

~~not~~ neither adjacent to u nor adjacent to v ,
 $1 \notin A \cup B$.

$$\text{So } |A \cup B| \leq n-1$$

We know that

$$\begin{aligned} |A \cup B| &= |A| + |B| - |A \cap B| \\ &= |A| + |B| \quad (\because |A \cap B| = 0) \end{aligned}$$

$$\Rightarrow |A| + |B| \leq n-1$$

$$\Rightarrow \deg(u) + \deg(v) \leq n-1$$

which is a contradiction to the given fact that $\deg(u) + \deg(v) \geq n$.

Therefore, G is a Hamilton graph. (proved)

Ramsey's theorem for graphs

Proposition: Among any six people, there are three any two of whom are friends, or there are three such that no two of them are friends.

Proof: Let $G = (V, E)$ be a graph and $|V| = 6$. For a vertex $v \in V$. We consider two cases.

Case-I: If the degree of v is at least 3, then consider three neighbors of v , call them x, y, z . If any two among $\{x, y, z\}$ are friends, we are done because they form a triangle together with v . If not, no two of $\{x, y, z\}$ are friend and we are done as well.

Case-II: If degree of v is at most 2, then there are at least three other vertices

which are not neighbors of v , call them u, y, z . In this case, the argument is complementary to the previous one. Either $\{u, y, z\}$ are mutual friends, in which case we are done. Or there are two among $\{u, y, z\}$ who are not friends, for example u and y , and then no two of $\{v, u, y\}$ are friends.

More generally, we color the edge K_n (a complete graph on n vertices) with a certain number of colors and we ask whether there is a complete subgraph (a clique) of certain size such that all its edges have same color.

Definition

A clique of size k is a set of k vertices such that all pairs among them are edges.

An independent set of size s is a set s vertices such that there is no edge between them.

Ramsey's theorem states that for any large enough graph, there is an independent set of size s or a clique of size k . The smallest number of vertices required to achieve this is called a Ramsey number.

Definition

The Ramsey number $R(s, t)$ is the minimum number n such that any graph on n vertices contains either an independent set of size s or a clique of size t .

Properties

$$(i) R(s, t) = R(t, s)$$

$$(ii) R(1, 1) = 1, R(2, 2) = 2, R(3, 3) = 6, R(4, 4) = 18.$$

$$(iii) R(s, 1) = R(1, t) = 1$$

$$(iv) R(2, t) = t; R(s, 2) = s$$

$$(v) R(s, t) \leq R(s-1, t) + R(s, t-1) \quad \forall s, t \geq 2 \text{ and } s, t \text{ are integers.}$$

Upper bound for $R(s, t)$

$$R(s, t) \leq \binom{s+t-2}{s-1} = {}^{s+t-2}C_{s-1} \quad (\text{where } C \text{ is combination})$$

Lower bound for $R(s, t)$

$$R(s, t) \geq 2^{\frac{k}{2}}, \quad k = \min\{s, t\}$$

That means

$$2^{\frac{k}{2}} \leq R(s, t) \leq {}^{s+t-2}C_{s-1}$$

Problem

Find the upper bound and Lower bound of Ramsey number $R(4, 4)$.

Solⁿ: upper bound

$$R(4, 4) \leq {}^{4+4-2}C_{4-1} = {}^8C_3 = \frac{8!}{(8-3)!3!} = \frac{8!}{5!3!} = \frac{6 \times 7 \times 8}{2 \times 3} = 56$$

Lower bound

$$k = \min\{4, 4\} = 4$$

$$R(4, 4) \geq 2^{\frac{4}{2}} = 2^2 = 4.$$

Ramsey's Theorem

For any $s, t \geq 1$, there is $R(s, t) < \infty$ such that any graph on $R(s, t)$ vertices contains either an independent set of size s or a clique of size t . In particular,

$$R(s, t) \leq \binom{s+t-2}{s-1}.$$

Proof: We show that $R(s, t) \leq R(s-1, t) + R(s, t-1)$. To see this, let $n = R(s-1, t) + R(s, t-1)$ and consider any graph G on n vertices. For a vertex $v \in V$, we consider two cases:

Case-I

There are at least $R(s, t-1)$ edges incident with v . Then we apply induction on the neighbors of v , which implies that either they contain an independent set of size s , or a clique of size $t-1$. In the second case, we extend the clique by adding v , and hence G contains either an independent set of size s or a clique of size t .

Case-II

There are at least $R(s-1, t)$ non-neighbors of v . Then we apply induction to the non-neighbors of v and we get either an independent set of size

(56)

$s-1$, or a clique of size ℓ . Again, the independent set can be extended by adding v and hence we are done.

In particular, $R(s, \ell) \leq R(s-1, \ell) + R(s, \ell-1)$, it follows ~~that~~ by induction that these Ramsey numbers are finite. Moreover, we get an explicit bound.

First, $R(s, \ell) \leq \binom{s+\ell-2}{s-1}$ holds for the base case, where $s=1$ or $\ell=1$ since every graph contains a clique or an independent set of size 1. The inductive step is as follows:

$$R(s, \ell) \leq R(s-1, \ell) + R(s, \ell-1)$$

$$\leq \binom{s+\ell-3}{s-2} + \binom{s+\ell-3}{s-1}$$

$$= \binom{s+\ell-2}{s-1}, \text{ where } \binom{\cdot}{\cdot} \text{ is combination.}$$

(proved)