


By choosing $m=2$, \exists a complete matching from the set V_1 to V_2 .

Problem

Prove that the bipartite graph shown in the following graph does not have a complete matching.



Solⁿ: We observe that the three vertices v_1, v_2, v_3 in V_1 are together joined to two vertices a_1, a_2 in V_2 . Thus, there is a subset of 3 vertices in V_1 which is collectively adjacent to 2 (< 3) vertices in V_2 .

Hence, by Hall's theorem, there does not exist a complete matching from V_1 to V_2 .

Matching in General Graphs

Definition

A factor of a graph G is a spanning subgraph of G . A k -factor is a spanning k -regular subgraph. An odd component of a graph is a component of odd order; the number of odd components of H is $o(H)$.

Remark

→ A 1-factor and a perfect matching are almost the same thing.

(13)

→ The precise definition is that "1-factor" is a spanning 1-regular subgraph of G , while "perfect matching" is the set of edges in such a subgraph.

→ A 3-regular graph that has a perfect matching ~~decomposition~~ decomposes into a 1-factor and 2-factor.

TUTTE'S 1-FACTOR THEOREM

Tutte found a necessary and sufficient condition for which graph have 1-factor and we consider a set $S \subseteq V(G)$, then every odd component of $G-S$ has a vertex matched to something outside it, which can only belong to S . Since these vertices of S must be distinct, $o(G-S) \leq |S|$.


The condition "For all $S \subseteq V(G)$, $o(G-S) \leq |S|$ " is Tutte's Condition.

Theorem

A graph G has a 1-factor if and only if $o(G-S) \leq |S|$ for every $S \subseteq V(G)$.

Definition

The join of simple graphs G and H , written $G \vee H$, is the graph obtained from the disjoint union $G+H$ by adding adding the edges $\{xy : x \in V(G), y \in V(H)\}$.



$P_4 \vee K_3$

Berge-Tutte Formula

The largest number of vertices saturated by a matching in G is $\max_{S \subseteq V(G)} \{n(G) - d(S)\}$, where $d(S) = o(G - S) - |S|$.

Proof: Given $S \subseteq V(G)$, at most $|S|$ edges can match vertices of S to vertices in odd components of $G - S$, so every matching has at least $o(G - S) - |S|$ unsaturated vertices. We want to achieve this bound.

Let $d = \max \{o(G - S) - |S| : S \subseteq V(G)\}$. The case $v = \emptyset$ gives $d \geq 0$. Let $G' = G \vee K_d$. Since $d(S)$ has the same parity as $n(G)$ for each S , we know that $n(G')$ is even. If G' satisfies Tutte's Condition, then we obtain a matching of the desired size in G from a perfect matching in G' , because deleting the d added vertices eliminates edges that saturate at most d vertices of G .

The condition $o(G' - S') \leq |S'|$ holds for $S' = \emptyset$ because $n(G')$ is even. If S' is nonempty but not contain all of K_d , then $G' - S'$ has only one component, and $1 \leq |S'|$. Finally, when $K_d \subseteq S'$, we let $S = S' - V(K_d)$. We have $G' - S' = G - S$, so $o(G' - S') = o(G - S) \leq |S| + d = |S'|$.

Petersen 1-factor Theorem

Every 3-regular graph with no cut-edge has a 1-factor.

Proof: Let G be a 3-regular graph with no cut-edge. We prove that G satisfies Tutte's Condⁿ. Given $S \subseteq V(G)$, we count the edges between S and the odd components of $G-S$. Since G is 3-regular, each vertex of S is incident to at most three such edges. If each odd component H of $G-S$ is incident to at least three such edges, then $3o(G-S) \leq 3|S|$ and hence $o(G-S) \leq |S|$.

Let m be the number of edges from S to H . The sum of the vertex degrees in H is $3n(H)-m$. Since H is a graph, the sum of ~~the~~^{its} vertex degrees must be even. Since $n(H)$ is odd, we conclude that m must also be odd. Since G has no cut edge, m can not equal 1. We conclude that there are at least three edges from S to H . Hence every 3-regular graph with no cut-edge has a 1-factor. (proved)

Petersen 2-factor Theorem

Every regular graph of even degree has a 2-factor.

Proof: Let G be a $2k$ -regular graph with vertices v_1, v_2, \dots, v_n . Every component of G is

(16)

Eulerian, with some Eulerian circuit C . For each component, define a bipartite graph H with vertices u_1, u_2, \dots, u_n and w_1, w_2, \dots, w_n by putting $u_i \leftrightarrow w_j$ if v_j immediately follows v_i somewhere on C . Because C enters and exits each vertex n times, H is n -regular.

Being a regular bipartite graph, H has a 1-factor M . The edge incident to w_i in H corresponding to an edge entering v_i in C . The edge incident to u_i in H corresponding to an edge exiting v_i . Thus ^{that} 1-factor in H transforms into a 2-regular spanning subgraph of this component of G . Doing this for each component of G yields a 2-factor of G . Hence every ~~component~~ regular graph of ~~an~~ even degree has a 2-factor. (proved)

Example (Construction of a 2-factor)

Consider the Eulerian circuit on $G = K_5$ that successively visits 1231425435. The corresponding bipartite graph H is on the right. For the 1-factor whose u, w -pairs are 12, 43, 25, 31, 54, the resulting 2-factor is the cycle $(1, 2, 5, 4, 3)$. The remaining edges form another 1-factor, which corresponds to the 2-factor ~~(1, 2, 3, 4, 5)~~ $(1, 4, 2, 3, 5)$ that remains in G .

