ANNEX I

SUMMARY OF PRODUCT CHARACTERISTICS

1. NAME OF THE VETERINARY MEDICINAL PRODUCT

Locatim Oral solution

2. QUALITATIVE AND QUANTITATIVE COMPOSITION

Active substances:

Bovine concentrated lactoserum containing specific immunoglobulins G against *E. coli* F5 (K99) adhesin ≥ 2.8* $\log_{10}$/ml.
* ELISA method

Excipients:

Qualitative composition of excipients and other constituents	**Quantitative composition if that information is essential for proper administration of the veterinary medicinal product**
Methyl parahydroxybenzoate	≤ 0.8 mg/ml.

Yellow liquid, slightly opalescent.

3. CLINICAL INFORMATION

3.1 Target species

Cattle (newborn calves) less than 12 hours of age

3.2 Indications for use for each target species

Reduction of mortality caused by enterotoxicosis associated with *E. coli* F5 (K99) adhesin during the first days of life as a supplement to colostrum from the dam.

3.3 Contraindications

None.

3.4 Special warnings

The veterinary medicinal product is produced from colostrum collected from cows kept under field conditions.
Consequently, in addition to antibodies to *E. coli* F5 (K99) it also contains antibodies to other organisms, as a result of vaccination and/or exposure of the donor cows to organisms in their environment.
This should be borne in mind when planning vaccination programmes for calves, which receive the veterinary medicinal product.

3.5 Special precautions for use

Special precautions for safe use in the target species:

This veterinary medicinal product may contain antibodies against Bovine Viral Diarrhoea (BVD) virus.

Special precautions to be taken by the person administering the veterinary medicinal product to animals:

Not applicable.

Special precautions for the protection of the environment:

Not applicable.

3.6 Adverse events

None known.

Reporting adverse events is important. It allows continuous safety monitoring of a veterinary medicinal product. Reports should be sent, preferably via a veterinarian, to either the marketing authorisation holder or its local representative or the national competent authority via the national reporting system. See the package leaflet for respective contact details.

3.7 Use during pregnancy, lactation or lay

Pregnancy and lactation:

The veterinary medicinal product is not intended for use during pregnancy and lactation.

3.8 Interaction with other medicinal products and other forms of interaction

No information is available on the safety and efficacy of this immunological veterinary medicinal product when used with any other veterinary medicinal product. A decision to use this immunological veterinary medicinal product before or after any other veterinary medicinal product therefore needs to be made on a case-by-case basis.

3.9 Administration routes and dosage

Oral administration of 60 ml as soon as possible, preferably given within the first 4 hours, but not later than 12 hours after birth.
The veterinary medicinal product should be administered neat or diluted in milk or in milk replacer within the first 12 hours of the cattle (newborn calve)'s life, preferably, as soon as it is receptive. If the cattle (newborn calve) is reluctant to take the product, it may be administered via an ordinary syringe placed in the mouth.
The cattle (newborn calve) must be given other normal colostrum in addition to the veterinary medicinal product.
In the absence of information specifically demonstrating the safety of more than one repeated dose, it is recommended that cattle (newborn calve) should only be dosed once.

3.10 Symptoms of overdose (and where applicable, emergency procedures and antidotes)

Transient effects of temperature increase and respiration rate increase have been seen when the veterinary medicinal product is administered in a double dose.

3.11 Special restrictions for use and special conditions for use, including restrictions on the use of antimicrobial and antiparasitic veterinary medicinal products in order to limit the risk of development of resistance

Not applicable.

3.12 Withdrawal periods

Zero days.

4. IMMUNOLOGICAL INFORMATION

4.1 ATCvet code: QI02AT01

The product supplements the protective properties of normal colostrum against E. coli F5 (K99) adhesin.

5. PHARMACEUTICAL PARTICULARS

5.1 Major incompatibilities

In the absence of compatibility studies, this veterinary medicinal product must not be mixed with other veterinary medicinal products.

5.2 Shelf life

Shelf life of the veterinary medicinal product as packaged for sale: 30 months.

5.3 Special precautions for storage

Store in a refrigerator (2 °C – 8 °C).
Do not freeze.
Keep the glass bottle in the outer carton.

5.4 Nature and composition of immediate packaging

Cardboard box with 1, 6, 12, 24 or 48 60 ml type III glass bottles closed with a polypropylene stopper with a polyethylene seal and a detachable lock-ring.

Not all pack sizes may be marketed.

5.5 Special precautions for the disposal of unused veterinary medicinal products or waste materials derived from the use of such products

Medicines should not be disposed of via wastewater or household waste.

Use take-back schemes for the disposal of any unused veterinary medicinal product or waste materials derived thereof in accordance with local requirements and with any national collection systems applicable to the veterinary medicinal product concerned.

6. NAME OF THE MARKETING AUTHORISATION HOLDER

Melchior santé animale

7. MARKETING AUTHORISATION NUMBER(S)

EU/2/99/011/001-005

8. DATE OF FIRST AUTHORISATION

Date of first authorisation: 29/03/1999.

9. DATE OF THE LAST REVISION OF THE SUMMARY OF THE PRODUCT CHARACTERISTICS

{JJ/MM/AAAA}

10. CLASSIFICATION OF VETERINARY MEDICINAL PRODUCTS

Veterinary medicinal product subject to prescription.

Detailed information on this veterinary medicinal product is available in the Union Product Database (https://medicines.health.europa.eu/veterinary).

ANNEX II

OTHER CONDITIONS AND REQUIREMENTS OF THE MARKETING AUTHORISATION

None.

ANNEX III

LABELLING AND PACKAGE LEAFLET

A. LABELLING

PARTICULARS TO APPEAR ON THE OUTER PACKAGE

BOX 1 x 60 ml
BOX 6 x 60 ml
BOX 12 x 60 ml
BOX 24 x 60 ml
BOX 48 x 60 ml

1. NAME OF THE VETERINARY MEDICINAL PRODUCT

Locatim Oral solution

2. STATEMENT OF ACTIVE SUBSTANCES

Bovine concentrated lactoserum containing specific immunoglobulins G against E. coli F5 (K99) adhesin ≥2.8 $\log_{10}$/ml.

3. PACKAGE SIZE

60 ml.
6 x 60 ml
12 x 60 ml
24 x 60 ml
48 x 60 ml

4. TARGET SPECIES

Cattle (newborn calves) less than 12 hours of age.

5. INDICATIONS

6. ROUTES OF ADMINISTRATION

Oral solution

7. WITHDRAWAL PERIODS

Withdrawal periods: Zero days.

8. EXPIRY DATE

Exp. {mm/yyyy}

9. SPECIAL STORAGE PRECAUTIONS

Store in a refrigerator.

Keep the glass bottle in the outer carton.
Do not freeze.

10. THE WORDS "READ THE PACKAGE LEAFLET BEFORE USE"

Read the package leaflet before use.

11. THE WORDS "FOR ANIMAL TREATMENT ONLY"

For animal treatment only.

12. THE WORDS "KEEP OUT OF THE SIGHT AND REACH OF CHILDREN"

Keep out of the sight and reach of children.

13. NAME OF THE MARKETING AUTHORISATION HOLDER

Melchior santé animale

14. MARKETING AUTHORISATION NUMBERS

EU/2/99/011/001
EU/2/99/011/002
EU/2/99/011/003
EU/2/99/011/004
EU/2/99/011/005

15. BATCH NUMBER

Lot {number}

PARTICULARS TO APPEAR ON THE IMMEDIATE PACKAGE

Bottle 60 mL

1. NAME OF THE VETERINARY MEDICINAL PRODUCT

Locatim Oral solution

2. STATEMENT OF ACTIVE SUBSTANCES

Bovine concentrated lactoserum containing specific immunoglobulins G against E. coli F5 (K99) adhesin ≥2.8 $\log_{10}$/ml.

3. TARGET SPECIES

Cattle (newborn calves) less than 12 hours of age.

4. ROUTES OF ADMINISTRATION

Oral use
Read the package leaflet before use.

5. WITHDRAWAL PERIODS

Withdrawal periods: zero days.

6. EXPIRY DATE

Exp. {mm/yyyy}

7. SPECIAL STORAGE PRECAUTIONS

Store in a refrigerator.
Keep the glass bottle in the outer carton.
Do not freeze.

8. NAME OF THE MARKETING AUTHORISATION HOLDER

Melchior santé animale

9. BATCH NUMBER

Lot {number}

B. PACKAGE LEAFLET

PACKAGE LEAFLET

1. Name of the veterinary medicinal product

Locatim Oral solution

2. Composition

Bovine concentrated lactoserum containing specific immunoglobulins G against E. coli F5 (K99) adhesin ≥2.8 $\log_{10}$/ml.
* ELISA method

3. Target species

Cattle (newborn calves) less than 12 hours of age.

4. Indications for use

Reduction of mortality caused by enterotoxicosis associated with *E. coli* F5 (K99) adhesin during the first days of life as a supplement to colostrum from the dam.

5. Contraindications

None.

6. Special warnings

Special warnings:
The product is produced from colostrum collected from cows kept under field conditions. Consequently, in addition to antibodies to *E. coli* F5 (K99) it also contains antibodies to other organisms, as a result of vaccination and/or exposure of the donor cows to organisms in their environment. This should be borne in mind when planning vaccination programmes for calve, which receive the veterinary medicinal product.

Special precautions for safe use in the target species:
This product may contain antibodies against Bovine Viral Diarrhoea (BVD) virus.

Pregnancy and lactation:
The product is not intended for use during pregnancy and lactation.

Interaction with other medicinal products and other forms of interaction:
No information is available on the safety and efficacy of this product when used with any other veterinary medicinal product. A decision to use this product before or after any other veterinary medicinal product therefore needs to be made on a case-by-case basis.

Major incompatibilities:
In the absence of compatibility studies this veterinary medicinal product must not be mixed with other veterinary medicinal products.

Overdose:

Transient effects of temperature increase and respiration rate increase have been seen when the product is administered in a double dose.

7. Adverse events

None known.

Reporting adverse events is important. It allows continuous safety monitoring of a product. If you notice any side effects, even those not already listed in this package leaflet, or you think that the medicine has not worked, please contact, in the first instance, your veterinarian. You can also report any adverse events to the marketing authorisation holder using the contact details at the end of this leaflet, or via your national reporting system: {national system details}

8. Dosage for each species, routes and method of administration

Oral administration of 60 ml as soon as possible, preferably given within the first 4 hours, but not later than 12 hours after birth.

9. Advice on correct administration

The product should be administered neat or diluted in milk or in milk replacer within the first 12 hours of the cattle (newborn calve)'s life, preferably, as soon as it is receptive. If the cattle (newborn calve) is reluctant to take the product, it may be administered via an ordinary syringe placed in the mouth.

The cattle (newborn calve) must be given other normal colostrum in addition to the product.

In the absence of information specifically demonstrating the safety of more than one repeated dose, it is recommended that calves should only be dosed once.

10. Withdrawal periods

Zero days.

11. Special storage precautions

Keep out of the sight and reach of children.

Store in a refrigerator (2 °C – 8 °C).
Do not freeze.
Keep the glass bottle in the outer carton.

Do not use this veterinary medicinal product after the expiry date which is stated on the label after Exp. The expiry date refers to the last day of that month.

12. Special precautions for disposal

Medicines should not be disposed of via wastewater or household waste.

Use take-back schemes for the disposal of any unused veterinary medicinal product or waste materials derived thereof in accordance with local requirements and with any applicable national collection systems. These measures should help to protect the environment.

Ask your veterinary surgeon or pharmacist how to dispose of medicines no longer required.

13. Classification of veterinary medicinal products

Veterinary medicinal product subject to prescription.

14. Marketing authorisation numbers and pack sizes

EU/2/99/011/001-005

Cardboard box with 1, 6, 12, 24 or 48 60 ml type III glass bottles closed with a polypropylene stopper with a polyethylene seal and a detachable lock-ring.

Not all pack sizes may be marketed.

15. Date on which the package leaflet was last revised

21/05/2026

Detailed information on this veterinary medicinal product is available in the Union Product Database (https://medicines.health.europa.eu/veterinary).

16. Contact details

Marketing authorisation holder and contact details to report suspected adverse events:
Melchior santé animale
5 Rue Victor Hugo, 69002 Lyon
France
Tel: +33 6 18 15 03 91

Manufacturer responsible for batch release:
BIOKEMA EUROPE
Centre des Affaires des Lilas 77
Avenue des Lilas 64000 Pau
France

For any information about this veterinary medicinal product, please contact the local representative of the marketing authorisation holder:

België/Belgique/Belgien
Dopharma Holding B.V.
Zalmweg 24
4941 VX Raamsdonksveer - NL
Tél/Tel: +32 475 367776

Lietuva
Dopharma Polska
ul. Wojskowa 6/B02
60 792 Poznań, PL
Tel: +48 516 052 508

Република България
Dopharma Holding B.V.
Zalmweg 24
4941 VX Raamsdonksveer - NL
Тел: +31 162 58 20 00

Česká republika
Cymedica, spol. s r.o.
Pod Nádražím 308/24
268 01 Hořovice- CZ
Tel: + 420 311 706 211

Deutschland
Dopharma Deutschland GmbH
Hansestr. 53
DE-48165 Münster
Tel: +49 (0)2501 594 349 20

Eesti
Dopharma Polska
ul. Wojskowa 6/B02
60 792 Poznań - PL
Tel: +48 516 052 508

Ελλάδα
GEROLYMATOS INTERNATIONAL S.A.
Ασκληπιού 8, 145 68, Κρυονέρι
Αττική , EL
Τηλ: + 30 210 350 0854

España
Dopharma Iberia S.L.
Avenida de la Llana 123
08191, Rubí – Barcelona – ES
Tel: +34 637 37 04 92

France
Melchior santé animale
5 rue Victor Hugo
69002 Lyon – FR
Tél: + 33 6 18 15 03 91

Hrvatska
Arnika Veterina d.o.o.
Vidikovac 20
10000 Zagreb - HR
Tel: + 385 91 364 3731

Luxembourg/Luxemburg
Dopharma Holding B.V.
Zalmweg 24
4941 VX Raamsdonksveer - NL
Tél/Tel: +32 475 367776

Magyarország
Dopharma Vet S.R.L.
Str. Aeroport nr. 44
Localitatea Ghiroda
RO-307200 Județul Timiș
Tel.: +40 728 138 903

Nederland
Dopharma Holding B.V.
Zalmweg 24
4941 VX Raamsdonksveer NL
Tel: +31 162 58 20 00

Austria
Dopharma Deutschland GmbH
Hansestr. 53
DE-48165 Münster
Tel: +49 (0)2501 594 349 20

Polska
Dopharma Polska
ul. Wojskowa 6/B02
60 792 Poznań - PL
Tel: +48 516 052 508

Italia
Dopharma Italia S.r.l.
Via delle Porte Nuove, 20
50144 Firenze – IT
Tel: + 39 (346) 14 26 164

Portugal
Vetlima-Sociedade Distribuidora De Produtos Agro-Pecuários, S.A., Centro Empresarial de Rainha
Lote 27, 2050-501 Vila Nova da Rainha PT
Tel: + 351 964404163

România
Dopharma Vet S.R.L.
Str. Aeroport nr. 44
Localitatea Ghiroda
RO-307200 Județul Timiș
Tel: +40 728 138 903

Latvija
Dopharma Polska
ul. Wojskowa 6/B02
60 792 Poznań
PL
Tel: +48 516 052 508

United Kingdom (Northern Ireland)
FORTE Healthcare Limited,
Block 3, Unit 9,
CityNorth Business Campus,
Stamullen, Co. Meath. K32 D990
Republic of Ireland – UK
Tel: +353 1 841 7666

Κύπρος
GEROLYMATOS INTERNATIONAL S.A.
Ασκληπιού 8, 145 68, Κρυονέρι
Αττική , EL
Τηλ: + 30 210 350 0854

Slovenská republika
Cymedica, spol. s r.o.
Pod Nádražím 308/24
268 01 Hořovice- CZ
Tel: + 420 311 706 211

17. Other information

The product supplements the protective properties of normal colostrum against E. coli F5 (K99) adhesin.

Pack size: 1, 6, 12, 24 or 48 60 ml bottles.

The manufacture, import, possession, sale, supply and/or use of the veterinary medicinal product may be prohibited in a Member State on the whole or part of their territory pursuant to national legislation. Any person intending to manufacture, import, possess, sell, supply and use the veterinary medicinal product must consult the relevant Member State's competent authority on the current animal health policies prior to the manufacture, import, possession, sale, supply and/or use.

For any information about this veterinary medicinal product, please contact the local representative of the marketing authorisation holder.