
C9555 A Survival Guide to Simulations


Sampling in Molecular Dynamics

Ivo Kabelka


Examples of studied processes




Membrane adsorption




Conformational ensemble




Ligand binding




Aggregation




Membrane properties



Vesicle fusion




Genome release



Pore closure

MUNI | CEITEC

Non-Equilibrium Simulations




Examples of non-equilibrium simulations


- Multiple starting configurations!
  - To capture a range of possible responses
- Time-dependent observables
  - Fluxes, structural changes, or relaxation times
- External perturbations
  - Drive the system out of equilibrium
  - Temperature or pressure gradients, electric field
- Types:
  - Steady-state: perturbations are constantly applied to study the response under constant conditions
  - Transient: relaxation of a system towards equilibrium state after a sudden perturbation

MUNI | CEITEC


Equilibrium Simulations




Membrane adsorption



Conformational ensemble



Ligand binding



Membrane properties

Examples of equilibrium simulations

- Single simulation or multiple replicas
  - Start from a well-equilibrated state
- Time-averaged observables
  - Temperature, pressure, …
- System at equilibrium
  - System remains in a steady state as there are no external perturbations
- Thermodynamic properties
  - Calculate enthalpy and free energy

MUNI | CEITEC

Ergodic Hypothesis

- Time averaging is equivalent to ensemble averaging
  - Only if the sampling of system's properties is done over a long enough time period
  - Ensemble average = average over many different configurations
- Importance sampling
  - Over a sufficiently long simulation, the system explores all accessible states

Molecular Dynamics Simulations


Equilibrium

1. Initial configuration
2. Equilibration
3. Production run
4. Analysis

Non-Equilibrium

1. Multiple initial configurations
2. Production runs
3. Analysis

G Protein-Coupled Receptor (GPCR)




PDB Code: 5MZP

- GPCRs are involved in a wide range of biological processes, including neurotransmission, hormonal regulation, immune response, and sensory perception (e.g., vision, taste, smell)
- Categorized into five main classes (A, B, C, F, and T), with Class A (Rhodopsin-like receptors) being the largest and most studied group
- Approximately 30–40% of all approved drugs target GPCRs
- The estimated 800–1,000 GPCRs represent approximately 3–5% of all human genes

MUNI | CEITEC

G Protein-Coupled Receptor (GPCR)



PDB Code: 6GDG

- GPCRs are involved in a wide range of biological processes, including neurotransmission, hormonal regulation, immune response, and sensory perception (e.g., vision, taste, smell)
- Categorized into five main classes (A, B, C, F, and T), with Class A (Rhodopsin-like receptors) being the largest and most studied group
- Approximately 30–40% of all approved drugs target GPCRs
- The estimated 800–1,000 GPCRs represent approximately 3–5% of all human genes

MUNI | CEITEC

G Protein-Coupled Receptor (GPCR)




PDB Code: 5MZP and 6GDG

- GPCRs are involved in a wide range of biological processes, including neurotransmission, hormonal regulation, immune response, and sensory perception (e.g., vision, taste, smell)
- Categorized into five main classes (A, B, C, F, and T), with Class A (Rhodopsin-like receptors) being the largest and most studied group
- Approximately 30–40% of all approved drugs target GPCRs
- The estimated 800–1,000 GPCRs represent approximately 3–5% of all human genes

MUNI | CEITEC


G Protein-Coupled Receptor (GPCR)



PDB Code: 5MZP and 6GDG

- GPCRs are involved in a wide range of biological processes, including neurotransmission, hormonal regulation, immune response, and sensory perception (e.g., vision, taste, smell)
- Categorized into five main classes (A, B, C, F, and T), with Class A (Rhodopsin-like receptors) being the largest and most studied group
- Approximately 30–40% of all approved drugs target GPCRs
- The estimated 800–1,000 GPCRs represent approximately 3–5% of all human genes

G Protein-Coupled Receptor (GPCR)




PDB Code: 5MZP and 6GDG

- GPCRs are involved in a wide range of biological processes, including neurotransmission, hormonal regulation, immune response, and sensory perception (e.g., vision, taste, smell)
- Categorized into five main classes (A, B, C, F, and T), with Class A (Rhodopsin-like receptors) being the largest and most studied group
- Approximately 30–40% of all approved drugs target GPCRs
- The estimated 800–1,000 GPCRs represent approximately 3–5% of all human genes


MUNI | CEITEC

Dose-Response Curve




MUNI | CEITEC

Dose-Response Curve



Hinz, S., et al. (2014). Journal of Pharmacology and Experimental Therapeutics, 349(3), 427-436.



MUNI | CEITEC

Free Energy


- Is a state function
  - Depends only on the initial and final states, not on the path taken
- Explains how molecules populate (distribute among) different energy states

- The change in Gibbs free energy, ΔG, for a process is given by:
  - $\Delta G = \Delta H - T\Delta S$
    ΔG: Change in Gibbs Free Energy
    ΔH: Change in enthalpy
    ΔS: Change in entropy
    Determined under constant temperature and pressure
  - A negative ΔG indicates that the process is spontaneous
  - Spontaneity does not imply speed; it only indicates thermodynamic favorability

Collective Variable (CV)

- A parameter that describes the progress of a process
  - Reduces the complexity of many-particle systems to a few key degrees of freedom
  - Must accurately represent the process or transition of interest
  - Selecting a CV can be non-trivial and often requires domain expertise
- Used to simplify the analysis of a trajectory or for enhanced sampling
  - In enhanced sampling, CVs facilitate efficient sampling of rare events or slow processes
- Examples of common collective variables:
  - Distance between atoms or molecules
  - Angles and dihedral angles
  - Coordination numbers
  - Radii of gyration

Collective Variable (CV)



Receptor activation

The outward swing of TM6 in a receptor can be described as either a distance or an angle


Ligand binding

Distance of a ligand from a receptor binding pocket


MUNI | CEITEC

Populations and Free Energy

Dose-response curve




Free energy




MUNI | CEITEC

Populations and Free Energy

Dose-response curve




Free energy




MUNI | CEITEC

Populations and Free Energy

Dose-response curve




Free energy




MUNI | CEITEC

Populations and Free Energy

Dose-response curve




Free energy




MUNI | CEITEC

Populations and Free Energy

Dose-response curve




Free energy



MUNI | CEITEC

Connecting Simulations and Experiment




Dissociation constant (K_D)

Experimentally determined using, e.g., surface plasmon resonance (SPR), isothermal titration calorimetry (ITC), or fluorescence spectroscopy.

MUNI | CEITEC

Connecting Simulations and Experiment



Dissociation constant (K_D)


$$K_D = \frac{[L][R]}{[LR]}$$

Experimentally determined using, e.g., surface plasmon resonance (SPR), isothermal titration calorimetry (ITC), or fluorescence spectroscopy.

Dissociation constant K_D is calculated from the concentrations of free ligand, receptor, and ligand-receptor complex, representing the equilibrium populations of different states.


MUNI | CEITEC

Connecting Simulations and Experiment




Dissociation constant (K_D)

Experimentally determined using, e.g., surface plasmon resonance (SPR), isothermal titration calorimetry (ITC), or fluorescence spectroscopy



$$K_D = \frac{[L][R]}{[LR]}$$


Dissociation constant K_D is calculated from the concentrations of free ligand, receptor, and ligand-receptor complex, representing the equilibrium populations of different states.




$$\Delta G^\circ = RT \ln K_D$$

Relationship between the dissociation constant K_D and standard Gibbs free energy ΔG°.

MUNI | CEITEC




Dissociation constant (K_D)




Dissociation constant (K_D)

- Weakly binding ligand
  - Higher K_D
  - Less negative free energy
- Strongly binding ligand
  - Lower K_D
  - More negative free energy
- Commonly reported as pKD
  - $pK_D = -\log_{10} K_D$
  - Higher pK_D indicate stronger binding
  - If K_D = 1 μM, then pK_D = 6
  - If K_D = 1 nM, then pK_D = 9




Dissociation constant (K_D)




Dissociation constant (K_D)




MUNI | CEITEC



Dissociation constant (K_D)




Dissociation constant (K_D)




Inhibition constant (K_I)

- Inhibition constant, K_I
  - Binding affinity via ligand displacement
  - Quantifies how effectively a molecule can prevent a ligand with known affinity from binding


MUNI | CEITEC




Dissociation constant (K_D)




Dissociation constant (K_D)




Inhibition constant (K_I)



ΔG_{CFF}



ΔG_{ZMA}




$\Delta\Delta G = \Delta G_{ZMA} - \Delta G_{CFF}$

MUNI | CEITEC


Practical Example 1

Differences in ligand affinities for a receptor in its active versus inactive conformation




The distance of a ligand from a receptor binding pocket can be used as a collective variable to calculate the free energy of binding*.


* If all possible poses, orientations, and conformations are not sampled, the result will **NOT** represent the binding free energy but rather the free energy along the chosen coordinate.



MUNI | CEITEC




$$\Delta\Delta G = \Delta G_A - \Delta G_I$$



$$\Delta\Delta G = \Delta G_A - \Delta G_I$$

Practical Example 2


Ligand-induced change in relative populations of receptors in active and inactive conformations




The swing* of TM6 is used to describe the receptor activation.


* Simple collective variables are often insufficient for accurately capturing protein conformational changes using enhanced sampling methods.

Fleetwood, O., et al. (2021). eLife, 10, e60715




MUNI | CEITEC






$$\Delta\Delta G = \Delta G_{CFF} - \Delta G_{Apo}$$



$$\Delta\Delta G = \Delta G_{ZMA} - \Delta G_{Apo}$$


MUNI | CEITEC

Thermodynamic Cycle



MUNI | CEITEC

Thermodynamic Cycle





MUNI | CEITEC


Ergodic Hypothesis

- Time averaging is equivalent to ensemble averaging
  - Only if the sampling of system's properties is done over a long enough time period
  - Ensemble average = average over many different configurations
- Importance sampling
  - Over a sufficiently long simulation, the system explores all accessible states

Sampling of Receptor Conformations







MUNI | CEITEC


Convergence Tests


- Increase simulation length
  - Monitor if time-averaged quantities change
- Block averaging
  - Divide a trajectory into blocks of equal length and calculate averages for each block
  - Stratification / strata
- Multiple starting conditions
  - Simulations should yield similar averages
- Check for symmetries


Convergence Tests

- Increase simulation length
  - Monitor if time-averaged quantities change
- Block averaging
  - Divide a trajectory into blocks of equal length and calculate averages for each block
  - Stratification / strata
- Multiple starting conditions
  - Simulations should yield similar averages
- Check for symmetries

Sampling of Receptor Conformations










MUNI | CEITEC

Sampling of Receptor Conformations










MUNI | CEITEC

Sampling of Receptor Conformations










MUNI | CEITEC

Sampling of Receptor Conformations










MUNI | CEITEC

Sampling of Receptor Conformations










MUNI | CEITEC

Sampling of Receptor Conformations










MUNI | CEITEC

Sampling of Receptor Conformations










MUNI | CEITEC

Sampling of Receptor Conformations










MUNI | CEITEC

Sampling of Receptor Conformations










MUNI | CEITEC

Sampling of Receptor Conformations










MUNI | CEITEC

Sampling of Receptor Conformations










MUNI | CEITEC

Sampling of Receptor Conformations










MUNI | CEITEC

Sampling of Receptor Conformations









MUNI | CEITEC


Convergence Tests

- Increase simulation length
  - Monitor if time-averaged quantities change
- Block averaging
  - Divide a trajectory into blocks of equal length and calculate averages for each block
  - Stratification / strata
- Multiple starting conditions
  - Simulations should yield similar averages
- Check for symmetries

Sampling of Receptor Conformations










MUNI | CEITEC

Sampling of Receptor Conformations










MUNI | CEITEC

Sampling of Receptor Conformations










MUNI | CEITEC

Sampling of Receptor Conformations








MUNI | CEITEC

Sampling of Receptor Conformations



MUNI | CEITEC

MUNI | CEITEC

Thank you for your attention!