LITERATURE MAP · STATUS 22 SEPTEMBER 2026

Condensates in twisted-bilayer optical lattices

Superfluidity across the moiré pocket crossover: how a weakly interacting condensate responds as the lowest band flattens

Václav Svoboda
Wolfgang Pauli Institute (WPI), Vienna

Abstract. Twisted-bilayer optical lattices put two spin states of a Bose gas into square lattices rotated against each other and couple them with a microwave field, so that the Rabi coupling plays the part of interlayer tunnelling in twisted bilayer graphene. One experiment has been published, on ^{87}Rb at a twist of 5.21°; it reports two superfluid regimes and a superfluid–insulator transition that moves with the coupling. This literature map collects what is known about how such a condensate stays superfluid as its lowest band flattens. It brings together the experiment and the lattice and continuum theory written about it, the theory of superfluid weight and quantum geometry developed for magic-angle graphene, recent work on condensates in flat bands, and the measurements of the superfluid fraction in modulated Bose gases. For a square bilayer with uniform coupling there is no magic angle: the band flattens because the atoms localise in the AA registry pockets, and the pockets are an atomic-limit band, for which the flat-band theory leaves no geometric superfluid weight (the uniform-density no-go statement itself covers only uniform bands). The map argues that the open physics lies in the crossover, where band mass, density inhomogeneity and Josephson tunnelling between pockets compete, and it lists the calculations that would settle it. Every number traces to a checked source or to the project's own single-particle band calculation, which is marked as such.

Keywords: twisted-bilayer optical lattice; moiré superlattice; Bose–Einstein condensate; superfluid fraction; quantum metric; flat band; Rabi coupling; Josephson array

At a glance

10	28	8	161	161	22
chapters	pages of text	figures	sources	sources checked	open questions

Status: complete draft, first edition. Literature status 21 September 2026; checks of sources, numbers and quotations carried out on 21–22 September 2026 with AI assistance; see "How this map was made" on page i.

The problem

Twisted-bilayer optical lattices bring moiré physics into a Bose gas. In the one experiment realised so far [1], two spin states of ^{87}Rb sit in square lattices rotated by 5.21° and a microwave field couples them, in the role that interlayer tunnelling plays in twisted bilayer graphene. The experiment reports a moiré density pattern, two distinct superfluid regimes and a superfluid–insulator transition that shifts with the coupling. The question is how the superfluidity of a weakly interacting condensate evolves as the lowest band of such a lattice flattens.

It matters because it tests, for bosons and without disorder, strain or Coulomb interaction, how far the quantum-geometric picture of magic-angle graphene carries over, on the only moiré quantum gas with published data. It is hard because in a square bilayer with uniform coupling the band flattens by localisation into AA registry pockets rather than by interference at a magic angle. In the crossover window $\Omega/J \approx 5$–25, which overlaps the experiment's range ($\Omega/J \approx 3$–12 at $V_0 = 4\,E_R$; about 16–40 along path III), conventional (m/m^*), geometric (condensate metric), density-inhomogeneity (Leggett) and inter-pocket Josephson contributions to the superfluid weight compete, and inter-pocket tunnelling can fall below accessible temperatures.

Established: uniform coupling gives no magic angle and acts as a scalar moiré potential proportional to Ω; a single flat band at a time-reversal-invariant momentum with uniform Bloch-state density cannot be superfluid for bosons within Bogoliubov theory; lattice models and Monte Carlo find clustered, glassy phases and thin superfluid domains that thermal broadening strongly depletes. Not yet found: Bogoliubov spectra, sound and the superfluid-fraction tensor of the Rabi-coupled condensate, its bosonic quantum geometry, finite temperature, and a trapped simulation compared with the data. Progress would be a quantitative account of superfluid fraction, sound and depletion across the crossover that reproduces the two superfluid regimes and the shifted transition.

How this map was made

Scope, method and conventions

Scope. Bose gases in twisted-bilayer optical lattices (experiment, lattice and continuum theory), flat-band bosons and quantum geometry, measurements and definitions of the superfluid fraction, the electronic moiré context, numerical methods and the research field; fermionic and photonic analogues only where they bear on the bosonic question.

Literature cutoff. 21 September 2026. Preprints are included and marked as such; published versions are cited where they were found.

How it was compiled. The starting documents were five project documents and a two-page proposal naming about 100 sources. Web searches covering 2011–2026 extended them on twisted-bilayer optical-lattice theory, flat-band bosons, superfluid-fraction measurements, electronic moiré, numerical methods and the research field. We then looked up every source in a primary record (arXiv or ar5iv, publisher page, code repository) in eleven checking rounds with AI assistance. Of the 161 sources the map cites, 56 were confirmed or corrected, 105 partly verified and 0 not verifiable.

Numbers. Each number and quotation traces to a line of the project's facts ledger. Quotations are verbatim from the checked source; numbers confirmed only in part carry "as reported" or "as quoted".

Conventions. $E_R = h \times 3.67\,\mathrm{kHz}$ for $a = 395\,\mathrm{nm}$ [1]; J single-layer nearest-neighbour hopping; Ω Rabi coupling (Ω_R in the experiment's notation); L_M moiré period; m commensurate index, $\tan(\theta/2) = 1/m$. The tag [computed for this map] marks numbers reproduced from the project's single-particle band-structure grid (no interactions, trap or temperature).

In one page: the field and the argument

Moiré physics reached quantum gases in 2023, when a ^{87}Rb condensate was loaded into two square lattices twisted by 5.21° and coupled by a microwave field [1]. The experiment found "two forms of superfluid and a modified superfluid to insulator transition". Magic-angle graphene suggests reading this through flat bands and quantum geometry [2–4]. For this lattice we argue, in five steps, that the reading has to change.

1. **The lattice has no magic angle.** In twisted square bilayers flat bands need a spatially varying interlayer coupling with symmetry-enforced cancellation [5]; a uniform Rabi coupling acts instead on the lowest band as a scalar moiré potential proportional to Ω [6]. The band flattens because the atoms localise in the AA registry pockets: at $m = 8$, $V_0 = 8\,E_R$ the width falls to 0.25 of its uncoupled value at $\Omega/J \approx 16$ and to 0.004 at $\Omega/J \approx 32$, where the participation ratio (fraction of sites the band-bottom state occupies) is 0.008 and the Wannier spread $0.022\,L_M^2$ [computed for this map].

2. **Thesis 1: geometric superfluidity is excluded there.** For bosons, "There can therefore be no superfluidity in a single flat band at a TRIM where the Bloch state has uniform densities" (TRIM: time-reversal-invariant momentum), and stable two-dimensional flat-band superfluidity "requires at least three bands within Bogoliubov theory" [7] (preprint). A band of isolated pockets is the atomic limit, where "no current is possible." [4]. Huhtinen *et al.* conclude that "flat-band superfluidity is much more difficult to achieve than flat-band superconductivity." [7]

3. **The open physics is the crossover.** Between a dispersive band and isolated pockets, band mass, density inhomogeneity (the Leggett bounds [8, 9]) and tunnelling between pockets all carry superfluid weight. No calculation of this competition for the Rabi-coupled condensate was found in the literature checked for this map. The tools exist in simpler geometries: sound anisotropy and Leggett bounds in modulated two-dimensional gases [10–12], Josephson oscillations in a supersolid [13].

4. **Thesis 2, working hypothesis: SF-II is a Josephson array of AA pockets.** The experiment's second regime is described as "superfluid domains embedded in a gapped insulator" [1]. At $V_0 = 8\,E_R$ its moiré-visibility threshold $\Omega_R > 0.5\,E_R$ sits at $\Omega/J \approx 16$, where the lowest band detaches and has lost 75 % of its width [computed for this map]. The hypothesis fails if the superfluid fraction exceeds the pocket prediction (which lies below Leggett's bound), if sound scales as U (the flat-band signature [14]), or if the threshold does not track Ω/J as V_0 changes.

5. **Progress** would be a two-component Gross–Pitaevskii calculation of the superfluid-fraction tensor, sound and depletion across the crossover, with Leggett brackets and a trap, that reproduces the two superfluid regimes and the shifted transition [1, 10, 12]. A finite-temperature treatment would show whether inter-pocket tunnelling survives at the experiment's temperature, which the paper, as read for this map, does not state.

Where quantum geometry still enters

The condensate metric at the condensation momentum sets sound and stability in the bosonic theory [7, 14], so it remains a diagnostic of how far the crossover has progressed. Band touchings [15] and detuned couplings with Hofstadter-type spectra [16] are where a genuinely geometric window could still open; chapter 3 explains these ideas for Bose-gas readers.

Contents

The problem i

How this map was made i

In one page: the field and the argument ii

1 The topic in brief 1

What a twisted-bilayer optical lattice is, what the one experiment saw, and which variants of the idea have been proposed.

2 Foundations 3

The geometry of the twisted supercell, the Rabi-coupled Gross–Pitaevskii model, how the lowest band turns into pockets, and the definitions of the superfluid fraction used in the rest of the map.

3 Context: electronic moiré and quantum geometry for Bose-gas physicists 7

How flat moiré bands in graphene become superconducting through quantum geometry, and which parts of that account survive for a condensate in a twisted optical lattice.

4 Data, benchmarks and verification 10

What the one experiment measures, how a superfluid fraction is measured and computed in modulated Bose gases, and which cross-checks and pitfalls a calculation for twisted-bilayer lattices has to respect.

5 State of the art 12

One experiment, about two dozen theory papers written around it or its relatives, and a clear pattern in what they treat: single-particle bands and ground states, rarely excitations, temperature or dynamics.

6 Position: geometric superfluidity is excluded; the pocket crossover is the open problem 16

Why a uniformly coupled square bilayer cannot hold a geometric flat-band superfluid, and why the experiment's second superfluid is best read as a Josephson array of AA pockets.

7 Around the topic 19

Neighbouring systems that share one ingredient with the twisted-bilayer condensate: moiré patterns for light, quasiperiodic lattices, dipolar bilayers, coupled 1-D condensates and moiré magnets.

8 Research and industry perspective 21

Who works on twisted optical lattices and flat-band superfluids, which codes and budgets exist, and where the neighbouring superconductivity research stands in 2026.

9 Open questions 24

Twenty-two questions in five themes, from effective models of the pockets to finite temperature and geometry, followed by directions that no study checked for this map has taken up.

10 Reading paths and outlook 27

Where to start reading, depending on the field one comes from, and what the next experiment or the first superfluid-fraction calculation would change.

References **29**

Research connected with the WPI and Vienna **37**

A Glossary **39**

Terms from cold-atom physics, electronic moiré materials and quantum geometry, as this map uses them.

How to read this map

Twenty minutes. The problem, the one-page summary, then the key-points boxes at the end of each chapter.

New to electronic moiré and quantum geometry. Chapter 3, then chapter 6.

Looking for a research question. Chapters 6 and 9.

Boxes: key points, what a result means for the reader, open questions. Numbers tagged [computed for this map] come from the project's own band-structure grid.

1 The topic in brief

What a twisted-bilayer optical lattice is, what the one experiment saw, and which variants of the idea have been proposed.

Twisted bilayer graphene (TBG) is two graphene sheets rotated by a small angle and coupled by interlayer tunnelling; near a magic angle its lowest moiré bands become flat and the material turns superconducting [2, 3, 17]. A twisted-bilayer optical lattice (TBOL) copies the construction with cold atoms. The two "layers" are two internal states of the same atoms, each trapped in its own square lattice, the lattices are rotated against each other, and a microwave or Raman field couples the states in the place of interlayer tunnelling [1, 18]. Figure 1.1 follows the idea from graphene to what is measured, and Table 1.1 lists the variants.

1.1 The platform

The one experiment so far used a Bose–Einstein condensate (BEC) of ^{87}Rb with "state $|1\rangle \equiv |F = 1, m_F = 1\rangle$ and state $|2\rangle \equiv |F = 2, m_F = 0\rangle$" [1]. Each state feels one square lattice. The two lattices are formed at "$\lambda_1 = 790.02\,\mathrm{nm}$ and $\lambda_2 = 788.28\,\mathrm{nm}$", both tune-out wavelengths, which a companion preprint studied with Kapitza–Dirac scattering [19]. The lattice constant is $a = 395\,\mathrm{nm}$ and the recoil energy is $E_\mathrm{R} = h \times 3.67\,\mathrm{kHz}$; the experiment writes E_r, and we write E_R throughout. With a twist of $\theta = 5.21°$ (uncertainty about 0.05°) the measured moiré period is $4.35\,\mu\mathrm{m}$, eleven lattice constants. The condensate holds up to 5×10^5 atoms, and an accordion lattice with spacing tunable from 12 to $3\,\mu\mathrm{m}$ compresses it into a pancake with "$\omega_z \approx 2\pi \times 1\,\mathrm{kHz}$". A 6.8 GHz microwave couples the two states with a Rabi coupling Ω_R of up to $1\,E_\mathrm{R}$; at $V_0 = 4\,E_\mathrm{R}$ it was calibrated as $2\pi \times 893\,\mathrm{Hz} = 0.243\,E_\mathrm{R}$, or $\Omega_\mathrm{R}/t = 2.07$ in units of the hopping t. After ramps of the lattices (30 ms) and the microwave (10 ms) and a hold of up to 40 ms, the gas is imaged in situ or after 18 ms of expansion [1]. As read for this map, the paper does not state the temperature, the radial trap frequency or the atom number in the pancake.




Figure 1.1: From twisted bilayer graphene to what a twisted-bilayer optical lattice measures. Top row: the chain of the platform [1]; solid arrows follow it from idea to observables, and dashed arrows show where ideas from electronic moiré systems are carried over. Orange: where the electronic moiré ideas of chapter 3 are borrowed. Aqua: the quantity that links the bands to the observed phases and that has not been calculated for this platform in the literature checked for this map.

1.2 The four regimes

The paper reports "two forms of superfluid and a modified superfluid to insulator transition". Its labels are defined as follows: "SF, SF-II, MI, and I refer to superfluid, superfluid only with short-range coherence, Mott insulator, and insulator." [1] In SF, time of flight shows peaks at the primary lattice momenta and at the moiré momenta. SF-II is described as "superfluid domains embedded in a gapped insulator"; as the project documents read the data, the moiré peaks

are gone there while the primary peaks remain. In I the filling is integer but modulated on the moiré scale. The phase diagram extends to $24\,E_R$. Along path I the coupling is fixed at $\Omega_R = 0.24\,E_R$ and the lattice is deepened, which gives SF $\to$ SF-II $\to$ MI. Along path III the depth is fixed at 8 or $14\,E_R$ and the coupling is raised; the moiré visibility "presents the threshold behavior and emerges at $\Omega_R > 0.5\,E_r$". The paper models the gas with a two-component Gross–Pitaevskii equation (GPE) with SU(2)-symmetric interactions $g_{11} = g_{22} = g_{12}$ and with a single-site Gutzwiller map of a Bose–Hubbard model, at the commensurate stand-in angle $2\arctan(1/22)$ and zero detuning [1]. The same group's later experiment used an untwisted honeycomb lattice [20], and no second twisted-lattice experiment was found for this map.

1.3 What "flat band" means here

In TBG the band flattens by interference: tunnelling paths cancel while the states stay spread over the moiré cell (chapter 3). A uniform microwave coupling cannot do this in a square bilayer, because flat bands there need a spatially varying interlayer coupling with symmetry-enforced cancellation [5], and a uniform Ω acts on the lowest band as a scalar moiré potential proportional to Ω [6] (preprint). In the paper's own strong-coupling limit the lower dressed state feels $(V_1 + V_2)/2 - \Omega_R$, the average of the two lattices [1], which is deepest where their sites coincide, in the AA registry regions. So a flat band here is an array of weakly coupled AA pockets, the atomic limit, not a band held flat by interference.

The parameter that places the experiment on this scale is Ω/J, with J the single-layer hopping. At $V_0 = 4\,E_R$, where according to the project documents the moiré and superfluid data were taken, $\Omega_R \le 1\,E_R$ means $\Omega/J \le 12$ and the lowest band keeps at least 93 % of its uncoupled width at $m = 8$ [computed for this map]; at $m = 12$ it keeps 59 %, and the experiment's angle, $m = 22$, was not computed. At $V_0 = 8\,E_R$ the path-III threshold $\Omega_R = 0.5\,E_R$ is $\Omega/J \approx 16$, where the band detaches from the next one and has lost 75 % of its width [computed for this map]. The definitions and the numbers are in chapter 2.

1.4 Variants of the platform

Table 1.1 sorts the proposals by how the moiré pattern enters. Most use two internal states with a uniform coupling, on square or honeycomb lattices. For square lattices Paul, Recher and Santos write the uniform coupling as an interlayer hopping that decays with the distance between sites and find Bose-glass-like "mobility islands" even at commensurate angles [21]; their earlier work found multifractal single-particle states [22]. With a detuning, the imbalanced limit "can be exactly mapped to Hofstadter's original model plus a long-range hopping term" [16]. On honeycomb lattices a uniform coupling does produce magic flat bands [23, 24]. Other variants drop the tunnelling and let interactions generate the moiré pattern [25, 26], sum two rotated lattices for a single component [27, 28], rotate one lattice continuously in time [29], or move to spin-1 condensates in twisted trilayer lattices [30] and Floquet-modulated interactions [31] (preprints). Photonic moiré lattices share the question of localisation in a moiré potential [32, 33].

Key points

- One experiment: ^{87}Rb, two spin states in square lattices twisted by 5.21°, microwave coupling up to $1\,E_R$, four regimes SF, SF-II, MI and I [1].
- Uniform coupling has no magic angle in a square bilayer; the band flattens by localisation in AA pockets [5, 6].
- Variants differ in how the moiré pattern enters; only one has data.

Table 1.1: Variants of the twisted-bilayer platform. "Uniform" means the coupling does not vary in space. Particles as stated in the sources or, where marked, in the project documents.

Variant	Coupling	Particles	What is studied	Sources
square lattices, uniform Ω	microwave (or Raman), uniform	bosons (^{87}Rb)	experiment: moiré density, two superfluids, shifted transition; lattice theory: clusters, mobility islands	[1, 18, 21]
honeycomb spin-twisted lattices	uniform Ω	fermions; atoms; fermions (docs)	magic flat bands for $\theta \lesssim 6°$; topological vs trivial flat bands; fractional Chern insulator (preprint)	[23, 24, 34]
moiré without a twist	spatially modulated Raman	fermions	quasi-flat bands with Dirac cones	[35]
interaction-induced moiré	none (contact interaction only)	bosons	induced and "interlocked" Mott insulators, Bose-glass-like phases, quenches	[25, 26]
summed twisted potential	one component, no coupling	bosons	continuous-space QMC: thin superfluid domains, Bose glass; tight-binding fits	[27, 28]
dipolar twisted bilayer	no tunnelling; dipolar interaction	hard-core bosons	QMC: insulating islands surrounded by superfluid at 5.21°	[36]
one-dimensional moiré	bichromatic lattice	bosons	moiré flat bands broadened by weak repulsion; "local supersolid" (preprint)	[37, 38]
photonic moiré	not applicable	photons	localisation and delocalisation of light; solitons	[32, 33]

2 Foundations

The geometry of the twisted supercell, the Rabi-coupled Gross–Pitaevskii model, how the lowest band turns into pockets, and the definitions of the superfluid fraction used in the rest of the map.

Numbers tagged [computed for this map] in this chapter come from the project's single-particle plane-wave bands at the commensurate index $m = 8$, without interactions, trap or temperature; the experiment's $m = 22$ was not computed.

2.1 Commensurate supercells and the moiré period

Two square lattices of constant a, rotated by $\pm\theta/2$, share a common supercell when $\tan(\theta/2) = 1/m$ for an integer m. The supercell is a square of side $\sqrt{m^2+1}\,a$ and holds m^2+1 sites of each layer; the moiré period is $L_\mathrm{M} = a/[2\sin(\theta/2)] = \sqrt{m^2+1}\,a/2$. At the experiment's $\theta = 5.21°$ this gives 4.35 μm, or 11 a, and the paper's theory uses the commensurate stand-in $\theta = 2\arctan(1/22) = 5.205°$, a supercell of 22 a [1]. The map's own numbers are for $m = 8$ ($\theta = 14.25°$, $L_\mathrm{M} = 4.03\,a$) and, in part, $m = 12$ ($\theta = 9.53°$, $L_\mathrm{M} = 6.02\,a$) [computed for this map]. We speak of AA registry where the sites of the two lattices coincide and of AB registry where a site of one sits at a plaquette centre of the other (Fig. 2.1). Without coupling each layer's lowest band is folded into the small supercell Brillouin zone; at $m = 8$, $V_0 = 8\,E_\mathrm{R}$ the lowest folded band has a width $W_1(0) = 0.0347 \times 8J$ [computed for this map].

2.2 The Rabi-coupled Gross–Pitaevskii equation

The mean-field model of the experiment is a two-component GPE in which each component $\sigma = 1, 2$ feels its own lattice and the Rabi coupling converts one into the other ($\bar{\sigma}$ is the other component):

$$i\hbar\,\partial_t\psi_\sigma = \Big[-\frac{\hbar^2\nabla^2}{2m} + V_\sigma + V_T + \sum_{\sigma'} g_{\sigma\sigma'}|\psi_{\sigma'}|^2\Big]\psi_\sigma + \frac{\hbar\Omega}{2}\,\psi_{\bar{\sigma}}, \qquad V_\sigma(\mathbf{r}) = -V_0 \sum_{j=x,y} \cos^2(k_L\,\mathbf{e}_{\sigma j}\cdot\mathbf{r}). \tag{1}$$

Here $k_L = \pi/a$, the unit vectors $\mathbf{e}_{\sigma j}$ of lattice σ are rotated by $\pm\theta/2$, V_T is the trap, and we quote $\hbar\Omega$ in units of E_R and call it Ω. The paper quotes $\Omega_\mathrm{R}/t = 2.07$ at $\Omega_\mathrm{R} = 0.243\,E_\mathrm{R}$ [1], so its t is $0.117\,E_\mathrm{R}$ at $V_0 = 4\,E_\mathrm{R}$; that is the value of the asymptotic formula, which lies 36 % above the exact band hopping $J = 0.0862\,E_\mathrm{R}$ used here [computed for this map]. Thresholds in this map are quoted in Ω/J with the exact J, and we take Ω equal to the experiment's Rabi frequency Ω_R, which enters Eq. (1) as $\hbar\Omega/2$. The experiment's model is quasi-two-dimensional with SU(2)-symmetric interactions $g_{11} = g_{22} = g_{12}$ [1]. Without interactions and trap, Eq. (1) reduces to the 2×2 Hamiltonian whose bands the project computed. Without lattices a finite coupling "changes the first order demixing phase transition" of the mixture "to a second order" one [39]. With lattices it mixes, in addition, two different potentials. For strong coupling the lower dressed state sees $-(\hbar^2/2m)\nabla^2 + (V_1 + V_2)/2 - \Omega_\mathrm{R}$ [1]. The average of the two lattices is deepest at AA registry, so this is where the atoms collect.

2.3 A scalar moiré potential, and no magic angle

Wang, Gao and Shi project the coupled problem on the lowest band at small angle and obtain two decoupled branches in a moiré potential proportional to Ω [6]:

$$H_{\mathrm{eff},\pm} = \frac{p^2}{2m_\mathrm{eff}} \pm V_m(\mathbf{r}), \qquad V_m(\mathbf{r}) = 8\pi^{3/2}\,\Omega\,l_\mathrm{ho}^3 \sum_{\mathbf{G}} e^{-G^2 l_\mathrm{ho}^2}\, e^{i(R^{1/2} - R^{-1/2})\mathbf{G}\cdot\mathbf{r}}, \tag{2}$$

with l_ho the harmonic length of a lattice site, $\mathbf{G}$ the reciprocal lattice vectors and R the rotation by θ. The derivation is written for a three-dimensional quasi-moiré crystal and holds "When the twist angle is small such that $\theta \cdot l_\mathrm{ho}^{-1} \ll 1$"; applying it to the two-dimensional square bilayer is a step taken by the project, not by the preprint. The amplitude grows linearly with Ω and carries a Gaussian form factor; nothing in Eq. (2) singles out an angle. Kariyado and Vishwanath reach the same conclusion from symmetry: in twisted Bravais bilayers, flat bands need a spatially varying interlayer coupling whose matrix elements vanish for particular momenta and shifts, "an interference effect where symmetry forces exact cancellation", and a uniform coupling lacks this (our paraphrase) [5]. Paul, Recher and Santos write the same coupling in lattice language as an interlayer hopping $t_\perp e^{-|\mathbf{R}_{1j} - \mathbf{R}_{2j'}|^2/4l_0^2}$ that is largest between coinciding sites [21].

2.4 Pocket formation

The lowest band flattens when the potential localises the atoms at its minima (Fig. 2.2; all at $m = 8$, $V_0 = 8\,E_\mathrm{R}$) [computed for this map]. At $\Omega/J = 3.2$ the band keeps 98 % of its folded width, the band-mass ratio at Γ is $m/m^* = 0.28$, the participation ratio (PR) of the band-bottom density is 0.38 and the coarse-grained AA/AB density ratio 2.4. At $\Omega/J = 16.2$ the width is $0.25\,W_1(0)$, a gap of $0.0007\,E_\mathrm{R}$ has opened to the second band, $m/m^* = 0.12$, PR = 0.059 and AA/AB = 110; at $\Omega/J = 32.4$ the four numbers are 0.004, 0.003, 0.008 and 3.7×10^3, with a gap of $0.010\,E_\mathrm{R}$. The width halves at $\Omega/J \approx 12$ for $V_0 = 8\,E_\mathrm{R}$, at ≈ 9 for $12\,E_\mathrm{R}$ and at ≈ 35 for $4\,E_\mathrm{R}$ [computed for this map], so the threshold is set by Ω/J with a depth-dependent prefactor, not by the angle.

The dependence on m is not settled. At $V_0 = 4\,E_\mathrm{R}$, $\Omega = 1\,E_\mathrm{R}$ the width ratio is 1.02 ($m = 6$), 0.93 ($m = 8$) and 0.59 ($m = 12$) [computed for this map], so the project documents' claim of a nearly angle-independent threshold for $m \geq 6$ fails at $4\,E_\mathrm{R}$; $m = 22$ was not computed.





Figure 2.1: Schematic of pocket formation (not computed; drawn at $m = 6$, $\theta = 18.9°$, for visibility). Dots: the two rotated square lattices, which coincide at AA registry and alternate at AB. (a) For small Ω/J the condensate density is spread over the moiré cell with a mild AA excess. (b) For large Ω/J it sits in AA pockets connected by a weak tunnelling J_M, and the lowest band is an array of nearly isolated pockets.



single-particle bands; no interactions, trap or temperature; computed for this map from grid.csv

Figure 2.2: Pocket crossover in the single-particle bands [computed for this map]: (a) width of the lowest band relative to its folded value, (b) participation ratio of the band-bottom state, (c) band-mass ratio m/m^* at Γ, against Ω/J for $V_0 = 4$, 8 and 12 E_R at $m = 8$ (faint: $m = 6$). Shaded: the window $\Omega/J = 5$–25; dotted: the experiment's moiré-visibility threshold $\Omega_\mathrm{R} = 0.5\,E_\mathrm{R}$ at $V_0 = 8\,E_\mathrm{R}$ [1], $\Omega/J = 16.2$. No interactions, trap or temperature; the experiment's $m = 22$ was not computed. J is the exact single-layer hopping; the participation ratio is normalised so that a plane wave gives 1.

2.5 Superfluid fraction

The superfluid fraction f_s is the part of the gas that does not follow a slow motion of the potential. In a phase-twist calculation we impose a velocity v and read f_s off the energy, $E(v) = E(0) + f_\mathrm{s} N m v^2/2$ for $v \to 0$ [9]; in an anisotropic lattice f_s is a tensor [40]. Leggett bounded f_s from above by the density modulation alone [8] and defined it for an arbitrary many-body system at zero temperature [41]. In two dimensions, for a twist along x, Pérez-Cruz,

Astrakharchik and Massignan give the pair of bounds

$$f^{+} = \frac{1}{\bar{n}} \frac{1}{\left\langle 1/\langle n\rangle_{r_\perp}\right\rangle_x}, \qquad f^{-} = \frac{1}{\bar{n}} \left\langle \frac{1}{\langle 1/n\rangle_x} \right\rangle_{r_\perp}, \tag{3}$$

where $\langle\cdot\rangle_x$ and $\langle\cdot\rangle_{r_\perp}$ average along and across the twist, and they find that "The bracketing is tighter for a 2D optical potential than for a grid of columns" [9]. Experiments use sound. With c_x and c_y the sound velocities along and across a one-dimensional modulation, $f_{s,x} = c_x^2/c_y^2$ and $c_x^2 = f_{s,x}/(m\kappa)$, with κ the compressibility; for a weakly interacting condensate in a one-dimensional periodic potential Leggett's bound becomes an identity [10]. A Bragg-phonon measurement found that "the speed of sound saturates Leggett's bound" [11]. For a condensate in one Bloch state, Taylor and Zaremba derive the sound speed $s = \sqrt{(\bar{n}/m^*)(\partial\mu/\partial\bar{n})}$ with $(m^*)^{-1} = \hbar^{-2}\partial^2\tilde{\varepsilon}/\partial k^2$ the inverse band mass [42]. Comparing with the sound relation above gives $f_s = m/m^*$ for a Bloch condensate (our combination of the two results). The computed m/m^* is a single-particle number, and once interactions smooth the density it is only an estimate of f_s.

2.6 The Bose–Hubbard mapping and the energy scales

Lattice models replace Eq. (1) by a Bose–Hubbard model with hopping J and on-site interaction U; the experiment quotes $U/t = 1.67$ [1]. The exact single-layer hopping, taken from the one-dimensional Mathieu band, is $J = 0.0862$, 0.0308 and $0.01225\,E_R$ at $V_0 = 4$, 8 and $12\,E_R$ [computed for this map]. The asymptotic formula used in the project code, $J \approx (4/\sqrt{\pi})\,V_0^{3/4} e^{-2\sqrt{V_0}}\,E_R$ with V_0 in units of E_R, from the deep-lattice limit discussed by Zwerger [43], is too large by 36, 22 and 16 % at these depths [computed for this map], so every Ω/J in the literature depends on which J was used. The single-band description has limits of its own. In continuous-space quantum Monte Carlo (QMC) for hexagonal potentials the bands match tight binding for $V_0 \gtrsim 5\,E_R$, yet at $15\,E_R$ "the Bose–Hubbard approximation fails to describe but the first Mott lobe" [44].

With $E_R = h \times 3.67\,\mathrm{kHz}$ for ^{87}Rb at $a = 395\,\mathrm{nm}$ [1], the experiment's couplings $\Omega_R = 0.24$–$1\,E_R$ at $4\,E_R$ correspond to $\Omega/J = 2.8$–11.6 [computed for this map]. Between pockets the tunnelling can be much smaller. Johnstone *et al.* ran continuous-space worm QMC for single-component bosons in a summed twisted potential at $k_B T = 0.01$–$0.04\,E_R$, in boxes 5 and 15 moiré periods wide. They report maximal hoppings in the thin superfluid domains between 10^{-11} and $10^{-3}\,E_R$ and find that "thermal broadening strongly depletes the SF domains" [27]. The experiment's temperature is not stated. The measured scattering lengths we found are for another pair of ^{87}Rb states, $|F=1, m_F=-1\rangle$ and $|F=2, m_F=+1\rangle$, $a_{12} = 98.006(16)\,a_0$ and $a_{22} = 95.44(7)\,a_0$ [45], with earlier values of $97.66\,a_0$ and $95.0\,a_0$ respectively, as quoted there, and attributed by the project documents to [46]; values for the experiment's pair $|1,1\rangle$, $|2,0\rangle$ were not found for this map.

Key points

- Commensurate supercells: $\tan(\theta/2) = 1/m$, side $\sqrt{m^2+1}\,a$, $L_M = a/[2\sin(\theta/2)]$; the experiment is $m = 22$, the map's numbers are $m = 8$.
- A uniform Ω acts as a scalar moiré potential $\propto \Omega$; the band flattens by localisation in AA pockets, with its width halved at $\Omega/J \approx 12$ for $8\,E_R$ [computed for this map].
- The superfluid fraction has three working definitions: phase twist, Leggett brackets and sound anisotropy; for a Bloch condensate $f_s = m/m^*$.
- Exact and asymptotic J differ by 16–36 %, and inter-pocket tunnelling can fall far below J.

3 Context: electronic moiré and quantum geometry for Bose-gas physicists

How flat moiré bands in graphene become superconducting through quantum geometry, and which parts of that account survive for a condensate in a twisted optical lattice.

We borrow most of the vocabulary for twisted-bilayer optical lattices (TBOL) from twisted bilayer graphene (TBG). This chapter covers how a moiré band becomes flat, why a flat band can still carry superfluid weight, and what changes for a condensate. Figure 3.1 traces the argument; Table 3.1 translates the terms.

3.1 Flat bands by interference: the magic angle

In the Bistritzer–MacDonald (BM) continuum model the Dirac cones of the two graphene layers are coupled by an interlayer tunnelling w that varies with the moiré pattern [2]. One dimensionless number controls the low-energy bands, $\alpha = w/(vk_\theta)$, where v is the Dirac velocity and k_θ the moiré wavevector. The renormalised velocity is $v^*/v = (1-3\alpha^2)/(1+6\alpha^2)$, and in the authors' words, "For a discrete set of magic angles the velocity vanishes, the lowest moire band flattens". The first magic angle is $\theta \approx 1.05°$. Nothing localises here. Tunnelling paths interfere destructively while the states stay extended over the moiré cell, and this is exactly where the optical lattice will differ.

Experiments then found correlated insulators at half filling of the flat bands [47] and superconductivity at "twist angles of about 1.1°—the first 'magic' angle", with "a critical temperature of up to 1.7 kelvin" at "a carrier density of about 10^{11} per square centimetre" [3, 17].




Figure 3.1: From Bloch states to superfluid weight: the shared trunk (top), then the fermion branch (pairing) and the boson branch (condensation at $\mathbf{k}_c$; [14] and the preprint [7]).

3.2 Superfluid weight from quantum geometry

In a dispersive band $D_s \propto n/m^*$, so a flat band with $m^* \to \infty$ looks like a dead end, but it is not one. The quantum metric $g(\mathbf{k})$ measures how fast the Bloch state $|u_\mathbf{k}\rangle$ changes with $\mathbf{k}$, and the quantum distance $1 - |\langle u_\mathbf{k}|u_{\mathbf{k}'}\rangle|^2$ is its finite-step version. Peotta and Törmä showed that pairing in a flat band still produces a superfluid weight, linear in the attraction ("$D_s \propto U$ in a flat band", as the abstract puts it) and controlled by this geometry [48]. For TBG the superfluid weight has "a topological lower bound proportional to the pairing gap" [49]. From the other side, Verma, Hazra and Randeria bound the low-energy optical weight, and so the stiffness, of a trivial flat

band by the Wannier spread [50]: well-localised orbitals mean little stiffness. Huhtinen *et al.* then showed that the physically relevant quantity is the minimal quantum metric, which "is related to the superfluid weight in isolated flat bands", and that "a non-isolated flat band can actually be beneficial" [15]. The review [4] states the limiting case plainly: "a trivial flat band is equivalent to an insulator in the atomic limit of large atom-atom separation", and there "no current is possible."

The stiffness of magic-angle TBG has now been measured. Tanaka *et al.* combined "d.c. transport and microwave circuit quantum electrodynamics to directly measure the superfluid stiffness of superconducting MATBG through its kinetic inductance" and found it "much larger than expected from conventional Fermi liquid theory" [51]; a companion measurement did the same in twisted trilayer graphene [52, 53]. A BM-model decomposition attributes 22–26 % of D_s at charge neutrality to quantum geometry [54] (preprint).

3.3 Condensates in flat bands

A condensate has no pairing gap. It occupies one Bloch state at momentum $\mathbf{k}_c$, and in a flat band the interaction energy Un_0 is the only scale. Julku, Bruun and Törmä worked this out for the kagome lattice [14]. The projected Bogoliubov spectrum is

$$E_1(\mathbf{k}_c+\mathbf{q}) = \frac{Un_0}{3} D(\mathbf{q}), \qquad D(\mathbf{q}) = \sqrt{1-|\langle u_1(\mathbf{k}_c+\mathbf{q})|u_1(\mathbf{k}_c-\mathbf{q})\rangle|^2}, \tag{4}$$

so the excitations are set by the quantum distance, the sound velocity $c_s = (2Un_0/3)\sqrt{g(\mathbf{k}_c)}$ is linear in U rather than in $\sqrt{U}$, and the depletion as $U \to 0$ "is determined by the quantum geometry only". The companion general theory [55] assumes, as the project documents summarise it (the paper was not read for this map), a uniform condensate density over the orbitals, an isolated band and projected interactions. Julku, Salerno and Törmä added "orbital-position-independent generalizations of the quantum metric" and showed that D_s must be taken at fixed total particle number [56].

The 2026 preprint by Huhtinen *et al.* sharpens the picture [7]. For uniform-density condensates at weak coupling, $D_1 + D_2^{\mathrm{ndeg}} = (2Un_0^2/M)\,\mathcal{M}^D(\mathbf{k}_c)$, where $\mathcal{M}^D$ is the condensate quantum metric: a local quantity at $\mathbf{k}_c$, not an integral over the Brillouin zone (BZ). The global geometry enters the fluctuation term D_3', which in their model "is positive only at very small interactions, and turns negative around $U \approx 0.13t$". At a time-reversal-invariant momentum (TRIM), "There can therefore be no superfluidity in a single flat band at a TRIM where the Bloch state has uniform densities." And a large integrated metric does not help: on the checkerboard lattice, "even though the integrated quantum metric diverges due to the band touching, condensation and superfluidity remain impossible." The authors conclude that stable two-dimensional flat-band superfluidity "requires at least three bands within Bogoliubov theory" and that "flat-band superfluidity is much more difficult to achieve than flat-band superconductivity." A companion preprint phrases the criterion in real space: "frameworks consisting of triangles with nonzero area are promising for condensation, whereas … square frameworks indicate condensation in a single mode is impossible" [57].

Two further preprints add to this. For a two-band condensate, "in the case of U(2) invariant interactions, the superfluid density is zero" [58]; the TBOL experiment is modelled with SU(2)-symmetric interactions $g_{11} = g_{22} = g_{12}$ [1], but no source checked tests the result there. Burgher *et al.* relate the sound velocity in quasi-flat bands to a "quantum geometric kernel" [59].

Table 3.1: Dictionary between twisted bilayer graphene and twisted-bilayer optical lattices.

TBG	TBOL	What it means
layer	spin state, each with its own lattice [1]	the "layers" are internal states, not planes in space
interlayer tunnelling w	Rabi coupling Ω (microwave)	uniform in space for a microwave drive
twist angle, magic angle	twist θ, index m; no magic angle for uniform Ω [5]	flattening is set by Ω/J, not by θ
AA / AB stacking	AA / AB registry regions	where the two lattices' sites coincide or alternate
BM continuum model	scalar moiré potential $\propto \Omega$ [6]	one effective band in a moiré potential
superfluid weight D_s	superfluid fraction f_s (a tensor)	response to a phase twist
quantum metric	condensate quantum metric at $\mathbf{k}_c$ [7]	local geometry, not the BZ integral
Cooper pairs	condensate in one Bloch state	what the superfluid is made of
Coulomb interaction	contact interaction g	short range, tunable, no screening
transport, kinetic inductance [51]	sound velocity, phase twist, Leggett bounds [10, 11]	how the stiffness is measured

3.4 What carries over, and what does not

Both systems ask what carries superfluid weight once the kinetic energy of a moiré band is gone. TBG samples come with twist-angle disorder, strain and long-range Coulomb repulsion, and the superfluid is made of pairs. The optical lattice is clean, the interaction is a contact term, and the superfluid is one macroscopic state whose density can be imaged; its superfluid fraction follows from sound anisotropy, $f_{s,x} = c_x^2/c_y^2$, and from Leggett's bounds on that density [10–12].

The real difference lies in how the band flattens. Flat bands in twisted square bilayers need a spatially varying interlayer coupling whose symmetry forces exact cancellations [5]. A microwave Rabi coupling is uniform, and projected on the lowest band it acts as a scalar moiré potential of amplitude proportional to Ω (derived for 3-D quasi-moiré crystals and applied to 2-D by the project) [6]. The band flattens by localisation in the AA pockets, not by interference, and at $m = 8$, $V_0 = 8\,E_R$ the coarse-grained AA/AB density ratio is already 110 at $\Omega/J \approx 16$ [computed for this map]. The bosonic geometric results assume the opposite, a uniform density.

What this means for Bose-gas physicists

A flat-band condensate has a clear signature: sound linear in U and a depletion that stays finite as $U \to 0$ [14]. The metric that counts is the local one at $\mathbf{k}_c$, not the BZ integral of the fermionic literature. In a square lattice with uniform coupling the flat limit is the atomic limit, so the geometric formulas become diagnostics of the crossover, not a mechanism.

Open

No calculation of the condensate metric of the Rabi-coupled bands was found for this map. Open: what the uniform-density results say for AA pockets, whether [58] transfers, and how the geometric term compares with the Leggett bound.

Key points

- TBG flattens by interference, a uniform-Ω TBOL by localisation [2, 5, 6].
- Bosons: one flat band at a TRIM with uniform density is not superfluid [7].
- AA pockets break uniform density: the open physics is in the crossover.

4 Data, benchmarks and verification

What the one experiment measures, how a superfluid fraction is measured and computed in modulated Bose gases, and which cross-checks and pitfalls a calculation for twisted-bilayer lattices has to respect.

4.1 How the experiment identifies its phases

Meng *et al.* use two probes [1]. In situ, at 50× magnification, they resolve the moiré density pattern, whose period of 4.35 μm is eleven lattice constants of $a = 395$ nm. After 18 ms of time of flight (TOF), imaged state by state, the momentum distribution has peaks at the primary lattice momentum and at the moiré momentum; because the moiré period is $11a$, the second set of peaks sits at 1/11 of the first. The phases are defined by the visibilities of the two sets. The superfluid (SF) shows both. The second regime, SF-II, keeps the primary peaks and loses the moiré ones; the figure caption calls it "superfluid only with short-range coherence", and the Extended Data describe "superfluid domains embedded in a gapped insulator". A Mott insulator (MI) and a moiré-modulated insulator (I) complete the diagram. Along path I the coupling is held at $\Omega_R = 0.24\,E_\mathrm{R}$ while the depth rises (SF → SF-II → MI). Along path III the coupling rises at a depth in the Mott region, and "the visibility at the moiré momenta presents the threshold behavior and emerges at $\Omega_R > 0.5\,E_r$". The experiment writes E_r; we write E_R throughout.

The numeric anchor points of the phase diagram are in the Extended Data of the Nature version only, not in the arXiv PDF, and this map could not confirm them. Temperature, radial trap frequency and the atom number per pancake were not found in the pages read.

4.2 How the superfluid fraction is measured

Three groups have measured f_s in modulated Bose gases since 2023 (Table 4.1). Chauveau *et al.* imposed a 1-D modulation of period $d = 3.93(4)$ μm and amplitude up to 80 nK on a 2-D ^{87}Rb gas. They compared Leggett's formula, evaluated on the measured density, with the sound anisotropy $f_{s,x} = c_x^2/c_y^2$; for a 1-D modulation of a weakly interacting condensate "the bound reduces to an identity" [10]. Tao, Zhao and Spielman excited phonons by Bragg scattering in a 1-D lattice of period $a = 266$ nm (depth up to $10\,E_\mathrm{R}$, healing length $\xi = 170(20)$ nm, $N \approx 2\times10^5$) and found that "the speed of sound saturates Leggett's bound" [11]. Rabec *et al.* moved to a 2-D triangular lattice ($d = 6.0(1)$ μm, $\mu_0/k_B = 45(2)$ nK, $T < 20$ nK, depths up to $4.7\,\mu_0$, as reported). They obtained the tensor in two ways, statically from the density profile and the continuity equation, and dynamically from compressibility and sound, and both agree with Gross–Pitaevskii equation (GPE) simulations and with the Leggett bounds of the measured profiles [12]. In a dipolar supersolid, Biagioni *et al.* read f_s off the Josephson oscillations of single cells and found "a relatively large sub-unity superfluid fraction" [13]. One preprint tests Leggett's bound for strongly interacting bosons (not checked for this map) [60], and another proposes vortex necklaces as a probe [61].

4.3 How the superfluid fraction is computed

Four routes are in use. The phase twist gives $E(v) = E(0) + f_\mathrm{s} N m v^2/2$ for $v \to 0$ [9, 41]. Leggett's brackets give an upper bound f^+ and an explicit lower bound f^- from the density alone, and "The bracketing is tighter for a 2D optical potential than for a grid of columns" [8, 9]; a 2026 preprint studies how the brackets depend on angle in composite potentials [62]. Sound gives $c_x^2 = f_\mathrm{s}/(m\kappa)$ [10] and, in a lattice, $s = \sqrt{(\bar{n}/m^*)\,\partial\mu/\partial\bar{n}}$ [42, 63]. The band route ties nonclassical translational inertia to the effective mass and makes f_s a rank-2 tensor [40].

Table 4.1: Measured superfluid fractions in modulated Bose gases; none is moiré or Rabi-coupled.

Group / platform	Modulation	Period	Method	Compared with
Dalibard/Beugnon, 2-D ^{87}Rb [10]	1-D, $V_0 \leq 80\,\mathrm{nK}$	3.93(4) μm	density + Leggett; c_x^2/c_y^2	Leggett bound (exact in 1-D)
Spielman, ^{87}Rb [11]	1-D lattice, $\leq 10\,E_\mathrm{R}$	266 nm	Bragg phonons, sound speed	Leggett bound, GPE
Dalibard/Beugnon, 2-D ^{87}Rb [12]	triangular, $\leq 4.7\,\mu_0$	6.0(1) μm	continuity eq.; compressibility + sound	GPE, Leggett bounds
Modugno, dipolar supersolid [13]	self-organised	–	Josephson oscillations of cells	sub-unity f_s

Quantum Monte Carlo (QMC) takes $f_\mathrm{s} = \Upsilon/\rho$ from winding numbers, which sample the response to a twist [27, 64]. The unitary Fermi gas has its own Leggett test [65].

4.4 Cross-method checks

For the square Bose–Hubbard model, cluster Gutzwiller extrapolated from clusters of up to 4×4 sites gives $(J/U)_c = 0.0596(4)$, against the QMC value 0.0597 [66, 67]. The single-site Gutzwiller used in the experiment's theory has no such check for the moiré case. Continuous-space QMC shows where the lattice model ends. The bands match tight binding for $V_0 \gtrsim 5\,E_\mathrm{R}$, but at $15\,E_\mathrm{R}$ "the Bose–Hubbard approximation fails to describe but the first Mott lobe", and density-assisted tunnelling appears beyond it [44]. The Gaussian-overlap interlayer coupling of the two-layer tight-binding models [18, 21] reproduces the lowest bandwidth to 5–8 % for $\Omega/J \leq 3$, but it is 30 % low at $\Omega/J \approx 10$ and $10\times$ low at $\Omega/J \approx 32$ (project computation; the underlying table was not re-read for this map) [computed for this map]. Neural quantum states on 20×20 sites [68] and infinite projected entangled-pair states (iPEPS) [69] give variational references.

4.5 Pitfalls

Classical-field methods need "the number of field modes in the Wigner simulation [to] be smaller than the number of particles" [70], and "all c-field region modes" have to be "highly occupied" [71, 72]. In strongly interacting regimes near the Mott lobes the truncated Wigner approximation (TWA) is fragile. In a preprint, Nagao and Yunoki report that the SU(3) discrete TWA breaks down for $t \gtrsim 0.1\,\hbar/J_\mathrm{fin}$ in strongly interacting 2-D Bose–Hubbard quenches and deviates from experiment "even at relatively short times"; their cluster version uses 2×2 clusters on 20×20 sites [73].

Kibble–Zurek (KZ) exponents mislead. In a 3-D ^{87}Rb lattice the measured exponent is 0.31 ± 0.03, against the KZ value 3/4 [74]. For ^{39}K the exponents lie well below the KZ value and depend on the final interaction [75]. On the numerical side, iPEPS needs bond dimension $D = 11$–14, at a cost $\propto D^8$, to test the KZ prediction $\hat{\xi} \propto \tau_\mathrm{Q}^{0.40}$ [76]. Energy scales can also become vanishingly small. In twisted single-component potentials the superfluid domains carry tunnelling J_max between 10^{-11} and $10^{-3}\,E_\mathrm{R}$, and thermal broadening at $k_B T = 0.01$–$0.04\,E_\mathrm{R}$ "strongly depletes" them [27].

Key points

- Phases are defined by TOF visibilities at two momenta; numeric anchors only in the Nature Extended Data [1].
- f_s is measured via Leggett bounds, sound and Josephson oscillations, never yet in a

moiré gas [10–13].

- Gaussian-overlap couplings fail beyond $\Omega/J \approx 10$ [computed for this map]; TWA near Mott lobes, KZ exponents and nK tunnelling trap dynamics [27, 73, 74].

5 State of the art

One experiment, about two dozen theory papers written around it or its relatives, and a clear pattern in what they treat: single-particle bands and ground states, rarely excitations, temperature or dynamics.

5.1 The experiment

Meng *et al.* loaded a ^{87}Rb Bose–Einstein condensate (BEC) with "up to 5×10^5 atoms" into two square lattices of constant $a = 395\,\mathrm{nm}$, each felt by one spin state, $|1\rangle = |F{=}1, m_F{=}1\rangle$ and $|2\rangle = |F{=}2, m_F{=}0\rangle$ [1]. The lattice beams sit at the tune-out wavelengths 790.02 and 788.28 nm, which a companion metrology paper measured by Kapitza–Dirac scattering [19]. The recoil energy is $E_\mathrm{R} = h \times 3.67\,\mathrm{kHz}$ (written E_r in the paper). The twist is $\theta = 5.21^\circ$ with an uncertainty of about 0.05°, the measured moiré period is $4.35\,\mu\mathrm{m}$, and the paper's theory uses the commensurate angle $\theta = 2\arctan(1/22)$. An accordion lattice at 532 nm, tunable from 12 to 3 μm, makes the gas quasi-2-D with $\omega_z \approx 2\pi \times 1\,\mathrm{kHz}$. A 6.8 GHz microwave couples the layers with Ω_R up to $1\,E_\mathrm{R}$. At $4\,E_\mathrm{R}$ the calibrated Rabi frequency is $2\pi \times 893\,\mathrm{Hz}$, so $\Omega_R/t = 2.07$, and the paper quotes $U/t = 1.67$. The Rabi oscillations decay at $1200\,\mathrm{s}^{-1}$. In the sequence both lattices are ramped up in 30 ms and the microwave in 10 ms, after which the gas is held for up to 40 ms. The phase diagram spans depths up to $24\,E_\mathrm{R}$.

The paper's theory has three parts: a continuum band calculation with the strong-coupling limit $H^-_\mathrm{eff} \approx -(\hbar^2/2m)\nabla^2 + (V_1 + V_2)/2 - \Omega_R$, a two-component GPE with SU(2) interactions $g = g_{11} = g_{22} = g_{12}$, and a single-site Gutzwiller map. The headline is "two forms of superfluid and a modified superfluid to insulator transition". No second twisted-lattice experiment was found for this map; the same group has since studied a vortex at the Dirac point of an untwisted honeycomb lattice [20].

5.2 Single-particle continuum theory

Five results fix the band structure. First, the preprint by Wang, Gao and Shi projects the Rabi-coupled Hamiltonian on the lowest band at small angle and obtains a scalar moiré potential $V_m(\mathbf{r}) = 8\pi^{3/2}\Omega\, l_\mathrm{ho}^3 \sum_\mathbf{G} e^{-G^2 l_\mathrm{ho}^2} e^{i(R^{1/2} - R^{-1/2})\mathbf{G}\cdot\mathbf{r}}$, derived for a 3-D quasi-moiré crystal and valid for $\theta\, l_\mathrm{ho}^{-1} \ll 1$ [6]. Its amplitude is proportional to Ω times a Gaussian form factor. Second, Wan, Gao and Shi add a detuning δ. In the imbalanced limit ($\delta \gg V_0, \Omega$) they find an on-site term $v = \Omega^2 V_0/\delta^2$ that "can be exactly mapped to Hofstadter's original model plus a long-range hopping term"; the twist plays the role of the flux, and the spectra are fractal [16]. Third, Kariyado and Vishwanath showed that flat bands in twisted Bravais bilayers come from a spatially varying interlayer coupling whose symmetry forces exact cancellations, a mechanism that a uniform coupling lacks [5].

Fourth, hexagonal lattices behave differently. Luo and Zhang found that for spin-twisted hexagonal lattices "All twist angles $\theta \lesssim 6^\circ$ can become magic that support gapped flat bands", at a coupling that grows almost linearly with θ [23]. Sui *et al.* report for honeycomb lattices an isolated topological flat band at a critical coupling $\Omega_R \approx 0.11\,E_\mathrm{R}$ at $\theta \approx 4.4^\circ$, and trivial flat bands at strong coupling [24]. Fifth, for incommensurate twists the interlayer coupling acts

as effective disorder, and above a critical value part of the spectrum becomes multifractal [22]. Three further results complete the single-particle picture: the original proposal of twisted state-dependent potentials [18], a fermionic "twistronics without a twist" with moiré-modulated Raman coupling [35], and a continuously rotated lattice that acts as a position-dependent drive with ring-shaped localised states [29].

5.3 Lattice models with interactions

Paul, Recher and Santos wrote the two-layer Bose–Hubbard model with Gaussian interlayer hopping and density–density terms, $t_\perp(j,j') = t_\perp e^{-|\mathbf{R}_{1j}-\mathbf{R}_{2j'}|^2/4l_0^2}$, and treated it with cluster Gutzwiller after "site clusterization", in which pairs with coupling above $t_{\perp,\mathrm{cr}} = 0.03\,t_\perp$ (commensurate) or $0.06\,t_\perp$ (incommensurate) form clusters [21]. They find "Mott-like phases … resulting typically in a density wave modulation" and the "formation of mobility islands", where Bose-glass phases may occur. As the project documents summarise it, the runs use $t_\perp = U$, $V_\perp = 4U$ and $T = 0$, and dynamics is left open.

Zeng, Zhu and He removed the interlayer tunnelling altogether, so that the layers couple only through contact interactions, at angles 36.87°, 22.62° and 7.63°. Their Gutzwiller calculations give induced Mott insulators, an "interlocked" Mott insulator "with spontaneously formed moiré lattice" and self-localised Bose-glass-like phases [25]. The arXiv versions carry two titles, "Interaction induced moiré systems" and "Dynamical Moiré Systems". A follow-up finds that interactions can delocalise a disconnected Bose glass into a connected superfluid at low filling, with reentrance at higher filling, and it also studies quenches [26]. For spin-1 condensates in a twisted trilayer, depth quenches create vortex pairs that inherit the moiré periodicity [30].

5.4 Continuous space: quantum Monte Carlo and GPE

Johnstone *et al.* ran continuous-space worm QMC for single-component bosons in the summed twisted potential (depths 3–6 E_R, $\theta \approx 28.07°$ and an incommensurate 27.13°, $k_BT = 0.01$–$0.04\,E_\mathrm{R}$). Thin superfluid domains with J_max between 10^{-11} and $10^{-3}\,E_\mathrm{R}$ separate density-wave and Mott plateaus, and the incommensurate case leaves "a prominent Bose glass phase" [27]. A companion paper fits four forms of tight-binding model [28].

Real-time GPE work is scarce. Fan *et al.* solve the Meng-type coupled GPE (hexagonal lattices, $\Omega = 1\,E_\mathrm{R}$, $V_0 = 4\,E_\mathrm{R}$, $\theta = 9.43°$, $N = 2\times10^4$, trap $2\pi\times(20,20,1000)$ Hz) with a Floquet-modulated scattering length and obtain quasicrystal density patterns [31]. A preprint shows that the nonlinearity alone can self-organise a twisted-bilayer structure in a two-component BEC [77]. Gap solitons and vortices in moiré potentials form a separate strand; according to the abstracts, their stability is studied by linear or nonlinear methods on localised states [78–82].

5.5 One dimension and fermions

In 1-D, repulsion broadens the moiré flat band of a condensate in a $\mathcal{PT}$-symmetric bichromatic lattice [37]. A density-matrix renormalisation group (DMRG) preprint finds a "local supersolid" in a moiré-modulated Bose–Hubbard chain, with "A vanishing global structure factor in the thermodynamic limit, while the local structure factor remains finite" [38]. For fermions, Yang *et al.* (preprint, September 2026) take honeycomb lattices with uniform Ω and, as reported, a cross-lattice admixture $w \approx 0.05$. The single-particle spectrum "hosts a pair of quasi-degenerate nearly flat moiré bands"; Hartree–Fock interactions "spontaneously break time-reversal symmetry" and produce a Chern band; exact diagonalisation then finds a fractional Chern insulator (FCI), reported at hole filling $\nu_h = 1/3$, with a many-body gap "of order $10^{-3}\,E_R$" [34].




Figure 5.1: Coverage of twisted-bilayer optical-lattice theory and the experiment. Rows are studies (p: preprint), columns are, in order, Rabi/uniform Ω coupling, bosons, square lattice, interactions, excitations/Bogoliubov, superfluid fraction, $T > 0$, real-time dynamics/ramps, trap, experiment. Filled: treated; half: partly treated or single-particle only; empty: not treated, or not stated in the abstracts and pages read for this map.

5.6 What the coverage shows

Figure 5.1 shows the pattern: most TBOL studies treat single-particle bands or ground states at $T = 0$. The finite-temperature work (Johnstone) is for a single-component summed potential, and the dynamics (Fan, Ding, Li) concerns quenches of the scattering length, the interaction or the depth, not of Ω. The graphene literature and the superfluid-fraction experiments, placed in time in Figure 5.2, ran in parallel with the TBOL work [2, 3, 7, 10–12, 14, 48, 51, 83].

5.7 What is openly available

The worm QMC code for lattice bosons [64] is public under GPL-3.0, with 26 commits and no releases [84]. No TBOL data set has been released. The Nature paper states that "All data




Figure 5.2: Timeline 2011–2026 of the work this map draws on: graphene and quantum geometry (2011–2021), then the experiment (marked), the superfluid-fraction measurements and the theory papers (2023–2026). Salamon *et al.* and Yang *et al.* treat fermions.

generated or analysed during this study are included in this published article", with further data "available from the corresponding authors upon reasonable request", and it has no code-availability statement [1]. The neural-quantum-state code for the 2-D Bose–Hubbard model of [68] is archived on Zenodo as v1.0.0 under the Apache-2.0 licence [85].

Not found for this map

No calculation of the following was found in the literature checked for this map:

- Bogoliubov spectra, sound velocities or the superfluid-fraction tensor of the Rabi-coupled condensate;
- the quantum geometry of its bosonic bands (only fermionic honeycomb bands [34]);
- finite-temperature or truncated-Wigner behaviour of the Rabi-coupled model;
- ramps or quenches of Ω;
- an independent trapped simulation compared with the Nature data.

Key points

- One experiment: ^{87}Rb, 5.21°, $\Omega \leq 1\ E_R$, two superfluid regimes, no follow-up found [1].
- Uniform Ω acts as a scalar moiré potential; magic angles appear only in hexagonal lattices [5, 6, 23, 24].
- Interacting theory is Gutzwiller at $T = 0$, or QMC for single-component relatives with thin, fragile superfluid domains [21, 25, 27].
- No released data; the worm code is GPL-3.0 [84].

6 Position: geometric superfluidity is excluded; the pocket crossover is the open problem

Why a uniformly coupled square bilayer cannot hold a geometric flat-band superfluid, and why the experiment's second superfluid is best read as a Josephson array of AA pockets.

Our claim covers square twisted-bilayer optical lattices (TBOL) with a spatially uniform Rabi coupling at zero detuning. It rests on single-particle band structures and on the published theory of condensates in flat bands, so honeycomb lattices, modulated couplings and detuned drives lie outside it. We argue one thesis and hold a second as a working hypothesis with stated falsifiers. Figure 6.1 separates what is fixed from what is open; Table 6.1 records how the framing of the project changed.

6.1 Thesis 1: the flat limit is the atomic limit

Symmetry. Kariyado and Vishwanath showed that flat bands in twisted Bravais bilayers, the square lattice included, arise from an interlayer coupling that varies in space: symmetry then forces the coupling matrix elements to cancel for specific momenta and displacements, "an interference effect where symmetry forces exact cancellation" [5]. A microwave Rabi coupling has no spatial structure, so this route is closed.

Effective theory. For a 3-D quasi-moiré crystal, Wang, Gao and Shi project the two-component Hamiltonian onto the lowest band and obtain a scalar moiré potential whose amplitude is proportional to Ω times a Gaussian form factor, valid for $\theta \cdot l_{\mathrm{ho}}^{-1} \ll 1$ [6] (preprint). If we apply this to the 2-D square bilayer (our step, not theirs), a uniform coupling becomes an ordinary potential with minima in the AA registry regions.

Computation. The single-particle bands bear this out [computed for this map]. At $m = 8$ and $V_0 = 8\,E_{\mathrm{R}}$, the width of the lowest band relative to its uncoupled value is $W_1/W_1(0) = 0.85, 0.66, 0.25, 0.043$ and 0.004 at $\Omega/J = 6.5, 9.7, 16, 23$ and 32. The indirect gap to band 2 opens between $\Omega/J \approx 10$ and 16, and the mass ratio m/m^* at Γ falls from 0.28 to 0.003 over $\Omega/J = 3.2$–32. The band flattens by localisation. The coarse-grained AA/AB density ratio is about 110 at $\Omega/J \approx 16$, and at $\Omega/J \approx 32$ the Wannier spread is $0.022\,L_{\mathrm{M}}^2$ and the participation ratio 0.008. These numbers contain no interactions, trap or temperature, and the experiment's angle ($m = 22$) was not computed.

Theory. The preprint of Huhtinen *et al.* treats condensates in Bloch states of uniform density. It gives $D_1 + D_2^{\mathrm{ndeg}} = (2Un_0^2/M)\,\mathcal{M}^D(\mathbf{k}_c)$, with the condensate quantum metric at the condensation momentum, and states that "There can therefore be no superfluidity in a single flat band at a TRIM where the Bloch state has uniform densities."; stable two-dimensional flat-band superfluidity "requires at least three bands within Bogoliubov theory" [7]. The pocket band is not uniform in density, so this statement does not apply to it directly. What excludes the geometric mechanism there is the atomic limit. The review of Törmä, Peotta and Bernevig states that "a trivial flat band is equivalent to an insulator in the atomic limit of large atom-atom separation" and that there "no current is possible." [4]. In an isolated flat band the relevant metric is the minimal one, and a non-isolated band can do better [15]; for fermions, the stiffness of a trivial flat band is bounded by the Wannier spread [50], which here is about 2 % of L_{M}^2. Two further results point the same way. A two-band condensate has zero superfluid density "in the case of U(2) invariant interactions" [58] (preprint), and the experiment is modelled with SU(2)-symmetric $g_{11} = g_{22} = g_{12}$ [1]. We conclude that the geometric mechanism cannot carry D_{s} in the collapsed regime. No calculation of the condensate metric for the Rabi-coupled bands

was found for this map, so the conclusion rests on the atomic-limit argument and not on a computed $\mathcal{M}^D$.




Figure 6.1: Position diagram. Left: results that hold independently of this map; middle: single-particle bands at $m = 8$, $V_0 = 8\,E_R$ (no interactions, trap or temperature; the experiment's $m = 22$ was not computed); right: what requires interacting calculations; bottom: the calculations and measurements that would decide.

Table 6.1: How the framing changed between the project's first proposal (v1) and this map (v2).

Question	v1 framing	v2 framing (this map)
how the band flattens	interference, as in TBG; a magic angle	localisation in AA pockets; no magic angle
control parameter	twist angle θ	Ω/J, with a V_0-dependent prefactor
what carries D_s	flat band plus quantum metric	competition of m/m^*, Leggett and Josephson terms across $\Omega/J \approx$ 5–25
role of geometry	the mechanism, centrepiece of the project	a diagnostic of the crossover
reading of SF-II	geometric flat-band superfluid	Josephson array of AA pockets (hypothesis)
central calculation	quantum metric of the flat band	f_s tensor by phase twist, Leggett brackets, sound

The crossover window matters because the experiment sits in it. At $V_0 = 4\,E_R$, where the moiré and superfluid data were taken, $\Omega_R \leq 1\,E_R$ corresponds to $\Omega/J \leq 12$ and $W_1 \geq 0.93\,W_1(0)$ at $m = 8$ [computed for this map], so the band is still dispersive. The threshold also moves with depth. At $V_0 = 12\,E_R$ the band has already lost 96 % of its width at $\Omega/J \approx 16$; at $4\,E_R$ it halves only near $\Omega/J \approx 35$ [computed for this map].

6.2 Thesis 2 (working hypothesis): SF-II is a Josephson array of AA pockets

The experiment describes its second superfluid regime as "superfluid domains embedded in a gapped insulator" [1]. We read these domains as the AA pockets of Thesis 1: the AB regions insulate, and tunnelling J_M between pockets carries phase coherence. Three lattice and Monte Carlo studies found related textures. Paul, Recher and Santos find Bose-glass-like "mobility islands" in a two-layer Bose–Hubbard model with Gaussian-overlap interlayer hopping, even at commensurate angles [21]. For dipolar bosons in a twisted bilayer at 5.21°, "a periodic structure of insulating islands is formed. These insulating islands are surrounded by a superfluid." [36]. This is the same geometry with the roles exchanged. Continuous-space Monte Carlo of a single-component twisted potential finds density-wave plateaus separated by thin superfluid domains,

with maximal tunnelling $J_{\max}$ between 10^{-11} and $10^{-3}\,E_{\mathrm{R}}$ and thermal broadening that "strongly depletes the SF domains" [27].

The experiment's moiré visibility at $V_0 = 8\,E_{\mathrm{R}}$ "emerges at $\Omega_R > 0.5E_r$" [1], where E_r is the paper's symbol for the recoil energy. At $m = 8$ this is $\Omega/J \approx 16$, where the lowest band detaches from band 2 and has fallen to $W_1 = 0.25\,W_1(0)$ [computed for this map]. The coincidence is suggestive, not a derivation, because it compares an interacting, trapped measurement at $m = 22$ with non-interacting bands at $m = 8$.

Falsifiers. The hypothesis fails if any of the following is observed or computed.

F1 The measured or computed superfluid fraction exceeds the value the pocket (Josephson-array) picture predicts, which lies well below Leggett's upper bound [8, 9].

F2 The sound velocity scales linearly with U, the signature of a geometric flat-band condensate [14], rather than as $\sqrt{U}$ with a Josephson-renormalised band mass [42].

F3 The visibility threshold does not track Ω/J as V_0 changes. The experiment reports its $0.5\,E_{\mathrm{R}}$ threshold for path III, run at 8 or $14\,E_{\mathrm{R}}$ [1]; extrapolating the grid, the single-particle collapse at $14\,E_{\mathrm{R}}$ would already sit near 0.15–$0.2\,E_{\mathrm{R}}$ [computed for this map] (no grid point at $14\,E_{\mathrm{R}}$). Either interactions or temperature shift it, or the pocket reading is wrong.

F4 The inter-pocket J_{M} at the relevant Ω lies far below any plausible $k_B T$, on the scale of the Monte Carlo $J_{\max}$ [27]; SF-II would then be a finite-size or interaction-restored effect, not a thermodynamic Josephson array. The experiment does not state its temperature.

Where quantum geometry still enters

Geometry stays in the problem as a diagnostic. The bare metric at Γ peaks at about $0.027\,L_{\mathrm{M}}^2$ as the band collapses and decays beyond it [computed for this map], and the condensate metric at $\mathbf{k}_c$ sets sound and stability in the bosonic theory [7, 14]. A genuinely geometric window could open at band touchings, where "a non-isolated flat band can actually be beneficial" [15], and under detuning, which maps onto "Hofstadter's original model plus a long-range hopping term" [16]. Honeycomb bilayers are different: uniform coupling there gives "magic" isolated flat bands for all $\theta \lesssim 6^\circ$ [23] and an isolated topological flat band at a weak critical coupling [24].

Key points

- Uniform Ω on a square bilayer gives no magic angle; the band flattens by AA localisation [5, 6].
- In the collapsed regime the band is in the atomic limit, where no geometric superfluid weight survives [4].
- The open physics is the crossover $\Omega/J \approx 5$–25, which overlaps the experiment's range ($\Omega/J \approx 3$–12 at $V_0 = 4\,E_{\mathrm{R}}$; about 16–40 along path III).
- Working hypothesis: SF-II is a Josephson array of AA pockets; four falsifiers are stated.

7 Around the topic

Neighbouring systems that share one ingredient with the twisted-bilayer condensate: moiré patterns for light, quasiperiodic lattices, dipolar bilayers, coupled 1-D condensates and moiré magnets.

None of these systems is a Rabi-coupled moiré Bose gas; each offers a tool or a measurement to borrow.

7.1 Photonic moiré lattices

Light in a photonic moiré lattice obeys, in the paraxial approximation, an equation of the same form as the single-particle Schrödinger equation, and with a Kerr nonlinearity the form of the GPE. In such lattices Wang *et al.* observed "the two-dimensional localization–delocalization transition of light" [32]. The same group showed that the twist angle controls soliton formation, with "threshold properties that are a pristine direct manifestation of flat-band physics" [33]. In nanostructured moiré superlattices, "the confinement mechanism of magic-angle lasers does not rely on a full bandgap but on the mode coupling between two twisted layers" [86]. Reviews cover moiré photonics and optoelectronics [87] and dielectric photonic moiré metamaterials [88]; a 2024 editorial surveys moiré physics beyond van der Waals materials [89], and the project documents cite a 2024 review of photonic and polaritonic moiré whose identity was not confirmed [90]. The closest analogue to our problem is condensation itself, since exciton polaritons condense "in a perovskite moiré flat band at room temperature" [91] (title; authors not obtained for this map). Polaritons are driven and dissipative, however, so they do not test the equilibrium superfluid fraction.

7.2 Optical quasicrystals and the 2-D Bose glass

The incommensurate limit of a twisted lattice is a quasiperiodic potential. Yu *et al.* reported "the experimental realization of the two-dimensional Bose glass using ultracold atoms in an eight-fold symmetric quasicrystalline optical lattice" [83]. A 2026 preprint from the same group quenches across the Bose-glass transition and finds that the characteristic timescales "are governed primarily by quasi-disorder strength in the Bose glass phase, and by tunneling strength in the superfluid phase" [92]. Both are single-component and without a Rabi coupling, but they show that a Bose glass in 2-D is measurable, which matters if SF-II is to be told apart from one.

7.3 Dipolar twisted bilayers

Zhang, Fan, Capogrosso-Sansone and Deng studied hard-core dipolar bosons in two twisted layers where "particles cannot hop between layers", with path-integral QMC using the two-worm algorithm and the superfluid stiffness as observable [36]. At the experiment's angle of 5.21° "a periodic structure of insulating islands is formed. These insulating islands are surrounded by a superfluid." The layers couple only through the long-range interaction, as in the contact-coupled model of [25], and the resulting picture of insulating pockets in a superfluid network is the inverse of the superfluid pockets in an insulator that SF-II suggests.

Electronic moiré materials have a structural relative of this domain picture. Twisted bilayer graphene reconstructs at the atomic scale into domains separated by domain walls, which also changes its electronic structure [93]. In twisted trilayer graphene, a moiré-of-moiré lattice, the same reconstruction produces unconventional tessellations of domains [94].

7.4 Coupled one-dimensional condensates in Vienna

The relative phase of two coupled condensates is the one-dimensional cousin of the interlayer phase in a TBOL. On an atom chip, "a pair of tunnel-coupled one-dimensional atomic superfluids" in thermal equilibrium "can be seen as a quantum simulator of the sine-Gordon model" [158]. With Raman tunnel coupling, the commensurate–incommensurate transition "can be … described by the Pokrovsky-Talapov model for the relative phase", with "a soliton staircase in the phase difference" [159]. Mennemann *et al.* studied relaxation in this extended bosonic Josephson junction: "stochastic classical field simulations using Gross-Pitaevskii equations in three spatial dimensions reproduce the main experimental findings", the relaxation persists "even at zero temperature", and it could not be explained by the sine-Gordon model [149]. The stochastic classical-field method behind that work has a longer Vienna history: fluctuations and stochastic processes in one-dimensional many-body systems [148], and, for the hydrodynamic regime, a dissipative form of generalised hydrodynamics with a numerical scheme [150]. The same groups wrote the time-splitting and boundary-layer numerics a TBOL Gross–Pitaevskii code would inherit: time splitting for the three-dimensional Pauli equation [152], an adaptive absorbing boundary layer for the nonlinear Schrödinger equation [153], fast Gaussian-sum and FFT–Kronecker evaluations of non-local potentials [154, 155], and an early numerical study of a Schrödinger–Poisson–Xα model [151]; their hard-constrained physics-informed networks [156, 157] are among the machine-learning tools chapter 9 lists. In coherently (Rabi) coupled condensates the coupling gaps the spin mode, and "finite spin-flipping coupling changes the first order demixing phase transition … to a second order" [39]; experiments studied ferromagnetism in an extended coherently coupled superfluid [95] and observed "bubble nucleation in isolated and coherently coupled atomic superfluids", a false-vacuum decay [96]. These systems have the two ingredients of a TBOL, a Rabi coupling and a spatially varying relative phase, but no lattices.

7.5 Magnetic moiré

Twisted magnets form moiré patterns in the spin texture. A 2025 review covers twisted CrI_3 with "moiré magnetism in nanoscale domains and periodic magnetization patterns", "skyrmion lattices and helical or spin spirals", and "moiré magnon excitations" [97]. Moiré magnons are the closest analogue of the Bogoliubov modes of a moiré condensate; no Bogoliubov spectrum of the latter was found for this map.

Key points

- Photonic moiré lattices share the GPE mathematics and show a localisation–delocalisation transition and flat-band soliton thresholds [32, 33].
- The 2-D Bose glass has been observed in an optical quasicrystal, the incommensurate relative of a TBOL [83, 92].
- Dipolar twisted bilayers at 5.21° give insulating islands in a superfluid [36].
- Coupled 1-D condensates show that relative-phase dynamics needs 3-D classical-field simulation, not sine-Gordon alone [149].

8 Research and industry perspective

Who works on twisted optical lattices and flat-band superfluids, which codes and budgets exist, and where the neighbouring superconductivity research stands in 2026.

Every statement below comes from group pages, code repositories, programme and company statements or the published record. Versions and licences are as stated on the pages, accessed on 21 September 2026, and statements by companies are marked "according to the company".

8.1 Platforms and groups

One laboratory has realised a twisted-bilayer optical lattice (TBOL) with a Bose gas, the Shanxi–Chicago collaboration of Jing Zhang and Cheng Chin [1], with later work on an untwisted honeycomb lattice [20] and, from Shanxi, theory of twisted honeycomb lattices [24]. The superfluid-fraction measurements a TBOL study would copy come from three groups that have not worked on twisted lattices (Table 8.1). The Santos group page lists one twisted-bilayer paper [98]; the affiliation of the fermionic proposal [34] was not checked.

Table 8.1: Groups active on the topic or on the methods it needs, with the papers cited in this map. Affiliations of the Palaiseau, Paris, Cambridge and Vienna groups follow the project documents and were not re-checked.

Group	Work relevant here	Papers
Shanxi / Chicago: J. Zhang, C. Chin, Z. Meng	the TBOL experiment; untwisted honeycomb lattice; from Shanxi, twisted-honeycomb theory	[1, 19, 20, 24]
Palaiseau (CPHT): L. Sanchez-Palencia	continuous-space QMC of twisted potentials	[27, 28, 44]
Hannover: L. Santos	TBOL lattice models, cluster Gutzwiller	[21, 22]
Aalto: P. Törmä	flat-band superfluid weight, quantum geometry	[4, 14, 48]
ETH Zürich: S. D. Huber	Bose condensation in flat bands	[7, 99]
Paris (Collège de France): J. Dalibard, J. Beugnon	f_s from Leggett bounds and sound	[10, 12]
NIST / JQI: I. B. Spielman	f_s from Bragg phonons in a 1-D lattice	[11]
Florence (CNR-INO): G. Modugno, with M. Fattori	f_s from Josephson oscillations of a supersolid	[13]
Cambridge: U. Schneider	2-D Bose glass in an optical quasicrystal; quenches	[83, 92]
Vienna (TU Wien, Atominstitut): J. Schmiedmayer	tunnel-coupled 1-D condensates	[149, 158, 159]
Y. Xu and co-workers	fermionic TBOL, fractional Chern insulators	[34]

8.2 Open codes and data

No TBOL data set has been released. The experiment's data statement reads "All data generated or analysed during this study are included in this published article." [1]. Table 8.2 lists the codes a TBOL study would draw on. The worm quantum Monte Carlo (QMC) code builds on ALPSCore and shows "near-linear scaling" in volume and inverse temperature [64]. As far as the pages state, none of the listed Gross–Pitaevskii equation (GPE) codes offers the Rabi-coupled twisted lattice as a ready model.

Table 8.2: Open codes, with licence and version as stated on the pages (accessed 21 September 2026).

Code	Use here	Licence	Version, status
worm QMC [64, 84]	lattice bosons, ρ_s from windings	GPL-3.0	no releases
ALPSCore [100]	QMC libraries	MIT	v2.3.3, May 2026
NetKet [101]	neural quantum states	Apache-2.0	3.22.4, Aug 2026
TeNPy [102]	tensor networks	Apache-2.0	v1.1.0, Nov 2025
ITensors.jl [103]	tensor networks	Apache-2.0	v0.9.30
lattice-bosons [68, 85]	neural states, Bose–Hubbard	not checked	v1.0.0, June 2025 (Zenodo)
XMDS2 [104]	stochastic PDEs, TWA	GPL-2.0	docs 2.2.2; news ends 2014
GPELab [105]	MATLAB GPE, stochastic GPE	none stated	none stated
GPUE [106]	CUDA GPE	not checked	JOSS 2018
GPU Python GPE [107]	PyTorch GPE, cavity QED	not checked	preprint 2024
quTARANG [108]	HPC GPE turbulence	MIT	CPC 2025
FourierGPE.jl [109]	FFT GPE	MIT	"no longer actively maintained"
GeneralizedGross-Pitaevskii.jl [110]	stochastic, GPU GPE	MIT	not checked
Wannier90 [111]	maximally localised Wannier functions	LGPL-2.1	v3.1.0, Feb 2024

8.3 Compute and cost, as quoted

For the Kibble–Zurek problem in the 2-D Bose–Hubbard model, infinite projected entangled-pair states need bond dimension $D = 11$–14, and "The dominant numerical cost of the NTU procedure scales as D^8" [76]. A spin-1 stochastic projected GPE study uses, as reported, 2.5×10^5 steady-state samples on a 512×512 grid [112]. The cluster truncated Wigner method runs 2×2 clusters on 20×20 sites [73], and a neural backflow ansatz reaches lattices up to 20×20 [68]. The continuous-space QMC of twisted potentials uses boxes of 5 and 15 moiré cells at $k_B T = 0.01$–$0.04\, E_R$ and gives no cost statement [27]. The project's band code took 7–35 s per (V_0, Ω) point at $m = 8$ and 15–108 s at $m = 12$ on 2 CPU cores [computed for this map]. No timing figure for a GPU GPE solver was found for this map.

8.4 Programmes and prizes

The EU Quantum Flagship project PASQuanS2, coordinated by Immanuel Bloch, was launched in April 2023 for seven years with a first phase of 3.5 years. It aims at "quantum simulators with at least 2,000 atoms and a path towards 10,000"; the page gives no EU amount [113]. The Austrian cluster of excellence quantA started on 1 December 2023 with "up to 70 million euros" over ten years for six partners, including TU Wien and the University of Vienna [161]. In the US, the NSF Quantum Leap Challenge Institutes round of 25 August 2026 gives more than $290 M to eight institutes over five years, "between about $28 and $37 million" each [114]; the first award to the Robust Quantum Simulation institute was $25 M in 2021 [115]. The ERC Consolidator grant QTWIST (Manchester, 1 April 2021 to 31 March 2027) receives €2,880,313 for twistronics; the PI is not on the page [116]. No ERC grant on superfluidity of quantum gases in moiré lattices was found for this map. The 2026 Kavli Prize in Nanoscience went to Eva Andrei, Pablo Jarillo-Herrero and Allan MacDonald "for foundational work that established the field of Twistronics", worth USD 1,000,000 [117].

8.5 Companies

According to the company, planqc builds the MAQCS system for the Leibniz Supercomputing Centre in Garching, a €20 M project funded by the German research ministry (BMBF), with "two independent cores of 1,000 qubits in total" that "will employ neutral atoms in optical lattices as qubits" [118]. QuEra's Aquila, a "field-programmable qubit array" in the authors' words, offers analog simulation on up to 256 neutral-atom qubits through Amazon Braket [119] (preprint). Lattice quantum simulation and atomtronics are reviewed in [120, 121]; no company offering moiré simulation, and no substantive commercialisation of twistronics, was found for this map.

8.6 Flat-band and quantum-geometric superconductivity

Magic-angle graphene superconducts at twist angles of about 1.1° with "a critical temperature of up to 1.7 kelvin" [3]. Its superfluid stiffness, measured through the kinetic inductance, is "much larger than expected from conventional Fermi liquid theory" [51], with a companion measurement in twisted trilayer graphene [52]; an earlier circuit measurement [122] is described in [54] as "approximately ten times larger than a conventional BCS estimate". The single-author preprint [54] attributes 22–26 % of D_s at charge neutrality to quantum geometry within the flat bands and about 55–58 % once remote bands are included (model calculation). Twisted WSe_2 superconducts at 3.5° and 3.65° with an "optimal superconducting transition temperature [that] is about 200 mK in both cases" [123], at 5.0° with a "maximum critical temperature of 426 mK" [124], and "evolves smoothly with twist angle" in between [125]. Rhombohedral tetralayer and pentalayer graphene, "without moiré superlattice effects", shows chiral superconductivity with "Tc up to 300 mK" at densities down to $2.4 \times 10^{11}\,\mathrm{cm}^{-2}$ [126]; aligned to hBN, it hosts a quantised anomalous Hall state, a superconductor and a fractional Chern insulator at zero field [127]. Fractional quantum anomalous Hall states appear in twisted $MoTe_2$ without a magnetic field [128], with "nearly 20 hidden states at fractional fillings" at 3.1°, 3.7° and 5.5° [129]; a Perspective compares the two platforms [130], and theory predicts a non-Abelian fractional Chern insulator in the half-filled second miniband [131]. A kagome metal under pressure is a non-moiré flat-band superconductor [132]. The quantum-geometric view is collected in a review of superfluidity [133], a Perspective on geometry in materials [134] and a review in National Science Review whose authors were not read for this map [135]. Twisted bilayer systems as a platform are the subject of a 2026 review, authors not read for this map [136].

What the cold-atom platform offers this research

The atoms see no twist-angle disorder, strain or Coulomb interaction, Ω and V_0 can be tuned in one run, and the superfluid fraction is measured directly, from sound and Leggett bounds [10, 12], Bragg phonons [11] or Josephson oscillations [13]. The platform tests the bosonic side, where geometric superfluidity is predicted to be harder to reach [7].

Key points

- One experimental group, several theory groups; the f_s measurements come from three groups outside TBOL.
- Open QMC, tensor-network and GPE codes exist; no TBOL data set or TBOL-ready GPE code was found.
- Funding reaches quantum simulation broadly (PASQuanS2, quantA, NSF QLCI); twistronics has an ERC grant and the 2026 Kavli Prize.
- Electronic flat-band superconductivity supplies the questions: stiffness, geometry share, angle dependence.

9 Open questions

Twenty-two questions in five themes, from effective models of the pockets to finite temperature and geometry, followed by directions that no study checked for this map has taken up.

The questions follow from Chapter 6, and Figure 9.1 groups them. "Not found" refers to the literature checked for this map (cutoff 21 September 2026).

models	**Q1** pocket Wannier functions, J_M, U_M	**Q2** scalar potential at finite θ	**Q3** detuning with interactions	**Q4** incommensurate angles	**Q5** the experiment's $m = 22$
f_s and sound	**Q6** f_s tensor by phase twist	**Q7** Leggett brackets vs metric	**Q8** sound anisotropy, U scaling	**Q9** depletion in pockets	**Q10** relative-phase branch
experiment	**Q11** SF-II: Josephson array or Bose glass	**Q12** the Ω-shifted transition	**Q13** visibility threshold vs V_0	**Q14** trap, N, scattering lengths	
finite T, dynamics	**Q15** J_M vs k_BT; BKT of pockets	**Q16** Ω ramps, defect counts	**Q17** TWA at low filling	**Q18** extended-junction relaxation	
geometry	**Q19** condensate metric of Rabi bands	**Q20** a genuine geometric window	**Q21** bosonic fractional Chern states	**Q22** fermions, dipoles, photonics	

Figure 9.1: The 22 open questions, one row per theme; the rows follow the subsections below.

9.1 Single-particle theory and effective models

Q1 **Pocket Wannier functions.** What are the Wannier functions of the AA pockets, and how do the inter-pocket tunnelling $J_M(\Omega)$ and on-pocket interaction $U_M(\Omega)$ depend on the coupling? Fitted tight-binding models exist for single-component twisted potentials [28], and Wannier tooling exists [111], but not for the Rabi-coupled bands.

Q2 **The scalar potential at finite angle.** How accurate is the scalar moiré potential, derived for $\theta \cdot l_{ho}^{-1} \ll 1$ in three dimensions [6], for the 2-D square bilayer? The Gaussian-overlap interlayer coupling [18, 21] underestimates the bandwidth by 30 % at $\Omega/J \approx 10$ and tenfold at $\Omega/J \approx 32$ [computed for this map].

Q3 **Detuning with interactions.** For single particles the detuned model maps onto "Hofstadter's original model plus a long-range hopping term" [16]. How do interactions act

there? In quasicrystal limits, Gutzwiller studies find interaction-induced reentrance of the Bose glass [26].

Q4 **Incommensurate angles.** Incommensurate twists act as effective disorder with multifractal states [22], and continuous rotation gives ring-shaped localised states [29]. Beyond Gutzwiller, no interacting study of the Rabi-coupled continuum was found.

Q5 **The experiment's angle.** We computed bands for $m \leq 12$. At $V_0 = 4\,E_R$, $\Omega = 1\,E_R$ the ratio $W_1/W_1(0)$ drops from 0.93 at $m = 8$ to 0.59 at $m = 12$ [computed for this map]; the experiment's $m = 22$ was not computed.

9.2 Superfluid fraction and excitations

Q6 **The f_s tensor.** What is the superfluid-fraction tensor [40] of the two-component condensate under a phase twist [41] across $\Omega/J \approx 5$–25? It has been measured in full for a triangular lattice [12], not for a moiré lattice.

Q7 **Leggett brackets and the metric.** No paper linking the Leggett bounds [8, 9] to the band quantum metric for bosons was found, as the project documents state and the checks did not contradict. Angular dependence in composite potentials is a first step [62].

Q8 **Sound.** Does the Bogoliubov–de Gennes sound anisotropy $f_{s,x} = c_x^2/c_y^2$ [10] follow the Bloch-band mass [42], or does it turn linear in U as in flat bands [14, 59]?

Q9 **Depletion.** In a geometric flat band the depletion stays finite as $U \to 0$ [14]. How large is it in the pockets, and do the Bogoliubov modes carry topology, as in some flat-band condensates [137]?

Q10 **The relative-phase branch.** Rabi coupling turns the demixing transition from first to second order [39], and a moiré-modulated density makes the balance of coupling and interaction vary in space. Its effect on ferromagnetic order [95] and bubble nucleation [96] in a twisted lattice is open.

9.3 The experiment's phases

Q11 **The nature of SF-II.** Is SF-II [1] a Josephson array of AA pockets or a Bose glass? Lattice models find "mobility islands" [21], induced Mott phases [25] and, for dipoles, insulating islands in a superfluid [36].

Q12 **The shifted transition.** Can a continuum calculation reproduce the Ω-dependent superfluid–insulator line? The single-band Bose–Hubbard model "fails to describe but the first Mott lobe" at 15 E_R in hexagonal potentials [44], while cluster Gutzwiller reaches QMC accuracy for square lattices [66, 67].

Q13 **Threshold and depth.** Does the 0.5 E_R visibility threshold [1] track Ω/J between 8 and 14 E_R, as the pocket picture predicts (Chapter 6)?

Q14 **Trap, atom number and scattering lengths.** As read for this map, the paper states neither the radial trap frequency nor the pancake atom number nor the temperature [1]; one GPE study assumes a $2\pi \times 20$ Hz radial trap [31]. Scattering lengths are known for other hyperfine pairs [45, 46], but none for $|1,1\rangle/|2,0\rangle$ was found.

9.4 Finite temperature and dynamics

Q15 **Tunnelling against temperature.** Is J_{M} above $k_B T$? In single-component twisted potentials $J_{\max}$ spans 10^{-11}–$10^{-3}\,E_{\mathrm{R}}$ and temperature depletes the superfluid domains [27]. The Berezinskii–Kosterlitz–Thouless (BKT) transition of the pocket array would follow from Nelson–Kosterlitz [138] and the 2-D critical density [139]; the spin-1 stochastic GPE study [112] is a template.

Q16 **Ramps of Ω.** Ω drives a crossover, not a transition, so a ramp will not give clean Kibble–Zurek exponents. Lattice ramps already deviate [74, 75], and iPEPS needs large bond dimension [76]. Depth quenches produce moiré-periodic vortex pairs [30].

Q17 **Truncated Wigner at low filling.** Truncated Wigner needs fewer modes than particles [70, 72] and highly occupied modes [71]; SU(3) and cluster variants show "noticeable deviations from the experimental results even at relatively short times" in a strongly interacting 2-D quench [73]. Does a sparsely filled pocket array break it?

Q18 **The extended junction.** Two tunnel-coupled 1-D gases realise the sine-Gordon model [158], a Pokrovsky–Talapov transition [159] and relaxation to phase locking [149], with learned representations from snapshots [160]. The 2-D twisted analogue is open.

9.5 Geometry and beyond

Q19 **The condensate metric.** What is $\mathcal{M}^D(\mathbf{k}_c)$ [7] of the Rabi-coupled bands, with the orbital-independent metric [56], and does the U(2) result [58] transfer to non-uniform pockets? No calculation was found.

Q20 **A geometric window.** Band touchings [15], triangle frameworks [57], honeycomb magic couplings [23, 24] and Raman modulation (proposed for fermions) [35] are candidates for a regime where geometry carries D_{s}. Which is reachable with bosons? Kagome and Lieb lattices for bosons have been realised [140–142].

Q21 **Bosonic fractional Chern states.** Interactions break time reversal and give a Chern band for fermions in a TBOL [34]; a bosonic analogue was not found.

Q22 **Transfer.** Which results carry over to fermions [23], dipolar bilayers [36] and photonic moiré lattices [32, 33], where polaritons condense in a moiré flat band [91]?

9.6 Directions not yet systematically addressed

As far as the checks for this map could establish, no study has taken up the following for the Rabi-coupled twisted Bose gas.

D1 Bogoliubov spectra of the extended condensate; the abstracts of the gap-soliton studies report none [78–81].

D2 Superfluid fraction with Leggett brackets, and finite temperature or truncated Wigner, for the two-component Ω-coupled model.

D3 Ramps or quenches of Ω; quenches exist for U, depth and $g(t)$ [26, 30, 31].

D4 An independent trapped simulation compared with the published data [1].

D5 A single account joining the continuum potential [6], the Hofstadter limit [16] and the cluster picture [21].

D6 New f_s probes on moiré lattices: Josephson oscillations [13], sound in supersolids [63] and vortex necklaces [61].

D7 Neural-network surrogates for the TBOL GPE: physics-informed networks [143], deep ground-state solvers [144] and GPE applications [145, 146], with constrained variants from micromagnetics [156, 157].

Key points

- The effective model of the pockets (J_M, U_M) is the first missing piece.
- The f_s tensor, sound and depletion of the Rabi-coupled condensate have not been computed.
- The experiment's missing parameters (trap, temperature, scattering lengths) limit any comparison.
- Geometry may still matter at band touchings, under detuning and in honeycomb bilayers.

10 Reading paths and outlook

Where to start reading, depending on the field one comes from, and what the next experiment or the first superfluid-fraction calculation would change.

10.1 Where to start

A reader from Bose-gas theory should start with the experiment [1] and the reason its band has no magic angle [5, 6]. Then read the flat-band condensate of Julku, Bruun and Törmä [14], which shows what a geometric superfluid looks like (sound linear in U), and the preprint of Huhtinen *et al.* [7], which states when it cannot exist. The magic-angle paper of Bistritzer and MacDonald [2] supplies the vocabulary; chapter 3 is the short version.

For a condensed-matter moiré physicist the new ingredients are the superfluid fraction of a modulated condensate and how cold-atom experiments measure it. Start with Leggett's bound [8] and its two-dimensional brackets [9], then the sound-anisotropy measurement [10]. Read the experiment [1] next, and then the lattice model [21] and the continuous-space Monte Carlo study whose thin superfluid domains are depleted at accessible temperatures [27].

For numerical analysts and applied mathematicians the model is Eq. (1) on a commensurate supercell (chapter 2). For effective models with controlled errors, start with the derivation of the Bistritzer–MacDonald model as the effective dynamics of wave packets "with rigorously estimated error bounds" [147] and with the scalar moiré potential and its validity condition [6]. The detuned limit maps onto a Hofstadter model with a fractal spectrum [16]. Continuum and lattice descriptions are compared in [44], and an open worm-algorithm code for lattice quantum Monte Carlo is described in [64]. The numerical building blocks closest to home are the Vienna time-splitting and absorbing-layer schemes [152, 153] and the fast non-local kernels [154].

10.2 Outlook

The field rests on one experiment at one angle. A second experiment would matter most if it changed the lattice depth along the coupling path: if the moiré-visibility threshold then moves with Ω/J as the band-structure numbers predict, pocket formation is the mechanism; if it stays at a fixed coupling, interactions or temperature set it. A measured temperature and

a sound-anisotropy or Josephson measurement of f_s, as already done in simpler modulated gases [10–13], would turn the phase labels SF and SF-II into numbers.

On the theory side, no superfluid-fraction calculation for the Rabi-coupled condensate was found in the literature checked for this map. A first one, a phase twist of Eq. (1) on the $m = 8$ supercell with Leggett brackets across $\Omega/J \approx 5$–25, would show which contribution carries f_s in the crossover: band mass, density inhomogeneity or tunnelling between pockets. It would also test the working hypothesis of chapter 6, that SF-II is a Josephson array of AA pockets; the experiment's $m = 22$, a trap and finite temperature would then meet the data.

Key points

- Start from the experiment, then flat-band condensates, Leggett's bound or the effective models, depending on background.
- Next experiment: vary the depth along the coupling path; measure f_s and the temperature.
- Next calculation: f_s of the Rabi-coupled condensate across the crossover, with Leggett brackets.

References

[1] Zengming Meng, Liangwei Wang, Wei Han, Fangde Liu, Kai Wen, Chao Gao, Pengjun Wang, Cheng Chin, and Jing Zhang. Atomic Bose–Einstein condensate in twisted-bilayer optical lattices. *Nature*, 615:231–236, 2023. doi:10.1038/s41586-023-05695-4. arXiv:2110.00149.

[2] R. Bistritzer and A. H. MacDonald. Moiré bands in twisted double-layer graphene. *Proc. Natl. Acad. Sci. USA*, 108:12233, 2011. doi:10.1073/pnas.1108174108. arXiv:1009.4203; [Partly verified: PNAS reference from a search listing; arXiv submission date not checked.].

[3] Yuan Cao, Valla Fatemi, Shiang Fang, Kenji Watanabe, Takashi Taniguchi, Efthimios Kaxiras, and Pablo Jarillo-Herrero. Unconventional superconductivity in magic-angle graphene superlattices. *Nature*, 556:43–50, 2018. doi:10.1038/nature26160. arXiv:1803.02342 (ID not seen on a page).

[4] P. Törmä, S. Peotta, and B. A. Bernevig. Superfluidity and quantum geometry in twisted multilayer systems. *Nat. Rev. Phys.*, 4:528, 2022. arXiv:2111.00807; [Partly verified: journal volume and page from the docs, not seen on a primary record.].

[5] Toshikaze Kariyado and Ashvin Vishwanath. Flat band in twisted bilayer Bravais lattices. *Phys. Rev. Research*, 1:033076, 2019. doi:10.1103/PhysRevResearch.1.033076. arXiv:1905.02206; [Partly verified: journal reference from a search listing only; the docs' quotation was not found in the text.].

[6] Ce Wang, Chao Gao, and Zhe-Yu Shi. Quasiperiodic moiré patterns and dimensional localization in three-dimensional quasi-moiré crystals. Preprint, arXiv:2504.02574, 2025. [Partly verified: the validity condition $\Omega \ll$ gap used in the docs is not stated in the paper (the paper states $\theta/l_{\mathrm{ho}} \ll 1$).].

[7] Kukka-Emilia Huhtinen, Matteo Dürrnagel, Valerio Peri, and Sebastian D. Huber. Interplay of local and global quantum geometry in the stability of flat-band superfluids. Preprint, arXiv:2603.08791, 2026.

[8] A. J. Leggett. Can a solid be "superfluid"? *Phys. Rev. Lett.*, 25:1543, 1970. [Partly verified: from the reference list of Pérez-Cruz et al. (2025); primary page not opened; DOI not seen.].

[9] Daniel Pérez-Cruz, Grigori E. Astrakharchik, and Pietro Massignan. Superfluid fraction of interacting bosonic gases. *Phys. Rev. A*, 111:L011302, 2025. arXiv:2403.08416; [Partly verified: journal reference not confirmed on a primary record.].

[10] G. Chauveau, C. Maury, F. Rabec, C. Heintze, G. Brochier, S. Nascimbene, J. Dalibard, J. Beugnon, S. M. Roccuzzo, and S. Stringari. Superfluid fraction in an interacting spatially modulated Bose–Einstein condensate. *Phys. Rev. Lett.*, 130:226003, 2023. arXiv:2302.01776; [Partly verified: journal reference not confirmed on a primary record.].

[11] J. Tao, M. Zhao, and I. B. Spielman. Observation of anisotropic superfluid density in an artificial crystal. *Phys. Rev. Lett.*, 131:163401, 2023. arXiv:2301.01258; [Partly verified: journal reference not confirmed on a primary record.].

[12] F. Rabec, G. Brochier, S. Wattellier, G. Chauveau, Y. Li, S. Nascimbene, J. Dalibard, and J. Beugnon. Superfluid fraction of a 2D Bose–Einstein condensate in a triangular lattice. *Phys. Rev. Lett.*, 136:133401, 2026. doi:10.1103/g494-rj5k. arXiv:2511.04575; [Partly verified: journal reference and DOI from the published PDF (phase 3), not re-checked; $f_s^{(yy)} = 0.82(2)$ not found in the arXiv version.].

[13] G. Biagioni, N. Antolini, B. Donelli, L. Pezzè, A. Smerzi, M. Fattori, A. Fioretti, C. Gabbanini, M. Inguscio, L. Tanzi, and G. Modugno. Measurement of the superfluid fraction of a supersolid by Josephson effect. *Nature*, 629:773–777, 2024. doi:10.1038/s41586-024-07361-9. [Partly verified: "Josephson plasma frequency" not quoted verbatim on the page.].

[14] A. Julku, G. M. Bruun, and P. Törmä. Quantum geometry and flat band Bose–Einstein condensation. *Phys. Rev. Lett.*, 127:170404, 2021. doi:10.1103/PhysRevLett.127.170404. arXiv:2104.14257; [Partly verified: journal reference not read on a primary record.].

[15] K.-E. Huhtinen, J. Herzog-Arbeitman, A. Chew, B. A. Bernevig, and P. Törmä. Revisiting flat band superconductivity: dependence on minimal quantum metric and band touchings. *Phys. Rev. B*, 106:014518, 2022. arXiv:2203.11133; [Partly verified: journal reference not seen on a primary record.].

[16] Xu-Tao Wan, Chao Gao, and Zhe-Yu Shi. Fractal spectrum in twisted bilayer optical lattice. *Phys. Rev. Lett.*, 2026. doi:10.1103/nz52-ypsw. arXiv:2404.08211; PubMed 41651470; [Partly verified: volume, article number and year not confirmed on a primary record; DOI from a search listing.].

[17] Eva Y. Andrei and Allan H. MacDonald. Graphene bilayers with a twist. *Nat. Mater.*, 19: 1265–1275, 2020. doi:10.1038/s41563-020-00840-0. Review Article; arXiv:2008.08129 (ID not seen on a page); Author Correction doi:10.1038/s41563-020-00917-w.

[18] A. González-Tudela and J. I. Cirac. Cold atoms in twisted bilayer optical potentials. *Phys. Rev. A*, 100: 053604, 2019. doi:10.1103/PhysRevA.100.053604. arXiv:1907.06126; [Partly verified: title as on arXiv; the published title form and the journal reference not seen.].

[19] Kai Wen, Zengming Meng, Liangwei Wang, Liangchao Chen, Lianghui Huang, Pengjun Wang, and Jing Zhang. Experimental study of tune-out wavelengths for spin-dependent optical lattice in ^{87}Rb Bose–Einstein condensation. Preprint, arXiv:2109.05670, 2021.

[20] Yunda Li, Wei Han, Zengming Meng, Wenxin Yang, Cheng Chin, and Jing Zhang. Observation of quantized vortex in atomic Bose–Einstein condensate at Dirac point with emergent spin–orbit coupling. *Nat. Photon.*, 19:1264–1269, 2025. doi:10.1038/s41566-025-01763-5. arXiv:2411.16287; [Partly verified: arXiv ID from a search listing only.].

[21] Ganesh C. Paul, Patrik Recher, and Luis Santos. Interacting Bose gases in twisted-bilayer optical lattices. *Phys. Rev. A*, 2025. doi:10.1103/3z9t-xbph. arXiv:2507.12153; [Partly verified: volume and article number not found; DOI from a search listing only.].

[22] Ganesh C. Paul, Patrik Recher, and Luis Santos. Particle dynamics and ergodicity-breaking in twisted-bilayer optical lattices. *Phys. Rev. A*, 108: 053305, 2023. doi:10.1103/PhysRevA.108.053305. arXiv:2306.01588; [Partly verified: journal reference not seen on a primary record.].

[23] Xi-Wang Luo and Chuanwei Zhang. Spin-twisted optical lattices: Tunable flat bands and Larkin–Ovchinnikov superfluids. *Phys. Rev. Lett.*, 126: 103201, 2021. arXiv:2008.01351; [Partly verified: journal reference not seen on a primary record.].

[24] Wenjie Sui, Wei Han, Zheng Vitto Han, Zengming Meng, and Jing Zhang. Topologically nontrivial and trivial flat bands via weak and strong interlayer coupling in twisted bilayer honeycomb optical lattices for ultracold atoms. *Phys. Rev. A*, 111:063306, 2025. arXiv:2506.11520; [Partly verified: journal reference not seen on a primary record; the critical-coupling range 0.09–0.15 E_r not found (the paper gives 0.11 E_r at $\theta \approx 4.4^\circ$).].

[25] Jian-Hua Zeng, Qizhong Zhu, and Liang He. Interaction-induced moiré systems in twisted bilayer optical lattices. *Phys. Rev. A*, 111:063317, 2025. doi:10.1103/zpqd-ryjm. arXiv:2405.20732; the arXiv versions are titled "Interaction induced moiré systems in twisted bilayer optical lattices" (v4) and "Dynamical Moiré Systems in Twisted Bilayer Optical Lattices" (ar5iv); [Partly verified: published title, volume, article number and DOI not confirmed on a primary record.]; [Partly verified: PRA volume and DOI not confirmed.].

[26] Shi-Hao Ding, Li-Jun Lang, Qizhong Zhu, and Liang He. Interaction-induced reentrance of Bose glass and quench dynamics of Bose gases in twisted bilayer and quasicrystal optical lattices. *Phys. Rev. A*, 112:033322, 2025. doi:10.1103/fvny-58kf. arXiv:2503.03375; [Partly verified: quasicrystal limit and quench protocol not in the abstract.].

[27] Dean Johnstone, Shanya Mishra, Zhaoxuan Zhu, Hepeng Yao, and Laurent Sanchez-Palencia. Weak superfluidity in twisted optical potentials. *Phys. Rev. Research*, 6:L042066, 2024. arXiv:2406.12963; [Partly verified: journal reference not seen on a primary record.].

[28] Dean Johnstone, Shanya Mishra, Zhaoxuan Zhu, and Laurent Sanchez-Palencia. Effective tight-binding models in optical moiré potentials. *Phys. Rev. A*, 111:043305, 2025. doi:10.1103/PhysRevA.111.043305. arXiv:2410.21371.

[29] James Walkling, Antonio Štrkalj, and F. Nur Ünal. Localization and topological signatures under periodic twisting. *Phys. Rev. Research*, 8:023356, 2026. doi:10.1103/wt1y-69rf. arXiv:2509.18248; [Partly verified: journal reference not seen on a primary record.].

[30] Tian-Tian Li, Ze-Hong Guo, Xiao-Ning Wang, and Qizhong Zhu. Ground state phases and topological excitations of spin-1 Bose–Einstein condensate in twisted optical lattices. Preprint, arXiv:2412.14731, 2024. Trilayer optical lattice; [Partly verified: journal version not searched.].

[31] Zhenze Fan, Meiling Wang, Juan Wang, and Yan Li. Floquet-engineered moiré quasicrystal patterns of ultracold atoms in twisted bilayer optical lattices. Preprint, arXiv:2508.21093, 2025. Later arXiv versions (v4/v5) are titled "…of ultracold Bose gases in twisted bilayer optical lattices"; [Partly verified: title differs between arXiv versions; cited version not fixed.].

[32] P. Wang, Y. Zheng, X. Chen, C. Huang, Y. V. Kartashov, L. Torner, V. V. Konotop, and F. Ye. Localization and delocalization of light in photonic moiré lattices. *Nature*, 577:42–46, 2020. doi:10.1038/s41586-019-1851-6. [Partly verified: the docs' "same mathematics (NLSE)" framing not checked.].

[33] Q. Fu, P. Wang, C. Huang, Y. V. Kartashov, L. Torner, V. V. Konotop, and F. Ye. Optical soliton formation controlled by angle twisting in photonic moiré lattices. *Nat. Photon.*, 14:663–668, 2020. doi:10.1038/s41566-020-0679-9. [Partly verified: "threshold vanishes at the localisation–delocalisation transition" not verbatim on the page.].

[34] Yan-Bin Yang, Jiong-Hao Wang, Shicheng Ma, and Yong Xu. Fractional Chern insulators in twisted bilayer optical lattices. Preprint, arXiv:2609.09733, 2026. v1 9 September 2026, v2 16 September 2026; [Partly verified: model details (fermions, honeycomb, $w \approx 0.05$, 4×6 cells, $\nu = 1/3$) not in the abstract.].

[35] T. Salamon, A. Celi, R. W. Chhajlany, I. Frérot, M. Lewenstein, L. Tarruell, and D. Rakshit. Simulating twistronics without a twist. *Phys. Rev. Lett.*, 125:030504, 2020. arXiv:1912.12736; [Partly verified: journal reference not seen on a primary record.].

[36] Chao Zhang, Zhijie Fan, Barbara Capogrosso-Sansone, and Youjin Deng. Dipolar bosons in a twisted bilayer geometry. *Phys. Rev. B*, 111: 024511, 2025. doi:10.1103/PhysRevB.111.024511. arXiv:2405.16425; [Partly verified: journal reference from a search listing only.].

[37] Enhong Cheng, Yu Tan, Yanzhen Xu, and Li-Jun Lang. Fate of moiré flat bands for a weakly repulsive Bose–Einstein condensate in one-dimensional $\mathcal{PT}$-symmetric bichromatic optical lattices. *Phys. Rev. A*, 114:033310, 2026. doi:10.1103/ppvl-5lmg. arXiv:2608.01680.

[38] Siyu Xie, Qiang Xu, Qianqian Shi, and Wanzhou Zhang. Local supersolid in moiré modulated Bose–Hubbard model using density-matrix renormalization group method. Preprint, arXiv:2605.10238, 2026.

[39] Marta Abad and Alessio Recati. A study of coherently coupled two-component Bose–Einstein condensates. *Eur. Phys. J. D*, 67:148, 2013. doi:10.1140/epjd/e2013-40053-2. arXiv:1301.6864.

[40] P. B. Blakie. Superfluid fraction tensor of a two-dimensional supersolid. *J. Phys. B*, 2024. doi:10.1088/1361-6455/ad41c1. arXiv:2308.14001; [Partly verified: volume, article number and DOI not confirmed on a primary record.].

[41] A. J. Leggett. On the superfluid fraction of an arbitrary many-body system at $T = 0$. *J. Stat. Phys.*, 93: 927, 1998. doi:10.1023/B:JOSS.0000033170.38619.6c. [Partly verified: from a citing paper's reference list and a publisher listing; DOI read from the listing URL.].

[42] E. Taylor and E. Zaremba. Bogoliubov sound speed in periodically modulated Bose–Einstein condensates. *Phys. Rev. A*, 68:053611, 2003. doi:10.1103/PhysRevA.68.053611. arXiv:cond-mat/0308194; [Partly verified: journal reference not seen on a primary record.].

[43] W. Zwerger. Mott–Hubbard transition of cold atoms in optical lattices. *J. Opt. B: Quantum Semiclass. Opt.*, 5:S9, 2003. doi:10.1088/1464-4266/5/2/352. arXiv:cond-mat/0211314; [Partly verified: from a search listing only; the asymptotic hopping formula not checked in the text.].

[44] Danilo Nascimento Guimaraes and Laurent Sanchez-Palencia. Quantum phase diagrams for bosons in hexagonal optical potentials: A continuous-space quantum Monte Carlo study. *Phys. Rev. A*, 114:023312, 2026. arXiv:2605.06171; [Partly verified: journal reference not seen on a primary record.].

[45] M. Egorov, B. Opanchuk, P. Drummond, B. V. Hall, P. Hannaford, and A. I. Sidorov. Measurement of s-wave scattering lengths in a two-component Bose–Einstein condensate. *Phys. Rev. A*, 87:053614, 2013. arXiv:1204.1591; [Partly verified: journal reference not seen on a primary record; $a_{11} = 100.40\, a_0$ not in the paper text seen.].

[46] K. M. Mertes et al. Nonequilibrium dynamics and superfluid ring excitations in binary Bose–Einstein condensates. *Phys. Rev. Lett.*, 99:190402, 2007. doi:10.1103/PhysRevLett.99.190402. [Partly verified: from a search listing only; author list not confirmed.].

[47] Yuan Cao, Valla Fatemi, Ahmet Demir, Shiang Fang, Spencer L. Tomarken, Jason Y. Luo, Javier D. Sanchez-Yamagishi, Kenji Watanabe, Takashi Taniguchi, Efthimios Kaxiras, Ray C. Ashoori, and Pablo Jarillo-Herrero. Correlated insulator behaviour at half-filling in magic-angle graphene superlattices. *Nature*, 556:80–84, 2018. doi:10.1038/nature26154. arXiv:1802.00553 (ID not seen on a page).

[48] S. Peotta and P. Törmä. Superfluidity in topologically nontrivial flat bands. *Nat. Commun.*, 6:8944, 2015. doi:10.1038/ncomms9944. arXiv:1506.02815; [Partly verified: the docs' "$D_s \propto \Delta \int g$" form and the quotation attributed to it (ledger F127) not found in the abstract.].

[49] Fang Xie, Zhida Song, Biao Lian, and B. Andrei Bernevig. Topology-bounded superfluid weight in twisted bilayer graphene. *Phys. Rev. Lett.*, 124: 167002, 2020. doi:10.1103/PhysRevLett.124.167002. arXiv:1906.02213; [Partly verified: journal reference from a search listing only.].

[50] N. Verma, T. Hazra, and M. Randeria. Optical spectral weight, phase stiffness and t_c bounds for trivial and topological flat band superconductors. *Proc. Natl. Acad. Sci. USA*, 118:e2106744118, 2021. arXiv:2103.08540; [Partly verified: PNAS volume and article number from the research brief, not read.].

[51] M. Tanaka, J. Î-j. Wang, T. H. Dinh, D. Rodan-Legrain, S. Zaman, M. Hays, A. Almanakly, B. Kannan, D. K. Kim, B. M. Niedzielski, K. Serniak, M. E. Schwartz, K. Watanabe, T. Taniguchi, T. P. Orlando, S. Gustavsson, J. A. Grover, P. Jarillo-Herrero, and W. D. Oliver. Superfluid stiffness of magic-angle twisted bilayer graphene. *Nature*, 638:99–105, 2025. doi:10.1038/s41586-024-08494-7. [Partly verified: the quotation attributed to it in the docs (ledger F127) not found in the abstract.].

[52] M. Banerjee et al. Superfluid stiffness of twisted trilayer graphene superconductors. *Nature*, 638:93–98, 2025. doi:10.1038/s41586-024-08444-3. [Partly verified: bibliographic data from the reference list of the News & Views page; author list not confirmed.].

[53] Paritosh Karnatak and Christian Schönenberger. 'Unconventional' superconductivity probed in twisted graphene. *Nature*, 638:44–45, 2025. doi:10.1038/d41586-025-00057-8. News & Views, 5 February 2025.

[54] Jian Zhou. Band-basis decomposition of superfluid weight in magic-angle twisted bilayer graphene: Quantifying geometric and conventional contributions. Preprint, arXiv:2604.05994, 2026. [Partly verified: submission date not reread.].

[55] A. Julku, G. M. Bruun, and P. Törmä. Excitations of a Bose–Einstein condensate and the quantum geometry of a flat band. *Phys. Rev. B*, 104:144507, 2021. doi:10.1103/PhysRevB.104.144507. Erratum: Phys. Rev. B 107, 019901 (2023); [Partly verified: bibliographic data from search listings only; arXiv ID not found (it is not 2104.14257, the companion PRL).].

[56] A. Julku, G. Salerno, and P. Törmä. Supefluidity of flat band Bose–Einstein condensates revisited. *Low Temp. Phys.*, 49:701, 2023. arXiv:2210.11906; title spelled "Supefluidity" as on arXiv; [Partly verified: journal reference and published title form not seen.].

[57] K.-E. Huhtinen et al. Stability of flat-band Bose–Einstein condensation from the geometry of compact localized states. Preprint, arXiv:2603.09954, 2026. [Partly verified: author list not confirmed (only the first author seen).].

[58] Yi-Cai Zhang. Double commutator method for a two band Bose–Einstein condensate: superfluid density of a flat band superfluid. Preprint, arXiv:2503.23656, 2025. [Partly verified: the docs' "two-component continuum flat band" description is not the paper's (two-band BEC).].

[59] M. Burgher, S. Loddo, L. Vanderstraeten, N. Goldman, and I. Amelio. Competition between pair and single-particle superfluidity in bosonic quasi-flat bands: A Gaussian state approach. Preprint, arXiv:2605.26231, 2026. [Partly verified: "all-flat ladders" not stated in the text seen.].

[60] L. Pizzino, H. Pan, T. Giamarchi, and H. Yao. Leggett's bound and superfluidity in strongly interacting bosons. Preprint, arXiv:2512.14346, 2025. [Partly verified: title from a search listing only; authors not seen.].

[61] Probing superfluidity with quantum vortex necklaces. Preprint, arXiv:2505.16728, 2025. Published (per a search listing) as "Probing superfluidity with quantum vortex necklaces and Leggett's bounds", Phys. Rev. Research, doi:10.1103/r5x2-yy6v; [Partly verified: from search listings only; authors not seen.].

[62] Pérez-Cruz et al. Superfluid response of bosonic fluids in composite optical potentials: angular dependence and Leggett's bounds. Preprint, arXiv:2603.29603, 2026. [Partly verified: title from a search listing only; authors not seen.].

[63] Marija Šindik, Tomasz Zawiślak, Alessio Recati, and Sandro Stringari. Sound, superfluidity and layer compressibility in a ring dipolar supersolid. Preprint, arXiv:2308.05981, 2023. [Partly verified: journal version not found.].

[64] N. Sadoune and L. Pollet. Efficient and scalable path integral Monte Carlo simulations with worm-type updates for Bose–Hubbard and XXZ models. *SciPost Phys. Codebases*, page 9, 2022. arXiv:2204.12262; [Partly verified: Codebases number and GPL licence not seen on the paper record (SciPost blocked).].

[65] G. Orso and S. Stringari. Superfluid fraction and Leggett bound in a density modulated strongly interacting Fermi gas at zero temperature. *Phys. Rev. A*, 109:023301, 2024. doi:10.1103/PhysRevA.109.023301. arXiv:2311.16709; a unitary-Fermi-gas paper; [Partly verified: journal reference not seen on the page.].

[66] Dirk-Sören Lühmann. Cluster Gutzwiller method for bosonic lattice systems. *Phys. Rev. A*, 87:043619, 2013. arXiv:1302.6761; [Partly verified: journal reference not seen on a primary record.].

[67] Barbara Capogrosso-Sansone, Şebnem Güneş Söyler, Nikolay Prokof'ev, and Boris Svistunov. Monte Carlo study of two-dimensional Bose–Hubbard model. Preprint, arXiv:0710.2703, 2007. Identified from the unattributed dossier ID x0710.2703; [Partly verified: journal reference not shown on the arXiv record.].

[68] Zakari Denis and Giuseppe Carleo. Accurate neural quantum states for interacting lattice bosons. *Quantum*, 9:1772, 2025. doi:10.22331/q-2025-06-17-1772. arXiv:2404.07869; [Partly verified: the docs' "backflow-Jastrow" description is not the paper's wording.].

[69] Jacob Jordan, Román Orús, and Guifré Vidal. Numerical study of the hard-core Bose–Hubbard model on an infinite square lattice. Preprint, arXiv:0901.0420, 2009. Identified from the unattributed dossier ID x0901.0420; [Partly verified: journal reference not shown on the arXiv record.].

[70] Alice Sinatra, Carlos Lobo, and Yvan Castin. The truncated Wigner method for Bose condensed gases: limits of validity and applications. *J. Phys. B*, 35:3599, 2002. arXiv:cond-mat/0201217; HAL hal-00002253; [Partly verified: venue from the research brief; not confirmed (the arXiv text names an earlier J. Mod. Opt. paper).].

[71] P. B. Blakie, A. S. Bradley, M. J. Davis, R. J. Ballagh, and C. W. Gardiner. Dynamics and statistical mechanics of ultra-cold Bose gases using c-field techniques. *Adv. Phys.*, 57(5):363, 2008. arXiv:0809.1487; [Partly verified: first page from the docs, not seen on a primary record.].

[72] A. Polkovnikov. Phase space representation of quantum dynamics. *Ann. Phys.*, 325:1790, 2010. arXiv:0905.3384; merged with the unattributed dossier ID x0905.3384; [Partly verified: bibliographic data from a search listing only.].

[73] Kazuma Nagao and Seiji Yunoki. Two-dimensional correlation propagation dynamics with a cluster discrete phase-space method. Preprint, arXiv:2404.18594, 2024. [Partly verified: journal reference not seen; the "t above about $0.1\,\hbar/J_{\mathrm{fin}}$" breakdown sentence not in the excerpt read.].

[74] David Chen, Matthew White, Cecilia Borries, and Brian DeMarco. Quantum quench of an atomic Mott insulator. *Phys. Rev. Lett.*, 106:235304, 2011. arXiv:1103.4662; [Partly verified: journal reference not seen on a primary record.].

[75] S. Braun, M. Friesdorf, S. S. Hodgman, M. Schreiber, J. P. Ronzheimer, A. Riera, M. del Rey, I. Bloch, J. Eisert, and U. Schneider. Emergence of coherence and the dynamics of quantum phase transitions. *Proc. Natl. Acad. Sci. USA*, 112:3641, 2015. doi:10.1073/pnas.1408861112. arXiv:1403.7199; [Partly verified: PNAS volume and page not seen on a primary record; the 0.3–0.6 exponent range not confirmed verbatim.].

[76] Jacek Dziarmaga and Jakub M. Mazur. Tensor network simulation of the quantum Kibble–Zurek quench from the Mott to the superfluid phase in the two-dimensional Bose–Hubbard model. *Phys. Rev. B*, 107:144510, 2023. arXiv:2302.13347; [Partly verified: journal reference not seen on a primary record.].

[77] Rui Tian, Yue Zhang, Tianhao Wu, Min Liu, Yong-Chang Zhang, Shuai Li, and Bo Liu. Nonlinearity-induced dynamical self-organized twisted-bilayer lattices in Bose–Einstein condensates. Preprint, arXiv:2407.21466, 2024.

[78] Xiuye Liu and Jianhua Zeng. Matter-wave gap solitons and vortices of dense Bose–Einstein condensates in moiré optical lattices. *Chaos Solitons Fractals*, 174:113869, 2023. doi:10.1016/j.chaos.2023.113869. [Partly verified: bibliographic data from a bibliographic-database record (IDEAS/RePEc), not the publisher page.].

[79] Pingping Fang, Chao Gao, and Ji Lin. Bifurcations and dynamics of nonlinear excitations in twisted-bilayer optical lattices. *Chaos Solitons Fractals*, 195:116314, 2025. doi:10.1016/j.chaos.2025.116314. Merged with the duplicate key anon2025chaos; [Partly verified: bibliographic data from a bibliographic-database record (IDEAS/RePEc).].

[80] Pu Tu, Qing-Qing Wang, Jin-Ping Ma, Kai-Hua Shao, Xi Zhao, Bao-Long Xi, Xiao-Fei Zhang, and Yu-Ren Shi. Vector gap solitons of two-component Bose gas in twisted-bilayer optical lattice. *Chaos Solitons Fractals*, 190, 2025. doi:10.1016/j.chaos.2024.115773. [Partly verified: article number not shown; bibliographic-database record; assignment of the a/b keys inferred.].

[81] Pu Tu, Jin-Ping Ma, Xi Zhao, Bao-Long Xi, Kai-Hua Shao, Xiao-Fei Zhang, and Yu-Ren Shi. Gap solitons of spin–orbit coupled Bose–Einstein condensates with Rabi coupling in twisted-bilayer optical lattices. *Physica A*, 666, 2025. doi:10.1016/j.physa.2025.130504. [Partly verified: article number not shown; bibliographic-database record; assignment of the a/b keys inferred.].

[82] Wang et al. Gap solitons and vortices in two-dimensional spin-orbit-coupled Bose–Einstein condensates loaded onto moiré optical lattices. *Quantum Front.*, 4:9, 2025. doi:10.1007/s44214-025-00082-8. [Partly verified: from a search listing only; full author list not obtained (publisher page refused).].

[83] Jr-Chiun Yu, Shaurya Bhave, Lee Reeve, Bo Song, and Ulrich Schneider. Observing the two-dimensional Bose glass in an optical quasicrystal. *Nature*, 633(8029):338–343, 2024. doi:10.1038/s41586-024-07875-2. [Partly verified: the docs' framing ("the incommensurate end of the same physics") and the Cambridge affiliation not checked.].

[84] LodePollet. LodePollet/worm: Worm algorithm for Bose–Hubbard and XXZ models. https://github.com/LodePollet/worm. GitHub repository; GPL-3.0; 26 commits; no releases; accessed 21 September 2026.

[85] Zakari Denis. cqsl/lattice-bosons: v1.0.0. https://doi.org/10.5281/zenodo.15632176, 2025. Zenodo software record, 10 June 2025; Apache License 2.0; accessed 21 September 2026.

[86] X.-R. Mao, Z.-K. Shao, H.-Y. Luan, S.-L. Wang, and R.-M. Ma. Magic-angle lasers in nanostructured moiré superlattice. *Nat. Nanotechnol.*, 16: 1099–1105, 2021. doi:10.1038/s41565-021-00956-7.

[87] Luojun Du, Maciej R. Molas, Zhiheng Huang, Guangyu Zhang, Feng Wang, and Zhipei Sun. Moiré photonics and optoelectronics. *Science*, 379(6639):eadg0014, 2023. doi:10.1126/science.adg0014. Review.

[88] Mingjie Zhang and Eric Mazur. Dielectric photonic moiré metamaterials. *npj Metamaterials*, 2:20, 2026. doi:10.1038/s44455-026-00029-6. Review.

[89] Nature Materials. Moiré beyond van der Waals. *Nat. Mater.*, 23:1151, 2024. doi:10.1038/s41563-024-01999-6. Editorial (unsigned).

[90] [title to be confirmed]. *Nat. Mater.*, 2024. Review of photonic and polaritonic moiré; [Partly verified: identity not confirmed; candidate: M. Oudich, X. Kong, T. Zhang, C. Qiu, Y. Jing, "Engineered moiré photonic and phononic superlattices", Nat. Mater. 23, 1169–1178 (2024), doi:10.1038/s41563-024-01950-9.].

[91] Exciton polariton condensation in a perovskite moiré flat band at room temperature. *Sci. Adv.*, 11:eadx2361, 2025. doi:10.1126/sciadv.adx2361. Published 8 August 2025; 13 authors (NTU and NUS, Singapore); [Partly verified: author names not obtained; volume from the ADS bibcode.].

[92] Bo Song, Shaurya Bhave, Emmanuel Gottlob, Leanne Reeve, and Ulrich Schneider. Quantum quenches across the Bose-glass transition. Preprint, arXiv:2609.05589, 2026.

[93] H. Yoo, R. Engelke, S. Carr, S. Fang, K. Zhang, P. Cazeaux, S. H. Sung, R. Hovden, A. W. Tsen, T. Taniguchi, K. Watanabe, G.-C. Yi, M. Kim, M. Luskin, E. B. Tadmor, E. Kaxiras, and P. Kim. Atomic and electronic reconstruction at the van der Waals interface in twisted bilayer graphene. *Nat. Mater.*, 18:448–453, 2019. doi:10.1038/s41563-019-0346-z.

[94] Daesung Park, Changwon Park, Kunihiro Yananose, et al. Unconventional domain tessellations in moiré-of-moiré lattices. *Nature*, 641: 896–903, 2025. doi:10.1038/s41586-025-08932-0. Twisted trilayer graphene; [Partly verified: author list not confirmed (21 authors, last H. Yoo).].

[95] R. Cominotti et al. Ferromagnetism in an extended coherently-coupled atomic superfluid. *Phys. Rev. X*, 13:021037, 2023. arXiv:2209.13235; [Partly verified: volume and article number from the docs (a listing shows Phys. Rev. X, June 2023); author list not confirmed.].

[96] A. Zenesini, A. Berti, R. Cominotti, C. Rogora, I. G. Moss, T. P. Billam, I. Carusotto, G. Lamporesi, A. Recati, and G. Ferrari. False vacuum decay via bubble formation in ferromagnetic superfluids. *Nat. Phys.*, 20:558–563, 2024. doi:10.1038/s41567-023-02345-4. arXiv:2305.05225 (arXiv title "Observation of false vacuum decay via bubble formation in ferromagnetic superfluids").

[97] Paula Mellado. Magnetic moiré systems: a review. *J. Phys.: Condens. Matter*, 2025. arXiv:2506.05620; [Partly verified: journal from the docs; not shown on the arXiv record; volume and article number not found.].

[98] Luis Santos group, ITP, Leibniz Universität Hannover. Publikationen – Institut für Theoretische Physik – Leibniz Universität Hannover. https://www.itp.uni-hannover.de/arbeitsgruppen/luis-santos/publikationen. Group publication list; accessed 21 September 2026; [Partly verified: does not support "interaction-induced moiré" (lists one twisted-bilayer paper, PRA 108, 053305).].

[99] S. D. Huber and E. Altman. Bose condensation in flat bands. *Phys. Rev. B*, 2010. arXiv:1007.4640; [Partly verified: volume and article number not found.].

[100] ALPSCore: core ALPS libraries. https://github.com/ALPSCore/ALPSCore, 2026. MIT License; release v2.3.3 of 11 May 2026; accessed 21 September 2026.

[101] NetKet 3.22.4. https://pypi.org/project/NetKet/, 2026. PyPI release of 17 August 2026; Apache 2.0; Python $\geq$ 3.11; accessed 21 September 2026.

[102] TeNPy: Tensor Network Python. https://github.com/tenpy/tenpy, 2025. GitHub repository; latest release v1.1.0 (11 November 2025); Apache-2.0; accessed 21 September 2026.

[103] ITensors.jl. https://github.com/ITensor/ITensors.jl. GitHub repository; Apache-2.0; latest release v0.9.30 (30 May, year not shown); accessed 21 September 2026.

[104] XMDS2 (xpdeint): (stochastic) partial differential equation integrator. https://github.com/GrahamDennis/xpdeint. GPL-2.0; no release tag; documentation (version 2.2.2) https://xmds2.readthedocs.io/en/latest/; accessed 21 September 2026.

[105] R. Duboscq and X. Antoine. GPELab: a free Matlab toolbox to compute and simulate Bose–Einstein condensates. https://gpelab.math.cnrs.fr/. Companion papers in Comput. Phys. Commun. (pii S0010465514002318, S0010465515001113); no licence or version stated; accessed 21 September 2026.

[106] James R. Schloss and Lee J. O'Riordan. GPUE: Graphics Processing Unit Gross–Pitaevskii Equation solver. *J. Open Source Softw.*, 3(32):1037, 2018. doi:10.21105/joss.01037. Software archive doi:10.5281/zenodo.2182729; [Partly verified: the "performance greater than other available suites" quotation not on the landing page.].

[107] L. Fioroni, L. Gravina, J. Stefaniak, A. Baumgärtner, F. Finger, D. Dreon, and T. Donner. A Python GPU-accelerated solver for the Gross–Pitaevskii equation and applications to many-body cavity QED. Preprint, arXiv:2404.14401, 2024. Package TorchGPE.

[108] Sachin Singh Rawat, Shawan Kumar Jha, Mahendra Kumar Verma, and Pankaj Kumar Mishra. quTARANG: A high-performance computing Python package to study turbulence using the Gross–Pitaevskii equation. *Comput. Phys. Commun.*, 315:109725, 2025. doi:10.1016/j.cpc.2025.109725. Code MIT-licensed; earlier preprint arXiv:2301.08275 with a different title.

[109] A. S. Bradley. FourierGPE.jl: simple Gross–Pitaevskii equation solver using FFTW. https://github.com/AshtonSBradley/FourierGPE.jl. MIT License; no longer actively maintained; companion ProjectedGPE.jl; accessed 21 September 2026.

[110] GeneralizedGrossPitaevskii.jl: high-performance Julia package for quantum fluid dynamics simulations. https://github.com/marcsgil/GeneralizedGrossPitaevskii.jl. MIT License; accessed 21 September 2026.

[111] Wannier90: maximally-localised generalised Wannier functions code. https://github.com/wannier-developers/wannier90, 2024. Official repository; LGPL-2.1; latest release v3.1.0 (15 February 2024); code paper: Pizzi et al., J. Phys.: Condens. Matter 32, 165902 (2020); accessed 21 September 2026; [Partly verified: the pywannier alternative mentioned in the inventory not checked.].

[112] Andrew P. C. Underwood and P. B. Blakie. Stochastic Gross–Pitaevskii theory for a spin-1 Bose gas: Application to superfluidity in two dimensions. *Phys. Rev. A*, 112:023315, 2025. doi:10.1103/slxb-df37. arXiv:2502.18792; [Partly verified: journal reference not seen on the arXiv page; numerical parameters (γ, k_{cut}, grid, samples) not in the excerpt read.].

[113] Quantum Flagship. PASQuanS2.1 – Programmable Atomic Large-scale Quantum Simulation 2.1. https://qt.eu/projects/simulation/PASQuans2_1. CORDIS 101113690; project launched April 2023; accessed 21 September 2026.

[114] U.S. National Science Foundation. Eight NSF research institutes to propel U.S. quantum science with $290M investment. https://www.nsf.gov/news/eight-nsf-research-institutes-propel-us-quantum-science-290m, 2026. 25 August 2026; accessed 21 September 2026.

[115] University of Maryland, Department of Physics. UMD leads new $25M NSF Quantum Leap Challenge Institute for Robust Quantum Simulation. https://umdphysics.umd.edu/about-us/news/department-news/1733-q-leap.html, 2021. 2 September 2021; accessed 21 September 2026.

[116] CORDIS. QTWIST – Quantum nanomaterials by twistronics. https://cordis.europa.eu/project/id/101001515. ERC Consolidator Grant 101001515, The University of Manchester, 1 April 2021 – 31 March 2027; accessed 21 September 2026.

[117] The Kavli Prize. 2026 Kavli Prize in Nanoscience. https://www.kavliprize.org/prizes/nanoscience/2026, 2026. Accessed 21 September 2026; also reported by ACerS Ceramic Tech Today, 23 June 2026, https://ceramics.org/ceramic-tech-today/twistronics-kavli-prize-2026/.

[118] planqc GmbH. planqc to build 1,000-qubit neutral atom quantum computer for LRZ. https://www.planqc.eu/news/planqc-to-build-1-000-qubit-neutral-atom-quantum-computer, 2024. News release, 13 November 2024; accessed 21 September 2026.

[119] Jonathan Wurtz, Alexei Bylinskii, Boris Braverman, Jesse Amato-Grill, Sergio H. Cantu, Florian Huber, Alexander Lukin, Fangli Liu, Phillip Weinberg, John Long, Sheng-Tao Wang, Nathan Gemelke, and Alexander Keesling. Aquila: QuEra's 256-qubit neutral-atom quantum computer. Preprint, arXiv:2306.11727, 2023.

[120] C. Gross and I. Bloch. Quantum simulations with ultracold atoms in optical lattices. *Science*, 357: 995, 2017. doi:10.1126/science.aal3837. Erratum: Science 357, eaap9526; [Partly verified: from a search listing only.].

[121] L. Amico et al. Colloquium: Atomtronic circuits: From many-body physics to quantum technologies. *Rev. Mod. Phys.*, 94:041001, 2022. doi:10.1103/RevModPhys.94.041001. [Partly verified: from a search listing only; author list not confirmed.].

[122] H. Tian et al. Evidence for Dirac flat band superconductivity enabled by quantum geometry. *Nature*, 614:440, 2023. doi:10.1038/s41586-022-05576-2. A published comment exists: arXiv:2303.06054; [Partly verified: from a search listing only; author list not confirmed.].

[123] Yiyu Xia, Zhongdong Han, Kenji Watanabe, Takashi Taniguchi, Jie Shan, and Kin Fai Mak. Superconductivity in twisted bilayer WSe_2. *Nature*, 637:833–838, 2025. doi:10.1038/s41586-024-08116-2. Published online 30 October 2024.

[124] Yinjie Guo, Jordan Pack, Joshua Swann, Luke Holtzman, Matthew Cothrine, Kenji Watanabe, Takashi Taniguchi, David G. Mandrus, Katayun Barmak, James Hone, Andrew J. Millis, Abhay Pasupathy, and Cory R. Dean. Superconductivity in 5.0° twisted bilayer WSe_2. *Nature*, 637:839–845, 2025. doi:10.1038/s41586-024-08381-1. Published online 22 January 2025.

[125] Yinjie Guo, John Cenker, Ammon Fischer, Daniel Muñoz-Segovia, Jordan Pack, Luke Holtzman, Lennart Klebl, Kenji Watanabe, Takashi Taniguchi, Katayun Barmak, James Hone, Angel Rubio, Dante M. Kennes, Andrew J. Millis, Abhay Pasupathy, and Cory R. Dean. Angle evolution of the superconducting phase diagram in twisted bilayer WSe_2. *Nature*, 652:622–627, 2026. doi:10.1038/s41586-026-10357-2.

[126] Tonghang Han, Zhengguang Lu, Zach Hadjri, et al. Signatures of chiral superconductivity in rhombohedral graphene. *Nature*, 643:654–661, 2025. doi:10.1038/s41586-025-09169-7. Rhombohedral tetralayer and pentalayer graphene without moiré superlattice effects; [Partly verified: author list not confirmed (23 authors).].

[127] Youngjoon Choi, Ysun Choi, Marco Valentini, Caitlin L. Patterson, Ludwig F. W. Holleis, Owen I. Sheekey, Hari Stoyanov, Xiang Cheng, Takashi Taniguchi, Kenji Watanabe, and Andrea F. Young. Superconductivity and quantized anomalous Hall effect in rhombohedral graphene. *Nature*, 639:342–347, 2025. doi:10.1038/s41586-025-08621-y.

[128] Jiaqi Cai, Eric Anderson, Chong Wang, Xiaowei Zhang, Xiaoyu Liu, William Holtzmann, Yinong Zhang, Fengren Fan, Takashi Taniguchi, Kenji Watanabe, Ying Ran, Ting Cao, Liang Fu, Di Xiao, Wang Yao, and Xiaodong Xu. Signatures of fractional quantum anomalous Hall states in twisted $MoTe_2$. *Nature*, 622:63–68, 2023. doi:10.1038/s41586-023-06289-w.

[129] Yiping Wang, Jeongheon Choe, Eric Anderson, Weijie Li, Julian Ingham, Eric A. Arsenault, Yiliu Li, Xiaodong Hu, Takashi Taniguchi, Kenji Watanabe, Xavier Roy, Dmitri Basov, Di Xiao, Raquel Queiroz, James C. Hone, Xiaodong Xu, and X.-Y. Zhu. Hidden states and dynamics of fractional fillings in twisted $MoTe_2$ bilayers. *Nature*, 641:1149–1155, 2025. doi:10.1038/s41586-025-08954-8. arXiv:2502.21153 (arXiv version titled "Hidden States and Dynamics of Fractional Fillings in t$MoTe_2$ Moiré Superlattices"; ID not seen on a page); Author Correction 29 May 2026, doi:10.1038/s41586-026-10717-y.

[130] Tingxin Li, Jianpeng Liu, Jian Xie, and Xiaobo Lu. Fractional quantum anomalous Hall effect in moiré fractional Chern insulators. *Nat. Mater.*, 2026. doi:10.1038/s41563-026-02649-9. Perspective; published online 3 July 2026, no volume assigned yet.

[131] Feng Chen, Wei-Wei Luo, Wei Zhu, and D. N. Sheng. Robust non-Abelian even-denominator fractional Chern insulator in twisted bilayer $MoTe_2$. *Nat. Commun.*, 16:2115, 2025. doi:10.1038/s41467-025-57326-3.

[132] Siqi Wu, Chenchao Xu, Xiaoqun Wang, Hai-Qing Lin, Chao Cao, and Guang-Han Cao. Flat-band enhanced antiferromagnetic fluctuations and superconductivity in pressurized $CsCr_3Sb_5$. *Nat. Commun.*, 16:1375, 2025. doi:10.1038/s41467-025-56582-7.

[133] Sebastiano Peotta, Kukka-Emilia Huhtinen, and Päivi Törmä. Quantum geometry in superfluidity and superconductivity. Preprint, arXiv:2308.08248, 2023. Review.

[134] Nishchhal Verma, Philip J. W. Moll, Tobias Holder, and Raquel Queiroz. Quantum geometry and the hidden scales in materials. *Nat. Rev. Phys.*, 8:226–239, 2026. doi:10.1038/s42254-026-00923-y. Perspective.

[135] Quantum geometry in condensed matter. *Natl. Sci. Rev.*, 12(3):nwae334, 2025. doi:10.1093/nsr/nwae334. Review; PubMed 39958148; [Partly verified: from search listings only; authors not seen.].

[136] Twisted bilayer systems: a versatile platform for exploring emergent quantum phenomena. *Natl. Sci. Rev.*, 13(4):nwag057, 2026.

doi:10.1093/nsr/nwag057. PubMed 41704745; [Partly verified: from search listings only; authors not seen; DOI read from a listing URL.].

[137] Zahra Jalali-Mola, Tobias Grass, Valentin Kasper, Maciej Lewenstein, and Utso Bhattacharya. Topological Bogoliubov quasiparticles from Bose–Einstein condensate in a flat band system. *Phys. Rev. Lett.*, 131:226601, 2023. doi:10.1103/PhysRevLett.131.226601. arXiv:2302.09910; [Partly verified: journal reference from a search listing, not checked.].

[138] D. R. Nelson and J. M. Kosterlitz. Universal jump in the superfluid density of two-dimensional superfluids. *Phys. Rev. Lett.*, 39:1201, 1977. doi:10.1103/PhysRevLett.39.1201. [Partly verified: from a search listing only.].

[139] Nikolay Prokof'ev, Oliver Ruebenacker, and Boris Svistunov. Critical point of a weakly interacting two-dimensional Bose gas. *Phys. Rev. Lett.*, 87:270402, 2001. arXiv:cond-mat/0106075; [Partly verified: journal reference not seen on a primary record.].

[140] Jo et al. Ultracold atoms in a tunable optical kagome lattice. *Phys. Rev. Lett.*, 108:045305, 2012. doi:10.1103/PhysRevLett.108.045305. arXiv:1109.1591; [Partly verified: from search listings only; author list not confirmed.].

[141] Taie et al. Coherent driving and freezing of bosonic matter wave in an optical Lieb lattice. *Sci. Adv.*, 1:e1500854, 2015. doi:10.1126/sciadv.1500854. arXiv:1506.00587; [Partly verified: from search listings only; author list not confirmed.].

[142] Tsz-Him Leung, Malte N. Schwarz, Shao-Wen Chang, Charles D. Brown, Govind Unnikrishnan, and Dan Stamper-Kurn. Interaction-enhanced group velocity of bosons in the flat band of an optical kagome lattice. *Phys. Rev. Lett.*, 125:133001, 2020. arXiv:2007.05928; [Partly verified: journal reference not seen on a primary record.].

[143] M. Raissi, P. Perdikaris, and G. E. Karniadakis. Physics-informed neural networks: A deep learning framework for solving forward and inverse problems involving nonlinear partial differential equations. *J. Comput. Phys.*, 378:686–707, 2019. https://www.sciencedirect.com/science/article/pii/S0021999118307125; [Partly verified: from search listings only.].

[144] Computing ground states of Bose–Einstein condensation by normalized deep neural network. Preprint, arXiv:2410.05319, 2024. Published in J. Comput. Phys. (ScienceDirect pii S0021999124007344); [Partly verified: from search listings only; authors, volume and year of the journal version not seen.].

[145] Physics-informed neural networks for the quantum droplets in binary Bose–Einstein condensates. *Chaos Solitons Fractals*, 2026. arXiv:2602.04590; ScienceDirect pii S0960077926001736; [Partly verified: from search listings only; authors and volume not seen.].

[146] PINN-based framework for soliton solutions of Gross Pitaevskii and nonlinear Schrödinger equations. Preprint, arXiv:2607.22685, 2026. [Partly verified: title from a search listing only; authors not seen.].

[147] Alexander B. Watson, Tianyu Kong, Allan H. MacDonald, and Mitchell Luskin. Bistritzer–MacDonald dynamics in twisted bilayer graphene. *J. Math. Phys.*, 64:031502, 2023. doi:10.1063/5.0115771. arXiv:2207.13767.

Research connected with the WPI and Vienna

Papers and programmes in this map whose authors or hosts are at the Wolfgang Pauli Institute, the University of Vienna or TU Wien (Atominstitut). First the work of the Mauser, Stimming, Mazets and Exl groups (stochastic classical fields and dissipative hydrodynamics of 1-D gases, the extended bosonic Josephson junction, time-splitting and absorbing-layer numerics, fast non-local kernels, physics-informed networks with hard constraints), then the Schmiedmayer atom-chip quantum field simulator and the Austrian cluster of excellence quantA. Numbering continues from the main list.

[148] H.-P. Stimming, N. J. Mauser, J. Schmiedmayer, and I. E. Mazets. Fluctuations and stochastic processes in one-dimensional many-body quantum systems. *Phys. Rev. Lett.*, 105:015301, 2010. doi:10.1103/PhysRevLett.105.015301. arXiv:0910.5337. [Partly verified: volume and article number from a search listing.].

[149] J.-F. Mennemann, I. E. Mazets, M. Pigneur, H. P. Stimming, N. J. Mauser, J. Schmiedmayer, and S. Erne. Relaxation in an extended bosonic Josephson junction. *Phys. Rev. Research*, 3:023197, 2021. doi:10.1103/PhysRevResearch.3.023197. arXiv:2012.05885; [Partly verified: journal reference not seen on a primary record; the 20 nK and 1-D-reduction statements are summary paraphrases.].

[150] F. Møller, N. Besse, I. E. Mazets, H. P. Stimming, and N. J. Mauser. The dissipative generalized hydrodynamic equations and their numerical solution. *J. Comput. Phys.*, 493, 2023. arXiv:2212.12349. [Partly verified: volume from the project documents; article number and DOI not confirmed.].

[151] W. Bao, N. J. Mauser, and H. P. Stimming. Effective one particle quantum dynamics of electrons: a numerical study of the Schrödinger-Poisson-Xα model. *Commun. Math. Sci.*, 1(4):809–828, 2003. [Partly verified: authors, volume and pages from the project documents and a listing; publisher page not reached.].

[152] T. S. Gutleb, N. J. Mauser, M. Ruggeri, and H. P. Stimming. A time splitting method for the three-dimensional linear Pauli equation. *Comput. Methods Appl. Math.*, 24(2):399–412, 2024. doi:10.1515/cmam-2023-0094. arXiv:2005.06072.

[153] H. P. Stimming, Xin Wen, and N. J. Mauser. Adaptive absorbing boundary layer for the nonlinear Schrödinger equation. *Comput. Methods Appl. Math.*, 24(3):663–690, 2024. doi:10.1515/cmam-2023-0096.

[154] L. Exl, N. J. Mauser, and Y. Zhang. Accurate and efficient computation of nonlocal potentials based on Gaussian-sum approximation. *J. Comput. Phys.*, 327:629–642, 2016. arXiv:1501.04438. [Partly verified: volume and first page from a listing; DOI not confirmed.].

[155] L. Exl, C. Abert, N. J. Mauser, T. Schrefl, H. P. Stimming, and D. Suess. FFT-based Kronecker product approximation to micromagnetic long-range interactions. *Math. Models Methods Appl. Sci.*, 24:1877–1901, 2014. doi:10.1142/S0218202514500109. arXiv:1212.3509.

[156] Sebastian Schaffer, Thomas Schrefl, Harald Oezelt, Norbert J. Mauser, and Lukas Exl. Physics aware machine learning for micromagnetic energy minimization: recent algorithmic developments. Preprint, arXiv:2409.12877, 2024.

[157] Alexander Kovacs, Lukas Exl, Alexander Kornell, Johann Fischbacher, Markus Hovorka, Markus Gusenbauer, Leoni Breth, Harald Oezelt, Masao Yano, Noritsugu Sakuma, Akihito Kinoshita, Tetsuya Shoji, Akira Kato, and Thomas Schrefl. Conditional physics informed neural networks. *Commun. Nonlinear Sci. Numer. Simul.*, 2022. arXiv:2104.02741; [Partly verified: volume, article number and year not seen.].

[158] T. Schweigler, V. Kasper, S. Erne, I. Mazets, B. Rauer, F. Cataldini, T. Langen, T. Gasenzer, J. Berges, and J. Schmiedmayer. Experimental characterization of a quantum many-body system via higher-order correlations. *Nature*, 545: 323–326, 2017. doi:10.1038/nature22310. [Partly verified: “atom chip” not in the abstract.].

[159] V. Kasper, J. Marino, S. Ji, V. Gritsev, J. Schmiedmayer, and E. Demler. Simulating a quantum commensurate-incommensurate phase transition using two Raman-coupled one-dimensional condensates. *Phys. Rev. B*, 101: 224102, 2020. doi:10.1103/PhysRevB.101.224102. arXiv:2002.12311; [Partly verified: journal reference from a search listing only.].

[160] Frederik Møller, Gabriel Fernández-Fernández, Thomas Schweigler, Paulin de Schoulepnikoff, Jörg Schmiedmayer, and Gorka Muñoz-Gil. Learning minimal representations of many-body physics from snapshots of a quantum simulator. *Phys. Rev. Research*, 2025. doi:10.1103/r7pj-gl7r. arXiv:2509.13821; [Partly verified: volume and article number not found; year is that of the arXiv version.].

[161] Universität Innsbruck. Austria’s quantum science is taking off. `https://www.uibk.ac.at/en/newsroom/2023/austrias-quantum-science-is-taking-off/`, 2023. Newsroom, 1 December 2023; accessed 21 September 2026; [Partly verified: Schmiedmayer’s research on coupled 1-D condensates not checked on this page.].

A Glossary

Terms from cold-atom physics, electronic moiré materials and quantum geometry, as this map uses them.

Moiré pattern. The long-wavelength beat between two periodic structures that are rotated or stretched against each other; its period L_M is much larger than the lattice constant at small twist.

Commensurate supercell. The common superlattice of two lattices at a commensurate twist, for square lattices $\tan(\theta/2) = 1/m$ with integer m; it makes band calculations periodic.

Registry (AA, AB). How the sites of the two layers line up locally: on top of each other in AA regions, alternating in AB regions. AA regions become the pockets where atoms collect.

Rabi coupling. A coherent drive (microwave or two-photon Raman) that converts atoms between two internal states; in a twisted-bilayer optical lattice it plays the role of interlayer tunnelling. Written Ω.

Tune-out wavelength. A laser wavelength at which the light shift of one atomic state vanishes, so that a lattice at that wavelength traps only the other state; used to give each spin state its own lattice.

Recoil energy. $E_R = \hbar^2 k^2/2m$ for the lattice wavevector k; the natural energy unit of optical lattices. Lattice depths V_0 are quoted in E_R.

Bose–Hubbard model. Lattice model of bosons with nearest-neighbour hopping J and on-site repulsion U; valid when the lowest band is deep and well separated.

Gutzwiller approximation. Mean-field product of single-site states; captures the Mott transition but no inter-site correlations.

Cluster Gutzwiller. Gutzwiller theory with small clusters of sites treated exactly, which restores short-range correlations.

Worm quantum Monte Carlo. Path-integral Monte Carlo that samples open world-line ends; exact for bosons without a sign problem and gives the superfluid fraction from winding numbers.

TWA and SPGPE. Truncated Wigner approximation: classical fields with quantum noise in the initial state. Stochastic projected GPE: a classical-field theory coupled to a thermal reservoir.

GPE. Gross–Pitaevskii equation: the mean-field equation for the condensate wavefunction; here with two components and a Rabi term.

BdG (Bogoliubov–de Gennes). Linearisation around a mean-field state; for a condensate it gives the Bogoliubov excitation spectrum, sound velocity and depletion.

Superfluid fraction. f_s: the part of the fluid that does not follow a slowly moving boundary, or equivalently the response to a phase twist; a tensor in anisotropic potentials.

Superfluid weight. D_s: the coefficient of the energy cost of a phase twist (superfluid stiffness); f_s is D_s normalised by its free-particle value.

Leggett bounds. Upper and lower bounds on f_s computed from the density profile alone; the upper bound is exact for a one-dimensional modulation of a weakly interacting condensate.

Quantum metric. $g_{\mu\nu}(\mathbf{k})$: the real part of the quantum geometric tensor; it measures how fast the Bloch state changes with $\mathbf{k}$.

Condensate quantum metric. The metric evaluated at the condensation momentum $\mathbf{k}_c$ in the basis of the condensate; it sets sound and stability of a flat-band condensate.

Minimal quantum metric. The integrated metric in the orbital basis that makes its diagonal components smallest; the quantity tied to the superfluid weight of paired fermions in an isolated flat band.

Quantum distance. $1 - |\langle u_{\mathbf{k}} | u_{\mathbf{k}'} \rangle|^2$ between two Bloch states; its small-step limit is the quantum metric.

Wannier function and spread. A localised orbital built from the Bloch states of a band; the spread is its mean-square size. A small spread means a band of nearly isolated orbitals.

Participation ratio. A measure of how many sites or cells a density spreads over, normalised to one for a uniform density; it falls towards zero as atoms collect in pockets.

TRIM. Time-reversal-invariant momentum: a point of the Brillouin zone equal to its negative up to a reciprocal-lattice vector (Γ, X, M in a square lattice).

BM model. Bistritzer–MacDonald continuum model of twisted bilayer graphene: Dirac cones of the two layers coupled by moiré-periodic tunnelling.

Magic angle. A twist angle at which the Dirac velocity of the BM model vanishes and the lowest moiré band flattens; the first lies near $1.05°$ [2].

Flat band. A Bloch band whose width is small compared with the interaction energy, so that interactions decide the ground state.

FCI / FQAH. Fractional Chern insulator, fractional quantum anomalous Hall state: fractional quantum Hall physics in a flat Chern band without an external magnetic field.

Bose glass. An insulating but compressible phase of bosons in a disordered or quasiperiodic potential, with local superfluid puddles but no global phase coherence.

Josephson array. A lattice of superfluid islands coupled by tunnelling; its superfluid weight is set by the inter-island Josephson coupling.