

Latent Semantic Indexing: A Probabilistic Analysis

C. Papadimitriou, P. Raghavan, H. Tamaki, S. Vempala

Seminar “Theoretical Topics in Data Science”

Vahe Eminyan

vahe.eminyan@rwth-aachen.de

30.01.2024

Overview

Introduction

LSI Background

Original Paper Overview and Emphasized Aspect

LSI by Random Projection

Summary and Newer Approaches

References

Introduction

Motivation

- Large datasets, often organized in tabular form, represented as **matrices**
 - Term-document matrix representing word occurrence in documents
 - Movie-user matrix representing watched movies of users

	Doc 1	Doc 2	...	Doc m
Term 1	0	1	...	1
Term 2	1	0	...	1
...
Term n	1	0	...	0



Documents

Terms $\begin{pmatrix} 0 & 1 & \dots & 1 \\ 1 & 0 & \dots & 1 \\ \vdots & \vdots & \ddots & \vdots \\ 1 & 0 & \dots & 0 \end{pmatrix}$

$n \times m$

Introduction

Motivation

- Large datasets, often organized in tabular form, represented as **matrices**
 - Term-document matrix representing word occurrence in documents
 - Movie-user matrix representing watched movies of users
- Interesting aspects
 - **Find** documents semantically associated with a **query**
 - **Recommend** a new movie to a user

	Doc 1	Doc 2	...	Doc m
Term 1	0	1	...	1
Term 2	1	0	...	1
...
Term n	1	0	...	0



Documents

Terms $\begin{pmatrix} 0 & 1 & \dots & 1 \\ 1 & 0 & \dots & 1 \\ \vdots & \vdots & \ddots & \vdots \\ 1 & 0 & \dots & 0 \end{pmatrix}$

$n \times m$

Latent Semantic Indexing

- LSI as an information retrieval method

Latent Semantic Indexing

- **LSI** as an information retrieval method
- Finds the **latent (hidden) semantic structure** of textual data. Solves the following problems:
 - Synonymy
 - Polysemy

Latent Semantic Indexing

- **LSI** as an information retrieval method
- Finds the **latent (hidden) semantic structure** of textual data. Solves the following problems:
 - **Synonymy**
 - **Polysemy**
- Represent term-document matrix as **product of three matrices**

Latent Semantic Indexing

- **LSI** as an information retrieval method
- Finds the **latent (hidden) semantic structure** of textual data. Solves the following problems:
 - **Synonymy**
 - **Polysemy**
- Represent term-document matrix as **product of three matrices**
- Answer queries with help of these matrices

Latent Semantic Indexing

- **LSI** as an information retrieval method
- Finds the **latent (hidden) semantic structure** of textual data. Solves the following problems:
 - **Synonymy**
 - **Polysemy**
- Represent term-document matrix as **product of three matrices**
- Answer queries with help of these matrices
- Based on **singular value decomposition** of the matrix

Singular Value Decomposition (SVD) [7]

- Any n by m matrix of rank r can be factored into

$$A_{n \times m} = U_{[n \times r]} D_{[r \times r]} (V_{[m \times r]})^T.$$

Singular Value Decomposition (SVD) [7]

- Any n by m matrix of rank r can be factored into

$$A_{n \times m} = U_{[n \times r]} D_{[r \times r]} (V_{[m \times r]})^T.$$

- U column-orthonormal matrix: **left singular vectors**

Singular Value Decomposition (SVD) [7]

- Any n by m matrix of rank r can be factored into

$$A_{n \times m} = U_{[n \times r]} D_{[r \times r]} (V_{[m \times r]})^T.$$

- U column-orthonormal matrix: left singular vectors
- V column-orthonormal matrix: right singular vectors

Singular Value Decomposition (SVD) [7]

- Any n by m matrix of rank r can be factored into

$$A_{n \times m} = U_{[n \times r]} D_{[r \times r]} (V_{[m \times r]})^T.$$

- U column-orthonormal matrix: **left singular vectors**
- V column-orthonormal matrix: **right singular vectors**
- D diagonal matrix: **Singular values** $\sigma_1 \geq \sigma_2 \geq \dots \geq \sigma_r$ in decreasing order

Singular Value Decomposition (SVD) [7]

- Any n by m matrix of rank r can be factored into

$$A_{n \times m} = U_{[n \times r]} D_{[r \times r]} (V_{[m \times r]})^T.$$

- U column-orthonormal matrix: **left singular vectors**
- V column-orthonormal matrix: **right singular vectors**
- D diagonal matrix: **Singular values** $\sigma_1 \geq \sigma_2 \geq \dots \geq \sigma_r$ in decreasing order
- Vector notation

$$A = U D V^T = \sum_{i=1}^r \sigma_i u_i v_i^T$$

Singular Value Decomposition (SVD) Example: Matrix A with rank $r = 3$

$$\begin{array}{c} \text{Terms} \end{array} \begin{array}{c} \text{Documents} \\ A \end{array} \begin{pmatrix} 1 & 0 & 1 \\ 1 & 1 & 0 \\ 0 & 1 & 0 \\ 0 & 1 & 1 \end{pmatrix} = \begin{array}{c} \text{Term-Topic similarity} \\ U \end{array} \begin{pmatrix} -0.48 & -0.79 & -0.11 \cdot 10^{-14} \\ -0.58 & 0.16 & 0.71 \\ \mathbf{-0.34} & \mathbf{0.56} & 0.42 \cdot 10^{-15} \\ -0.56 & 0.16 & -0.71 \end{pmatrix} \times \begin{array}{c} \text{Topic "importance"} \\ D \end{array} \begin{pmatrix} 2.1 & 0 & 0 \\ 0 & 1.26 & 0 \\ 0 & 0 & 1 \end{pmatrix}$$
$$\times \begin{array}{c} \text{Topic-Document similarity} \\ V^T \end{array} \begin{pmatrix} -0.5 & \mathbf{-0.71} & -0.5 \\ -0.5 & \mathbf{0.71} & -0.5 \\ 0.71 & 0.67 \cdot 10^{-15} & -0.711 \end{pmatrix}$$

Latent Semantic Indexing based on SVD

- LSI considers A_k the rank k approximation of A (l.e. keep only k most relevant topics)

$$A_k = U_k D_k V_k^T = \sum_{i=1}^k \sigma_i u_i v_i^T$$

$$\begin{array}{c} \text{Terms} \end{array} \begin{array}{c} \text{Documents} \\ \left(\begin{array}{ccc} 1.0 & 0.01 & 1.0 \\ 0.51 & 1.01 & 0.51 \\ 0.0 & 1.01 & 0.0 \\ 0.49 & 0.98 & 0.49 \end{array} \right) \\ A_k \end{array} = \begin{array}{c} \text{Term-Topic similarity} \\ \left(\begin{array}{cc} -0.48 & -0.79 \\ -0.58 & 0.16 \\ \mathbf{-0.34} & \mathbf{0.56} \\ -0.56 & 0.16 \end{array} \right) \\ U_k \end{array} \times \begin{array}{c} \text{Topic "importance"} \\ \left(\begin{array}{cc} 2.1 & 0 \\ 0 & 1.26 \end{array} \right) \\ D_k \end{array} \times \begin{array}{c} \text{Topic-Document similarity} \\ \left(\begin{array}{ccc} -0.5 & \mathbf{-0.71} & -0.5 \\ -0.5 & \mathbf{0.71} & -0.5 \end{array} \right) \\ V_k^T \end{array}$$

Latent Semantic Indexing based on SVD

- LSI considers A_k the rank k approximation of A (i.e. keep only k most relevant topics)

$$A_k = U_k D_k V_k^T = \sum_{i=1}^k \sigma_i u_i v_i^T$$

$$\begin{array}{c} \text{Terms} \end{array} \begin{array}{c} \text{Documents} \\ \left(\begin{array}{ccc} 1.0 & 0.01 & 1.0 \\ 0.51 & 1.01 & 0.51 \\ 0.0 & 1.01 & 0.0 \\ 0.49 & 0.98 & 0.49 \end{array} \right) \\ A_k \end{array} = \begin{array}{c} \text{Term-Topic similarity} \\ \left(\begin{array}{cc} -0.48 & -0.79 \\ -0.58 & 0.16 \\ \mathbf{-0.34} & \mathbf{0.56} \\ -0.56 & 0.16 \end{array} \right) \\ U_k \end{array} \times \begin{array}{c} \text{Topic "importance"} \\ \left(\begin{array}{cc} 2.1 & 0 \\ 0 & 1.26 \end{array} \right) \\ D_k \end{array} \times \begin{array}{c} \text{Topic-Document similarity} \\ \left(\begin{array}{ccc} -0.5 & \mathbf{-0.71} & -0.5 \\ -0.5 & \mathbf{0.71} & -0.5 \end{array} \right) \\ V_k^T \end{array}$$

- **Map** a query to k dimensional space with U_k , apply **cosine similarity** to find similar documents in $D_k V_k^T$

Latent Semantic Indexing based on SVD

Theorem (Eckart and Young [3])

Among all $n \times m$ matrices C of rank at most k , A_k is the one that minimizes $\|A - C\|_F^2 = \sum_{i,j} (A_{ij} - C_{ij})^2$, where F denotes the Frobenius norm of a matrix.

$$\begin{array}{c} \text{Terms} \end{array} \begin{array}{c} \text{Documents} \\ \begin{pmatrix} 1.0 & 0.01 & 1.0 \\ 0.51 & 1.01 & 0.51 \\ 0.0 & 1.01 & 0.0 \\ 0.49 & 0.98 & 0.49 \end{pmatrix} \\ A_k \end{array} \approx \begin{array}{c} \text{Terms} \end{array} \begin{array}{c} \text{Documents} \\ \begin{pmatrix} 1 & 0 & 1 \\ 1 & 1 & 0 \\ 0 & 1 & 0 \\ 0 & 1 & 1 \end{pmatrix} \\ A \end{array}$$

Original Paper Overview and Emphasized Aspect

- Strong empirical results of LSI

Original Paper Overview and Emphasized Aspect

- Strong empirical results of LSI
- Two important aspects
 - Why does LSI find **semantically related** documents?
 - How to **reduce the computational complexity** ?

Original Paper Overview and Emphasized Aspect

- Strong empirical results of LSI
- Two important aspects
 - Why does LSI find **semantically related** documents?
 - How to **reduce the computational complexity** ?
- Papadimitriou et al. [6] investigated both aspects:
 1. Under certain constraints semantically related documents are mapped to **similar vectors**
 2. Instead of LSI use **LSI by random projection**.
 - Map the original term-document matrix into a lower dimensional space
 - Use LSI on the lower dimensional matrix

Original Paper Overview and Emphasized Aspect

- Strong empirical results of LSI
- Two important aspects
 - Why does LSI find **semantically related** documents?
 - How to **reduce the computational complexity** ?
- Papadimitriou et al. [6] investigated both aspects:
 1. Under certain constraints semantically related documents are mapped to **similar vectors**
 2. Instead of LSI use **LSI by random projection**.
 - Map the original term-document matrix into a lower dimensional space
 - Use LSI on the lower dimensional matrix
- We focus on the **second** aspect

Random Projection for Dimensionality Reduction

Given a matrix $A \in \mathbb{R}^{n \times m}$ and a matrix $R \in \mathbb{R}^{\ell \times n}$. Use matrix R to **reduce the dimensionality** of matrix A while preserving pairwise distances between any two points:

$$B = \sqrt{\frac{n}{\ell}} \cdot R^T A \in \mathbb{R}^{\ell \times m}$$

Random Projection for Dimensionality Reduction

Given a matrix $A \in \mathbb{R}^{n \times m}$ and a matrix $R \in \mathbb{R}^{\ell \times n}$. Use matrix R to **reduce the dimensionality** of matrix A while preserving pairwise distances between any two points:

$$B = \sqrt{\frac{n}{\ell}} \cdot R^T A \in \mathbb{R}^{\ell \times m}$$

Lemma (Johnson and Lindenstrauss [4])

Let $v \in \mathbb{R}^n$ be a unit vector, let H be a random ℓ -dimensional subspace through the origin, and let the random variable X denote the square of the length of the projection of v onto H . Suppose $0 < \epsilon < 0.5$, and $24 \log n < \ell < \sqrt{n}$. Then, $E[X] = \frac{\ell}{n}$, and

$$Pr\left(\left|X - \frac{\ell}{n}\right| > \epsilon \frac{\ell}{n}\right) < 2\sqrt{\ell} e^{-(\ell-1)\epsilon^2/4}$$

LSI by Random Projection

Two-Step LSI

1. Apply a **random projection** onto ℓ dimensions on A . ($\ell > k$)

$$B = \sqrt{\frac{n}{\ell}} \cdot \begin{pmatrix} | & | & \cdots & | \\ r_1 & r_2 & \cdots & r_\ell \\ | & | & \cdots & | \end{pmatrix}^T \cdot A$$

2. Apply **rank $O(k)$ LSI**

Two-Step LSI

1. Apply a **random projection** onto ℓ dimensions on A . ($\ell > k$)

$$B = \sqrt{\frac{n}{\ell}} \cdot \begin{pmatrix} | & | & \cdots & | \\ r_1 & r_2 & \cdots & r_\ell \\ | & | & \cdots & | \end{pmatrix}^T \cdot A$$

2. Apply **rank $O(k)$ LSI**

- Improved computational complexity
- With high probability the original matrix A almost as good recovered as by directly using LSI (Formulation and proof of theorem later)

LSI by Random Projection

Comparison of Computational Time

Given the term-document matrix $A \in \mathbb{R}^{n \times m}$.

Time complexity of **one-step** LSI:

- LSI computation: $O(mnc)$ if A is sparse with about c nonzero entries per column

LSI by Random Projection

Comparison of Computational Time

Given the term-document matrix $A \in \mathbb{R}^{n \times m}$.

Time complexity of **one-step** LSI:

- LSI computation: $O(mnc)$ if A is sparse with about c nonzero entries per column

Time complexity of LSI by **random projection** :

- Random projection to ℓ dimensions: $O(m\ell)$
- LSI computation: $O(m\ell^2)$
- Together: $O(m\ell + m\ell^2) = O(m(\ell + \ell^2))$, with $\ell \in \Omega(\frac{\log n}{\epsilon^2})$
- Hence we get a time complexity: $O(m(\log^2 n + c \log n))$

$O(m(\log^2 n + c \log n))$ **better than** $O(mnc)$

Comparison of Both Matrices

- **A** : original term-document matrix
- **B** : original term-document matrix after random projection and scaling

$$\begin{array}{ccc} \textbf{A} & \longrightarrow & \textbf{B} \\ \begin{pmatrix} 0 & 1 & \dots & 1 \\ 1 & 0 & \dots & 1 \\ \vdots & \vdots & \ddots & \vdots \\ 1 & 0 & \dots & 0 \end{pmatrix} & & \begin{pmatrix} 0 & 0.47 & \dots & 0.47 \\ \vdots & \vdots & \ddots & \vdots \\ 0.47 & 0 & \dots & 0 \end{pmatrix} \\ n \times m & & \ell \times m \end{array}$$

Comparison of Both Matrices

- **A** : original term-document matrix
- **B** : original term-document matrix after random projection and scaling
- $\ell \in \Omega(\frac{\log n}{\epsilon^2})$, with $\epsilon \in (0, 0.5)$

$$\begin{array}{ccc} \textbf{A} & & \textbf{B} \\ \begin{pmatrix} 0 & 1 & \dots & 1 \\ 1 & 0 & \dots & 1 \\ \vdots & \vdots & \ddots & \vdots \\ 1 & 0 & \dots & 0 \end{pmatrix} & \longrightarrow & \begin{pmatrix} 0 & 0.47 & \dots & 0.47 \\ \vdots & \vdots & \ddots & \vdots \\ 0.47 & 0 & \dots & 0 \end{pmatrix} \\ n \times m & & \ell \times m \end{array}$$

Comparison of Both Matrices

- **A** : original term-document matrix
- **B** : original term-document matrix after random projection and scaling
- $\ell \in \Omega(\frac{\log n}{\epsilon^2})$, with $\epsilon \in (0, 0.5)$
- Dimensionality reduction for each document ($\ell \ll n$)

$$\begin{array}{ccc} \textbf{A} & & \textbf{B} \\ \begin{pmatrix} 0 & 1 & \dots & 1 \\ 1 & 0 & \dots & 1 \\ \vdots & \vdots & \ddots & \vdots \\ 1 & 0 & \dots & 0 \end{pmatrix} & \longrightarrow & \begin{pmatrix} 0 & 0.47 & \dots & 0.47 \\ \vdots & \vdots & \ddots & \vdots \\ 0.47 & 0 & \dots & 0 \end{pmatrix} \\ n \times m & & \ell \times m \end{array}$$

Background and Notation for the Proof

Vector notations of SVD:

$$A = \sum_{i=1}^n \sigma_i u_i v_i^T, \quad A_k = \sum_{i=1}^k \sigma_i u_i v_i^T, \quad B = \sum_{i=1}^{\ell} \lambda_i a_i b_i^T, \quad B_{2k} = A \sum_{i=1}^{2k} b_i b_i^T.$$

Background and Notation for the Proof

Vector notations of SVD:

$$A = \sum_{i=1}^n \sigma_i u_i v_i^T, \quad A_k = \sum_{i=1}^k \sigma_i u_i v_i^T, \quad B = \sum_{i=1}^{\ell} \lambda_i a_i b_i^T, \quad B_{2k} = A \sum_{i=1}^{2k} b_i b_i^T.$$

- A : **original** term-document matrix

Background and Notation for the Proof

Vector notations of SVD:

$$A = \sum_{i=1}^n \sigma_i u_i v_i^T, \quad A_k = \sum_{i=1}^k \sigma_i u_i v_i^T, \quad B = \sum_{i=1}^{\ell} \lambda_i a_i b_i^T, \quad B_{2k} = A \sum_{i=1}^{2k} b_i b_i^T.$$

- A : **original** term-document matrix
- A_k : **rank k approximation** of A

Background and Notation for the Proof

Vector notations of SVD:

$$A = \sum_{i=1}^n \sigma_i u_i v_i^T, \quad A_k = \sum_{i=1}^k \sigma_i u_i v_i^T, \quad B = \sum_{i=1}^{\ell} \lambda_i a_i b_i^T, \quad B_{2k} = A \sum_{i=1}^{2k} b_i b_i^T.$$

- A : **original** term-document matrix
- A_k : **rank k approximation** of A
- B : matrix after **randomly projecting** and **scaling** A

Background and Notation for the Proof

Vector notations of SVD:

$$A = \sum_{i=1}^n \sigma_i u_i v_i^T, \quad A_k = \sum_{i=1}^k \sigma_i u_i v_i^T, \quad B = \sum_{i=1}^{\ell} \lambda_i a_i b_i^T, \quad B_{2k} = A \sum_{i=1}^{2k} b_i b_i^T.$$

- A : **original** term-document matrix
- A_k : **rank k approximation** of A
- B : matrix after **randomly projecting** and **scaling** A
- B_{2k} : **rank $2k$ approximation** of A

Background and Notation for the Proof

Lemma (3)

Let ϵ be an arbitrary positive constant. If $\ell \geq c((\log n)/\epsilon^2)$ for a sufficiently large constant c then, for $p = 1, \dots, \ell$

$$\lambda_p^2 \geq \frac{1}{k} \left[(1 - \epsilon) \sum_{i=1}^k \sigma_i^2 - \sum_{j=1}^{p-1} \lambda_j^2 \right].$$

Background and Notation for the Proof

Lemma (3)

Let ϵ be an arbitrary positive constant. If $\ell \geq c((\log n)/\epsilon^2)$ for a sufficiently large constant c then, for $p = 1, \dots, \ell$

$$\lambda_p^2 \geq \frac{1}{k} \left[(1 - \epsilon) \sum_{i=1}^k \sigma_i^2 - \sum_{j=1}^{p-1} \lambda_j^2 \right].$$

Corollary (4)

$$\sum_{p=1}^{2k} \lambda_p^2 \geq (1 - \epsilon) \|A_k\|_F^2.$$

Background and Notation for the Proof

Lemma (5)

$$\|A - A_k\|_F^2 = \sum_{i=k+1}^n \sigma_i^2.$$

Background and Notation for the Proof

Lemma (5)

$$\|A - A_k\|_F^2 = \sum_{i=k+1}^n \sigma_i^2.$$

Theorem (Parsevals identity [2])

Let b_1, \dots, b_n be an orthonormal basis for a space S . Then for each $s \in S$, $|s|^2 = \sum_{i=1}^n (sb_i)^2$.

LSI by Random Projection

Main Theorem

Theorem (Papadimitriou et al. [6])

$$\|A - B_{2k}\|_F^2 \leq \|A - A_k\|_F^2 + 2\epsilon \|A\|_F^2$$

where $\epsilon \in (0, 0.5)$

Informally, the theorem states that the original matrix A after applying **random projection** and then **LSI** is with high probability almost as good **recovered** as by using **one-step LSI** on the original matrix.

LSI by Random Projection

Theorem (Papadimitriou et al. [6])

$$\|A - B_{2k}\|_F^2 \leq \|A - A_k\|_F^2 + 2\epsilon \|A\|_F^2$$

where $\epsilon \in (0, 0.5)$

Proof

We have

$$A = \sum_{i=1}^n \sigma_i u_i v_i^T, \quad A_k = \sum_{i=1}^k \sigma_i u_i v_i^T, \quad B = \sum_{i=1}^{\ell} \lambda_i a_i b_i^T, \quad B_{2k} = A \sum_{i=1}^{2k} b_i b_i^T.$$

b_1, \dots, b_n Are orthonormal vectors **spanning** the **row space** of A and B_{2k} .

Hence using **the Parseval's** identity we can write:

$$\|A - B_{2k}\|_F^2 = \sum_{i=1}^n |(A - B_{2k})b_i|^2. \tag{1}$$

LSI by Random Projection

Theorem (Papadimitriou et al. [6])

$$\|A - B_{2k}\|_F^2 \leq \|A - A_k\|_F^2 + 2\epsilon \|A\|_F^2$$

where $\epsilon \in (0, 0.5)$

Proof

We have

$$A = \sum_{i=1}^n \sigma_i u_i v_i^T, \quad A_k = \sum_{i=1}^k \sigma_i u_i v_i^T, \quad B = \sum_{i=1}^\ell \lambda_i a_i b_i^T, \quad B_{2k} = A \sum_{i=1}^{2k} b_i b_i^T.$$

b_1, \dots, b_n Are orthonormal vectors **spanning** the **row space** of A and B_{2k} .

Hence using **the Parseval's** identity we can write:

$$\|A - B_{2k}\|_F^2 = \sum_{i=1}^n |(A - B_{2k})b_i|^2. \quad (1)$$

For $i = 1, \dots, 2k$, because $b_i^T b_i = 1$, we have

$$(A - B_{2k})b_i = Ab_i - Ab_i = 0, \quad (2)$$

LSI by Random Projection

Theorem (Papadimitriou et al. [6])

$$\|A - B_{2k}\|_F^2 \leq \|A - A_k\|_F^2 + 2\epsilon \|A\|_F^2$$

where $\epsilon \in (0, 0.5)$

Proof

We have

$$A = \sum_{i=1}^n \sigma_i u_i v_i^T, \quad A_k = \sum_{i=1}^k \sigma_i u_i v_i^T, \quad B = \sum_{i=1}^{\ell} \lambda_i a_i b_i^T, \quad B_{2k} = A \sum_{i=1}^{2k} b_i b_i^T.$$

b_1, \dots, b_n Are orthonormal vectors **spanning** the **row space** of A and B_{2k} .

Hence using **the Parseval's** identity we can write:

$$\|A - B_{2k}\|_F^2 = \sum_{i=1}^n |(A - B_{2k})b_i|^2. \quad (1)$$

For $i = 1, \dots, 2k$, because $b_i^T b_i = 1$, we have

$$(A - B_{2k})b_i = Ab_i - Ab_i = 0, \quad (2)$$

and for $i = 2k + 1, \dots, n$, because $b_j^T b_i = 0$, we have

$$(A - B_{2k})b_i = Ab_i. \quad (3)$$

LSI by Random Projection

Theorem (Papadimitriou et al. [6])

$$\|A - B_{2k}\|_F^2 \leq \|A - A_k\|_F^2 + 2\epsilon \|A\|_F^2$$

where $\epsilon \in (0, 0.5)$

Proof (continued)

Now we continue from the equation

$$\|A - B_{2k}\|_F^2 = \sum_{i=1}^n |(A - B_{2k})b_i|^2 \tag{4}$$

$$= \sum_{i=2k+1}^n |Ab_i|^2 \tag{5}$$

(7)

LSI by Random Projection

Theorem (Papadimitriou et al. [6])

$$\|A - B_{2k}\|_F^2 \leq \|A - A_k\|_F^2 + 2\epsilon \|A\|_F^2$$

where $\epsilon \in (0, 0.5)$

Proof (continued)

Now we continue from the equation

$$\|A - B_{2k}\|_F^2 = \sum_{i=1}^n |(A - B_{2k})b_i|^2 \tag{4}$$

$$= \sum_{i=2k+1}^n |Ab_i|^2 \tag{5}$$

$$= \sum_{i=1}^n |Ab_i|^2 - \sum_{i=1}^{2k} |Ab_i|^2 \tag{6}$$

(7)

LSI by Random Projection

Theorem (Papadimitriou et al. [6])

$$\|A - B_{2k}\|_F^2 \leq \|A - A_k\|_F^2 + 2\epsilon \|A\|_F^2$$

where $\epsilon \in (0, 0.5)$

Proof (continued)

Now we continue from the equation

$$\|A - B_{2k}\|_F^2 = \sum_{i=1}^n |(A - B_{2k})b_i|^2 \quad (4)$$

$$= \sum_{i=2k+1}^n |Ab_i|^2 \quad (5)$$

$$= \sum_{i=1}^n |Ab_i|^2 - \sum_{i=1}^{2k} |Ab_i|^2 \quad (6)$$

$$\stackrel{\text{Parseval's id.}}{=} \|A\|_F^2 - \sum_{i=1}^{2k} |Ab_i|^2 \quad (7)$$

LSI by Random Projection

Theorem (Papadimitriou et al. [6])

$$\|A - B_{2k}\|_F^2 \leq \|A - A_k\|_F^2 + 2\epsilon \|A\|_F^2$$

where $\epsilon \in (0, 0.5)$

Proof (continued)

On the other hand, we have

$$\|A - A_k\|_F^2 \stackrel{\text{Lemma 5}}{=} \sum_{i=k+1}^n \sigma_i^2 \tag{8}$$

(9)

LSI by Random Projection

Theorem (Papadimitriou et al. [6])

$$\|A - B_{2k}\|_F^2 \leq \|A - A_k\|_F^2 + 2\epsilon \|A\|_F^2$$

where $\epsilon \in (0, 0.5)$

Proof (continued)

On the other hand, we have

$$\|A - A_k\|_F^2 \stackrel{\text{Lemma 5}}{=} \sum_{i=k+1}^n \sigma_i^2 \quad (8)$$

$$\stackrel{\text{Frob. norm [5]}}{=} \|A\|_F^2 - \|A_k\|_F^2. \quad (9)$$

LSI by Random Projection

Theorem (Papadimitriou et al. [6])

$$\|A - B_{2k}\|_F^2 \leq \|A - A_k\|_F^2 + 2\epsilon \|A\|_F^2$$

where $\epsilon \in (0, 0.5)$

Proof (continued)

Now we consider

$$\|A - B_{2k}\|_F^2 - \|A - A_k\|_F^2 = \|A\|_F^2 - \sum_{i=1}^{2k} |Ab_i|^2 - (\|A\|_F^2 - \|A_k\|_F^2) \quad (10)$$

(11)

LSI by Random Projection

Theorem (Papadimitriou et al. [6])

$$\|A - B_{2k}\|_F^2 \leq \|A - A_k\|_F^2 + 2\epsilon \|A\|_F^2$$

where $\epsilon \in (0, 0.5)$

Proof (continued)

Now we consider

$$\|A - B_{2k}\|_F^2 - \|A - A_k\|_F^2 = \|A\|_F^2 - \sum_{i=1}^{2k} |Ab_i|^2 - (\|A\|_F^2 - \|A_k\|_F^2) \quad (10)$$

$$= \|A_k\|_F^2 - \sum_{i=1}^{2k} |Ab_i|^2, \quad (11)$$

LSI by Random Projection

Theorem (Papadimitriou et al. [6])

$$\|A - B_{2k}\|_F^2 \leq \|A - A_k\|_F^2 + 2\epsilon \|A\|_F^2$$

where $\epsilon \in (0, 0.5)$

Proof (continued)

Now we consider

$$\|A - B_{2k}\|_F^2 - \|A - A_k\|_F^2 = \|A\|_F^2 - \sum_{i=1}^{2k} |Ab_i|^2 - (\|A\|_F^2 - \|A_k\|_F^2) \quad (10)$$

$$= \|A_k\|_F^2 - \sum_{i=1}^{2k} |Ab_i|^2, \quad (11)$$

that is equivalent to

$$\|A - B_{2k}\|_F^2 = \|A - A_k\|_F^2 + (\|A_k\|_F^2 - \sum_{i=1}^{2k} |Ab_i|^2) \quad (12)$$

LSI by Random Projection

Theorem (Papadimitriou et al. [6])

$$\|A - B_{2k}\|_F^2 \leq \|A - A_k\|_F^2 + 2\epsilon \|A\|_F^2$$

where $\epsilon \in (0, 0.5)$

Proof (continued)

For the next step, we show

$$(1 + \epsilon) \sum_{i=1}^{2k} |Ab_i|^2 \geq \sum_{i=1}^{2k} \lambda_i^2. \tag{13}$$

We write

(16)

LSI by Random Projection

Theorem (Papadimitriou et al. [6])

$$\|A - B_{2k}\|_F^2 \leq \|A - A_k\|_F^2 + 2\epsilon \|A\|_F^2$$

where $\epsilon \in (0, 0.5)$

Proof (continued)

For the next step, we show

$$(1 + \epsilon) \sum_{i=1}^{2k} |Ab_i|^2 \geq \sum_{i=1}^{2k} \lambda_i^2. \quad (13)$$

We write

$$\sum_{i=1}^{2k} \lambda_i^2 \stackrel{|Bb_i|=\lambda_i}{=} \sum_{i=1}^{2k} |Bb_i|^2 \quad (14)$$

(16)

LSI by Random Projection

Theorem (Papadimitriou et al. [6])

$$\|A - B_{2k}\|_F^2 \leq \|A - A_k\|_F^2 + 2\epsilon \|A\|_F^2$$

where $\epsilon \in (0, 0.5)$

Proof (continued)

For the next step, we show

$$(1 + \epsilon) \sum_{i=1}^{2k} |Ab_i|^2 \geq \sum_{i=1}^{2k} \lambda_i^2. \quad (13)$$

We write

$$\sum_{i=1}^{2k} \lambda_i^2 \stackrel{|Bb_i|=\lambda_i}{=} \sum_{i=1}^{2k} |Bb_i|^2 \quad (14)$$

$$\stackrel{\text{subst. B}}{=} \sum_{i=1}^{2k} \left| \sqrt{\frac{n}{\ell}} R^T (Ab_i) \right|^2 \quad (15)$$

$$(16)$$

LSI by Random Projection

Theorem (Papadimitriou et al. [6])

$$\|A - B_{2k}\|_F^2 \leq \|A - A_k\|_F^2 + 2\epsilon \|A\|_F^2$$

where $\epsilon \in (0, 0.5)$

Proof (continued)

For the next step, we show

$$(1 + \epsilon) \sum_{i=1}^{2k} |Ab_i|^2 \geq \sum_{i=1}^{2k} \lambda_i^2. \quad (13)$$

We write

$$\sum_{i=1}^{2k} \lambda_i^2 \stackrel{|Bb_i|=\lambda_i}{=} \sum_{i=1}^{2k} |Bb_i|^2 \quad (14)$$

$$\stackrel{\text{subst. B}}{=} \sum_{i=1}^{2k} \left| \sqrt{\frac{n}{\ell}} R^T(Ab_i) \right|^2 \quad (15)$$

$$= \sum_{i=1}^{2k} \frac{n}{\ell} |R^T(Ab_i)|^2 \quad (16)$$

LSI by Random Projection

Theorem (Papadimitriou et al. [6])

$$\|A - B_{2k}\|_F^2 \leq \|A - A_k\|_F^2 + 2\epsilon \|A\|_F^2$$

where $\epsilon \in (0, 0.5)$

Proof (continued)

Now from the **Johnson-Lindenstrauss lemma** [4] for very large $\ell \in \Omega((\log n)/\epsilon^2)$ we have for each i

$$\frac{n}{\ell} |R^T(Ab_i)|^2 \leq (1 + \epsilon) |Ab_i|^2 \quad (17)$$

with **high probability** .

LSI by Random Projection

Theorem (Papadimitriou et al. [6])

$$\|A - B_{2k}\|_F^2 \leq \|A - A_k\|_F^2 + 2\epsilon \|A\|_F^2$$

where $\epsilon \in (0, 0.5)$

Proof (continued)

Now from the **Johnson-Lindenstrauss lemma** [4] for very large $\ell \in \Omega((\log n)/\epsilon^2)$ we have for each i

$$\frac{n}{\ell} |R^T(Ab_i)|^2 \leq (1 + \epsilon) |Ab_i|^2 \quad (17)$$

with **high probability** .

Hence with a high probability

$$(1 + \epsilon) \sum_{i=1}^{2k} |Ab_i|^2 \geq \sum_{i=1}^{2k} \lambda_i^2. \quad (18)$$

LSI by Random Projection

Theorem (Papadimitriou et al. [6])

$$\|A - B_{2k}\|_F^2 \leq \|A - A_k\|_F^2 + 2\epsilon \|A\|_F^2$$

where $\epsilon \in (0, 0.5)$

Proof (continued)

Now we have

$$\sum_{i=1}^{2k} |Ab_i|^2 \geq \frac{1}{(1 + \epsilon)} \sum_{i=1}^{2k} \lambda_i^2 \tag{19}$$

(21)

LSI by Random Projection

Theorem (Papadimitriou et al. [6])

$$\|A - B_{2k}\|_F^2 \leq \|A - A_k\|_F^2 + 2\epsilon \|A\|_F^2$$

where $\epsilon \in (0, 0.5)$

Proof (continued)

Now we have

$$\sum_{i=1}^{2k} |Ab_i|^2 \geq \frac{1}{(1 + \epsilon)} \sum_{i=1}^{2k} \lambda_i^2 \tag{19}$$

$$\stackrel{\text{Cor. 4}}{\geq} \frac{(1 - \epsilon)}{(1 + \epsilon)} \|A_k\|_F^2 \tag{20}$$

(21)

LSI by Random Projection

Theorem (Papadimitriou et al. [6])

$$\|A - B_{2k}\|_F^2 \leq \|A - A_k\|_F^2 + 2\epsilon \|A\|_F^2$$

where $\epsilon \in (0, 0.5)$

Proof (continued)

Now we have

$$\sum_{i=1}^{2k} |Ab_i|^2 \geq \frac{1}{(1 + \epsilon)} \sum_{i=1}^{2k} \lambda_i^2 \tag{19}$$

$$\stackrel{\text{Cor. 4}}{\geq} \frac{(1 - \epsilon)}{(1 + \epsilon)} \|A_k\|_F^2 \tag{20}$$

$$\geq (1 - 2\epsilon) \|A_k\|_F^2 \tag{21}$$

LSI by Random Projection

Theorem (Papadimitriou et al. [6])

$$\|A - B_{2k}\|_F^2 \leq \|A - A_k\|_F^2 + 2\epsilon \|A\|_F^2$$

where $\epsilon \in (0, 0.5)$

Proof (continued)

Now we have

$$\sum_{i=1}^{2k} |Ab_i|^2 \geq \frac{1}{(1 + \epsilon)} \sum_{i=1}^{2k} \lambda_i^2 \quad (19)$$

$$\stackrel{\text{Cor. 4}}{\geq} \frac{(1 - \epsilon)}{(1 + \epsilon)} \|A_k\|_F^2 \quad (20)$$

$$\geq (1 - 2\epsilon) \|A_k\|_F^2 \quad (21)$$

i.e.

$$\sum_{i=1}^{2k} |Ab_i|^2 \geq (1 - 2\epsilon) \|A_k\|_F^2 \quad (22)$$

LSI by Random Projection

Theorem (Papadimitriou et al. [6])

$$\|A - B_{2k}\|_F^2 \leq \|A - A_k\|_F^2 + 2\epsilon \|A\|_F^2$$

where $\epsilon \in (0, 0.5)$

Proof (continued)

Remember the Equation (12):

$$\|A - B_{2k}\|_F^2 = \|A - A_k\|_F^2 + (\|A_k\|_F^2 - \sum_{i=1}^{2k} |Ab_i|^2)$$

Now we **substitute** the result of Equation (22) in equation (12):

(24)

LSI by Random Projection

Theorem (Papadimitriou et al. [6])

$$\|A - B_{2k}\|_F^2 \leq \|A - A_k\|_F^2 + 2\epsilon \|A\|_F^2$$

where $\epsilon \in (0, 0.5)$

Proof (continued)

Remember the Equation (12):

$$\|A - B_{2k}\|_F^2 = \|A - A_k\|_F^2 + (\|A_k\|_F^2 - \sum_{i=1}^{2k} |Ab_i|^2)$$

Now we **substitute** the result of Equation (22) in equation (12):

$$\|A - B_{2k}\|_F^2 \leq \|A - A_k\|_F^2 + \|A_k\|_F^2 - (1 - 2\epsilon) \|A_k\|_F^2 \tag{23}$$

(24)

LSI by Random Projection

Theorem (Papadimitriou et al. [6])

$$\|A - B_{2k}\|_F^2 \leq \|A - A_k\|_F^2 + 2\epsilon \|A\|_F^2$$

where $\epsilon \in (0, 0.5)$

Proof (continued)

Remember the Equation (12):

$$\|A - B_{2k}\|_F^2 = \|A - A_k\|_F^2 + (\|A_k\|_F^2 - \sum_{i=1}^{2k} |Ab_i|^2)$$

Now we **substitute** the result of Equation (22) in equation (12):

$$\|A - B_{2k}\|_F^2 \leq \|A - A_k\|_F^2 + \|A_k\|_F^2 - (1 - 2\epsilon) \|A_k\|_F^2 \quad (23)$$

$$\iff \|A - B_{2k}\|_F^2 \leq \|A - A_k\|_F^2 + 2\epsilon \|A_k\|_F^2 \quad (24)$$

LSI by Random Projection

Theorem (Papadimitriou et al. [6])

$$\|A - B_{2k}\|_F^2 \leq \|A - A_k\|_F^2 + 2\epsilon \|A\|_F^2$$

where $\epsilon \in (0, 0.5)$

Proof (continued)

Remember the Equation (12):

$$\|A - B_{2k}\|_F^2 = \|A - A_k\|_F^2 + (\|A_k\|_F^2 - \sum_{i=1}^{2k} |Ab_i|^2)$$

Now we **substitute** the result of Equation (22) in equation (12):

$$\|A - B_{2k}\|_F^2 \leq \|A - A_k\|_F^2 + \|A_k\|_F^2 - (1 - 2\epsilon) \|A_k\|_F^2 \quad (23)$$

$$\iff \|A - B_{2k}\|_F^2 \leq \|A - A_k\|_F^2 + 2\epsilon \|A_k\|_F^2 \quad (24)$$

Due to the formulation of **Frobenius norm** as in Lemma 5, we have $\|A\|_F^2 \geq \|A_k\|_F^2$.

Hence

$$\|A - B_{2k}\|_F^2 \leq \|A - A_k\|_F^2 + 2\epsilon \|A\|_F^2. \quad (25)$$

□

Summary and Newer Approaches

- Latent semantic analysis: SVD-based technique for information retrieval

Summary and Newer Approaches

- Latent semantic analysis: SVD-based technique for information retrieval
- Papadimitriou et al. analysed two important aspects [6]
 - Why does LSI find **semantically related** documents?
 - How to **reduce the computational time** ? (Our main focus)

Summary and Newer Approaches


- Latent semantic analysis: SVD-based technique for information retrieval
- Papadimitriou et al. analysed two important aspects [6]
 - Why does LSI find **semantically related** documents?
 - How to **reduce the computational time** ? (Our main focus)
- LSI by random projection: **reduction of computational complexity**, while preventing the **expressiveness** of original matrix with high probability.


Summary and Newer Approaches


- Latent semantic analysis: SVD-based technique for information retrieval
- Papadimitriou et al. analysed two important aspects [6]
 - Why does LSI find **semantically related** documents?
 - How to **reduce the computational time** ? (Our main focus)
- LSI by random projection: **reduction of computational complexity**, while preventing the **expressiveness** of original matrix with high probability.
- There are newer techniques based on **neural networks** [8, 1]


References

 William L. Hamilton, Rex Ying, and Jure Leskovec.
Representation learning on graphs: Methods and applications.
IEEE Data Eng. Bull., 40(3):52–74, 2017.
URL: <http://sites.computer.org/debull/A17sept/p52.pdf>.


 Leslie Hogben, editor.
Handbook of Linear Algebra.
Chapman and Hall/CRC, 2nd edition, 2013.
<https://doi.org/10.1201/b16113>.

 C. Reinsch J. H. Wilkinson.
Handbook for Automatic Computation.
Springer Berlin, Heidelberg, volume ii: linear algebra edition, 1971.


 William B Johnson.
Extensions of lipshitz mapping into hilbert space.
In *Conference modern analysis and probability, 1984*, pages 189–206, 1984.

 Changxue Ma, Y. Kamp, and L.F. Willems.
A frobenius norm approach to glottal closure detection from the speech signal.
IEEE Transactions on Speech and Audio Processing, 2(2):258–265, 1994.
[doi:10.1109/89.279274](https://doi.org/10.1109/89.279274).

 Christos H. Papadimitriou, Prabhakar Raghavan, Hisao Tamaki, and Santosh Vempala.
Latent semantic indexing: A probabilistic analysis.
Journal of Computer and System Sciences, 61(2):217–235, 2000.
URL: <https://www.sciencedirect.com/science/article/pii/S0022000000917112>, [doi:10.1006/jcss.2000.1711](https://doi.org/10.1006/jcss.2000.1711).

 Gilbert Strang.
Linear Algebra and Its Applications.
Cengage Learning, 4th edition edition, 2005.

References

-  [Liang Yao, Chengsheng Mao, and Yuan Luo.](#)
Graph convolutional networks for text classification.
In Proceedings of the AAAI conference on artificial intelligence, volume 33, pages 7370–7377, 2019.