

INFLUENZA

A Comprehensive Evidence-Based Analysis of Human, Animal, and Wildlife Mortality

Molecular Mechanisms • Zoonotic Transmission • Clinical Management • Control of Spread •
Vaccination • One Health Surveillance • Pandemic Preparedness



3-7-12 Systems Framework

A proposed systems architecture for translating influenza evidence into measurable preparedness, response and research priorities.

Vahid Nezamivand Chegane

ORCID: 0009-0003-6477-091X

vahidmaykoltarh@yahoo.com

Evidence and data cutoff: 10 September 2026

Abstract

Influenza viruses remain a major cause of respiratory morbidity and mortality and a continuing zoonotic and pandemic preparedness challenge. The global seasonal burden is estimated at approximately one billion cases annually, including 3–5 million severe illnesses and 290,000–650,000 respiratory deaths. Historical pandemics demonstrate that mortality can increase dramatically when a novel influenza virus enters a population with limited pre-existing immunity.

This manuscript integrates human mortality measurement, animal and wildlife disease ecology, molecular mechanisms of host adaptation, clinical management, antiviral therapy, vaccination, laboratory diagnostics, genomic surveillance, outbreak control, risk communication and pandemic preparedness. Particular attention is given to avian influenza A(H5N1), for which WHO maintains a time-stamped global register of laboratory-confirmed human infections. The manuscript deliberately distinguishes confirmed-case counts from population infection denominators and separates biological animal mortality from disease-control depopulation.

The paper introduces the 3–7–12 Systems Framework as a proposed conceptual and operational model: three interacting system layers, seven intervention points and twelve measurable preparedness actions. The framework is not presented as an established scientific law; its value is a testable hypothesis that can be evaluated through implementation studies, response-time metrics, laboratory performance and clinical outcomes.

Keywords: Influenza; Influenza A; H5N1; H7N9; H9N2; zoonotic influenza; One Health; mortality; pandemic preparedness; vaccination; antivirals; genomic surveillance; outbreak control; 3–7–12 Systems Framework.

Scientific integrity principle. Numerical claims are classified as observed surveillance counts, model-based estimates, historical ranges, or proposed synthesis. Exactness is not substituted for evidence.


1. Scope, Research Questions and Evidence Architecture

The purpose of this review is not merely to catalogue influenza deaths. It is to connect mortality measurement to the biological, clinical, ecological and institutional mechanisms that determine whether an outbreak is detected early, contained effectively and treated appropriately.

Research domain	Core question	Preferred evidence
Mortality	How many people die, under which definition?	Vital statistics, surveillance, excess-mortality models
Zoonotic risk	Where are human–animal spillover interfaces?	Animal surveillance, exposure investigations, genomics
Molecular biology	Which mechanisms influence host range and phenotype?	Virology, structural biology, experimental systems
Clinical care	Which interventions reduce severe outcomes?	Guidelines, randomized trials, observational evidence
Control	Which actions shorten transmission and response time?	Field epidemiology, implementation science
Preparedness	How can systems detect and adapt before widespread transmission?	GISRS, risk assessment, exercises, M&E;

2. Human Influenza Mortality and Burden


WHO estimates that seasonal influenza produces around one billion cases each year, including 3–5 million severe illnesses and 290,000–650,000 respiratory deaths. These are population-level burden estimates, not a count of laboratory-confirmed influenza deaths. The respiratory endpoint also excludes some non-respiratory influenza-associated deaths, such as cardiovascular complications, depending on the modelling framework.



The distinction between influenza-attributable mortality and laboratory-confirmed influenza deaths is fundamental. Many deaths occur outside the testing pathway; conversely, a positive influenza test does not establish that influenza was the sole cause of death.

2.1 Historical Pandemics

Pandemic	Period	Virus	Global mortality evidence
1918	1918-1919	A(H1N1)	At least tens of millions; commonly cited estimates are approximately 40-50 million or higher depending on methodology.
1957	1957-1958	A(H2N2)	Low-million global mortality estimates; exact totals remain uncertain.
1968	1968-1969	A(H3N2)	Low-million global mortality estimates; exact totals remain uncertain.
2009	2009-2010	A(H1N1)pdm09	CDC estimated 151,700-575,400 global deaths during the first year of circulation.



The 2009 estimate is model-based rather than a universal death registry. CDC reported the range of 151,700-575,400 deaths worldwide in the first year of A(H1N1)pdm09 circulation.

2.2 Mortality Metrics

Case-fatality ratio: $CFR = D / C \times 100\%$, where D and C refer to deaths and cases defined within the same surveillance framework.

Infection-fatality ratio: $IFR = D / I \times 100\%$, where I includes detected and undetected infections. IFR therefore requires an estimated infection denominator.

Mortality rate: $M = D / N \times k$, where N is the population at risk and k is the reporting multiplier (for example, 100,000).

Excess mortality: Excess deaths = observed deaths – expected deaths under a counterfactual baseline. The result depends on baseline construction and model assumptions.

3. Zoonotic Influenza and A(H5N1)

A(H5N1) is a central One Health concern because highly pathogenic avian influenza viruses have caused extensive outbreaks in birds and infections in multiple mammalian species. Human cases remain uncommon relative to seasonal influenza, but severe outcomes among confirmed cases and the continuing animal–human interface justify sustained surveillance.

WHO publishes a monthly cumulative register of laboratory-confirmed human A(H5N1) infections. The 7 August 2026 update is explicitly time-stamped and forms the appropriate reference point for the current edition.

Interpretation rule: a confirmed-case CFR is not an infection-fatality ratio. It is affected by who is tested, clinical presentation, ascertainment, healthcare access and reporting practices.

3.1 Human A(H5N1) Mortality

Measure	Current evidence position
Surveillance endpoint	Laboratory-confirmed human A(H5N1) cases reported to WHO
Latest reference date used here	7 August 2026
WHO reporting principle	Member States report sporadic human H5N1 and novel influenza infections under the International Health Regulations
Interpretation	Surveillance counts should not be used as a population infection denominator
Risk assessment	Must integrate clinical severity, exposure route, virology, epidemiology and evidence of human-to-human transmission

3.2 Animal and Wildlife Mortality — Measurement Framework

Domain	What can be measured robustly	What should not be claimed without a standardized denominator
Poultry	Outbreak counts, infected premises, mortality, depopulation/culling	One global annual 'death' total combining biological mortality and culling
Wild birds	Documented die-offs, affected species, geographic spread	A complete global annual wildlife death count
Marine mammals	Confirmed infections and mortality events	A single global marine-mammal mortality denominator
Terrestrial mammals	Species-specific infection and mortality events	A universal mammalian CFR
Livestock	Herd/flock infection, clinical disease, mortality and control measures	Assuming infection always results in death

Animal-control depopulation is an important public-health intervention but is analytically different from biological mortality caused by infection. A defensible global total must keep these endpoints separate.

4. Molecular Biology and Cross-Species Transmission

Mechanism	Biological relevance	Interpretive safeguard
Hemagglutinin receptor interaction	Influences cell attachment and tissue tropism	Receptor preference alone does not establish efficient human transmission
HA cleavage activation	Important for activation and pathogenicity in particular host contexts	Phenotype is context-dependent
Polymerase adaptation	Can alter replication efficiency in mammalian cells	No single mutation is a deterministic 'pandemic switch'
Reassortment	Can create new gene constellations when compatible viruses co-infect a host	Requires ecological opportunity and viable segment combinations
Antigenic drift	Gradual antigenic change that contributes to seasonal vaccine mismatch	Does not automatically create pandemic potential
Antigenic shift	Abrupt antigenic novelty, often involving reassortment	Pandemic spread also requires susceptibility and efficient transmission
Host ecology	Exposure, density, movement, biosecurity and environmental persistence	Requires One Health context

A scientifically defensible risk model should integrate molecular, epidemiological and ecological evidence rather than infer pandemic risk from a single sequence feature.

5. Clinical Presentation, Severity and Treatment

Clinical management must distinguish uncomplicated seasonal influenza from severe disease, high-risk patients and novel influenza A infections. WHO's 2024 clinical practice guideline covers severe and non-severe influenza and includes recommendations on antiviral use for exposed individuals in appropriate circumstances. The guideline also applies to novel influenza A viruses known to cause severe human illness, including zoonotic subtypes such as A(H5N1), A(H5N6) and A(H7N9).

5.1 Practical Clinical Pathway

Stage	Core action	Scientific purpose
Recognition	Assess acute respiratory symptoms, exposure history and risk factors	Early identification of high-risk or zoonotic disease
Triage	Identify hypoxaemia, respiratory distress, altered mental state, shock and other severe features	Prioritize urgent supportive care
Testing	Use validated molecular/antigen methods according to setting; confirm novel/zoonotic infection through appropriate reference pathways	Reduce diagnostic uncertainty
Antiviral management	Use current guideline-based antiviral therapy when indicated; treat high-risk/severe patients promptly	Reduce complications and severe outcomes
Supportive care	Oxygen/respiratory support, hydration and electrolyte management, monitoring and treatment of complications	Maintain organ function
Infection prevention	Appropriate isolation, PPE and occupational exposure assessment	Reduce healthcare-associated transmission
Escalation	Refer/transfer severe or rapidly deteriorating patients to appropriate higher-level care	Prevent avoidable deterioration

5.2 Antivirals and Therapeutics

Therapeutic class	Role	Evidence/implementation principle
Neuraminidase inhibitors	Established antiviral treatment for influenza when indicated	Use according to current clinical guidance, susceptibility and patient risk
Other antiviral classes	May be relevant depending on strain, availability and regulatory status	Resistance and evidence must be assessed for the specific virus
Combination strategies	Research area for severe or novel influenza	Require clinical trials and resistance monitoring
Host-directed approaches	Experimental research area	Must demonstrate clinical benefit and safety
Supportive critical care	Essential for severe disease	Includes oxygenation, hemodynamic support and management of complications

Important: antibiotics do not treat influenza viruses. They are used only when a bacterial co-infection or another bacterial indication is clinically suspected or established.

6. Vaccination and Medical Countermeasures

Vaccination is the principal preventive intervention for seasonal influenza. Because influenza viruses continually evolve, vaccine composition is reviewed and updated twice yearly through WHO's global surveillance and expert consultation process. For the 2026–2027 northern hemisphere season, WHO recommended updated H1N1pdm09, H3N2 and B/Victoria-lineage components, with distinct recommended strains depending on vaccine production platform.

Countermeasure	Function	Preparedness significance
Seasonal vaccines	Reduce risk of influenza illness and severe outcomes	Routine population prevention
Candidate vaccine viruses	Provide a starting point for pandemic vaccine development	Accelerate response to novel threats

Countermeasure	Function	Preparedness significance
Antivirals	Treat or prevent selected influenza infections according to guidance	Clinical and outbreak readiness
Diagnostics	Identify infection and subtype when appropriate	Enable treatment, isolation and surveillance
Monoclonal/novel agents	Potential targeted therapy for selected viral threats	Research and stockpile strategy where evidence supports
Non-pharmaceutical measures	Reduce exposure and transmission	Critical during surges and when countermeasures are limited


7. Diagnostics, Genomics and Global Surveillance

WHO's Global Influenza Surveillance and Response System (GISRS) has operated since 1952 and functions as a global mechanism for surveillance, preparedness and response to seasonal, zoonotic and pandemic influenza. WHO currently describes GISRS as comprising 166 institutions across 136 Member States.

Surveillance layer	Data stream	Decision enabled
Syndromic	Influenza-like illness / severe acute respiratory infection	Detect unusual activity
Laboratory	Molecular and antigenic testing	Confirm infection and characterize virus
Genomic	Sequencing and phylogenetic analysis	Track introductions, evolution and reassortment
Serologic	Antibody studies in defined populations	Estimate exposure and immune response
Animal	Poultry, swine, wild birds and mammal surveillance	Identify spillover interfaces
Environmental	Selected sampling of relevant interfaces	Complement direct host surveillance
Integrated analytics	Human + animal + genomic + contextual data	Support risk assessment and response

WHO's integrated GISRS guidance describes an end-to-end surveillance process from systematic screening and specimen collection through testing, sequencing, data sharing and analysis. Unity Studies provide targeted investigations that supplement routine surveillance when a novel influenza threat emerges.

Operational influenza preparedness cycle



8. Control of Transmission, Outbreak Response and Pandemic Escalation

Influenza control is a dynamic process. The objective is not simply to reduce cases after an outbreak is obvious; it is to detect unusual signals early, characterize the virus, assess transmission and severity, implement proportionate interventions, and continuously update the response.

Control domain	Operational measures	Key outcome
Source control	Animal-health measures, biosecurity, exposure reduction	Lower spillover probability
Early detection	Sentinel surveillance, rapid diagnostics, event-based intelligence	Shorter detection delay
Case management	Prompt triage and evidence-based treatment	Lower severe-outcome risk
Healthcare IPC	Isolation, PPE, ventilation and exposure protocols	Lower healthcare transmission
Contact management	Exposure assessment and monitoring	Early detection of secondary cases
Vaccination	Seasonal vaccination and preparedness vaccination strategies	Lower susceptibility/severity
Risk communication	Timely, transparent, uncertainty-aware messaging	Improved adherence and trust
International coordination	Virus/data sharing and coordinated risk assessment	Faster collective response

The WHO Pandemic Influenza Preparedness Framework is an international instrument designed to improve virus sharing and equitable access to vaccines and other pandemic-related benefits. Its 2024–2030 implementation plan includes monitoring and evaluation of preparedness capacities.

9. Quantitative Epidemiology and Preparedness Formulas

The following equations are analytical tools for researchers and preparedness teams. They are not substitutes for validated epidemiological models.


- Case-fatality ratio:** $CFR = D / C \times 100\%$
- Infection-fatality ratio:** $IFR = D / I \times 100\%$
- Relative risk:** $RR = \text{Risk}_{\text{intervention}} / \text{Risk}_{\text{control}}$
- Relative risk reduction:** $RRR = 1 - RR$
- Vaccine effectiveness:** $VE = (1 - RR) \times 100\%$
- Basic transmission approximation:** $R_t \approx \beta(t) \times c(t) \times D_{\text{inf}} \times S(t)$
- Surveillance anomaly score:** $Z = (x - \mu) / \sigma$
- Excess mortality:** $E = O - B$, where O = observed deaths and B = expected baseline deaths
- State-space preparedness model:** $dx/dt = F[x(t), u(t), \theta] + \varepsilon(t)$; $y(t) = H[x(t)] + \eta(t)$
- Response-delay objective:** $\Delta T = T_{\text{intervention}} - T_{\text{signal}}$

Research principle: Every calculated metric should state its denominator, uncertainty, time window, data source and case definition.

10. The 3-7-12 Systems Framework

3-7-12 INFLUENZA PREPAREDNESS ARCHITECTURE

03 SYSTEM LAYERS



07 INTERVENTION POINTS



12 ACTIONS



Evidence → risk assessment → intervention → measurement → feedback

Status of the framework: This is a proposed conceptual and operational framework authored for research synthesis. It should be independently evaluated rather than presented as an established scientific doctrine.

Layer		Scope
1. Viral		Genome, replication, mutation, reassortment, phenotype and antiviral susceptibility
2. Host		Age, immunity, comorbidity, pregnancy, exposure, clinical severity and healthcare access
3. Environment		Animal reservoirs, density, climate/ecology, mobility, healthcare systems, policy and communication
Intervention point		Purpose
1. Prevention		Reduce exposure and susceptibility
2. Detection		Identify cases and unusual signals early
3. Treatment		Deliver evidence-based clinical care
4. Containment		Interrupt transmission and exposure chains
5. Communication		Provide accurate, uncertainty-aware public information
6. Vaccination		Maintain and deploy effective vaccine protection
7. Research		Generate evidence for improved diagnostics, vaccines and therapeutics
#	Action	Example measurable indicator
1	Integrated surveillance	Signal-to-confirmation time
2	Genomic surveillance	Sequencing coverage of representative samples
3	Risk assessment	Time from characterization to formal risk assessment
4	Rapid diagnostics	Specimen-to-result time
5	Antiviral readiness	Time from triage to appropriate treatment
6	Vaccine readiness	Priority-group coverage / manufacturing readiness
7	One Health coordination	Joint human–animal risk assessments
8	Infection prevention and control	Healthcare-associated transmission rate
9	Data sharing	Time from generation to validated data release

#	Action	Example measurable indicator
10	Community engagement	Reach, trust and adherence indicators
11	Resistance monitoring	Sentinel susceptibility surveillance
12	Simulation exercises	Time-to-action in exercises

11. Advanced Research Agenda for Global Laboratories

Research priority	Question	Validation pathway
Universal / broadly protective vaccines	Can conserved-antigen or novel-platform strategies provide durable cross-subtype protection?	Randomized trials, immunogenicity, durability and clinical endpoints
Rapid diagnostics	Can point-of-care platforms maintain sensitivity across evolving strains?	Analytical validation + prospective clinical studies
Antiviral combinations	Can combination treatment reduce severe outcomes or resistance?	Randomized trials + resistance surveillance
Spillover prediction	Can integrated human-animal-genomic data improve early warning?	Prospective One Health validation
Genomic risk assessment	Which sequence/phenotype associations reproduce across datasets?	Laboratory phenotype + epidemiological validation
Wildlife mortality	Can standardized sampling improve mortality estimates?	Ecological surveillance + denominator standards
3-7-12 evaluation	Does the framework improve measurable preparedness outcomes?	Prospective implementation and comparative evaluation

AI and machine learning can be valuable for anomaly detection, forecasting, genomic classification and resource allocation, but should operate with calibrated uncertainty, human oversight, prospective validation and clear governance. An algorithmic prediction should not be treated as a diagnosis or an autonomous public-health decision.

12. Limitations, Uncertainty and Scientific Safeguards

Issue	Safeguard
Historical mortality estimates	Report ranges and methods rather than false precision
Confirmed-case CFR vs IFR	Never substitute the confirmed-case denominator for the infection denominator
Animal mortality vs culling	Report biological mortality and control depopulation separately
Wildlife surveillance gaps	Use species- and location-specific evidence; do not manufacture global totals
Rapidly changing 2026 data	Time-stamp every dataset and update before publication or policy use
Molecular risk interpretation	Avoid single-mutation deterministic claims
AI prediction	Require uncertainty, external validation and human oversight
Clinical recommendations	Follow current national/international guidance and patient-specific professional assessment

This manuscript is a scientific synthesis and preparedness framework, not individualized medical advice. Clinical decisions should follow current professional guidance, local epidemiology, regulatory requirements and qualified clinical judgment.

13. Conclusion

Influenza should be understood as a continuously evolving biological and public-health system rather than a single disease entity with a fixed mortality rate. Seasonal influenza causes a large annual respiratory mortality burden; historical pandemics demonstrate the capacity for extraordinary mortality under conditions of widespread susceptibility; and zoonotic influenza A viruses remain an important preparedness concern.

Effective control depends on an integrated chain: prevention, early detection, laboratory confirmation, genomic characterization, risk assessment, infection prevention, timely clinical management, vaccination, antiviral readiness, transparent communication and international cooperation. WHO's GISRS and Pandemic Influenza Preparedness Framework provide established international architecture for these functions.

The 3-7-12 Systems Framework is proposed as a compact research and implementation architecture. Its scientific test is empirical: whether it shortens detection and response intervals, improves interoperability, strengthens evidence-based treatment and vaccination delivery, and reduces preventable morbidity and mortality.

14. Selected Primary and High-Authority References

- [1] World Health Organization. Influenza (seasonal). Fact sheet, 28 February 2025. Seasonal burden: approximately one billion cases, 3–5 million severe illnesses, and 290,000–650,000 respiratory deaths annually.
- [2] World Health Organization. Recommended composition of influenza virus vaccines for use in the 2026–2027 northern hemisphere influenza season. 27 February 2026.
- [3] World Health Organization. Cumulative number of confirmed human cases for avian influenza A(H5N1) reported to WHO, 2003–2026, 7 August 2026. Emergency situation update.
- [4] World Health Organization. Clinical practice guidelines for influenza. 12 September 2024.
- [5] World Health Organization. Implementing integrated sentinel surveillance of influenza and other respiratory viruses of epidemic and pandemic potential by GISRS. 4 December 2024.
- [6] World Health Organization. Global Influenza Surveillance and Response System (GISRS). Current programme description.
- [7] World Health Organization. Pandemic Influenza Preparedness Framework. Framework and implementation resources.
- [8] World Health Organization. Framework for health emergency preparedness and response capabilities for national public health agencies. 23 March 2026.
- [9] Centers for Disease Control and Prevention. 2009 H1N1 Pandemic. Global first-year mortality estimate 151,700–575,400.
- [10] World Health Organization. WHO GISRS Unity Studies. Targeted investigations for pandemic preparedness and risk assessment.
- [11] World Health Organization. Candidate vaccine viruses and potency testing reagents for zoonotic influenza. Current candidate-virus programme.
- [12] WHO Global Influenza Programme. Clinical management of influenza. Technical guidance and supporting materials.
- [13] WHO Global Influenza Programme. Recommendations for influenza vaccine composition. Seasonal and zoonotic vaccine recommendations.
- [14] Historical pandemic literature should be interpreted using contemporary demographic reconstruction and model-based uncertainty; exact global totals for 1918, 1957 and 1968 are not treated as census counts in this manuscript.

Appendix A — Data Governance and Update Protocol

For each numerical claim, record: (1) source authority; (2) publication/update date; (3) data cutoff; (4) geographic scope; (5) case definition; (6) denominator; (7) uncertainty class; and (8) version of the dataset.

Recommended update chain: ingest surveillance data → quality control and deduplication → stratify by host and exposure → link laboratory and genomic data → estimate outcomes with uncertainty → conduct risk assessment → update intervention priorities → independently review major changes.

Appendix B — Author and Framework Note

Author: Vahid Nezamivand Chegane
ORCID: 0009-0003-6477-091X
Email: vahidmaykoltarh@yahoo.com

The 3–7–12 Systems Framework in this manuscript is presented as an author-proposed synthesis for research and preparedness applications. It does not imply endorsement, affiliation or authorship by WHO, CDC, Oxford, any university, laboratory or other institution.

This edition intentionally avoids unsupported numerical totals and separates established evidence from conceptual synthesis. The document is designed to support further scientific investigation, peer review and independent validation.

[15] World Health Organization. WHO guideline on public health and social measures for mitigating the risk and impact of epidemic and pandemic influenza. 23 April 2026. Evidence-based recommendations covering personal protection, environmental interventions, case-finding and contact management, social measures, and travel-related actions.