



Math for the people, by the people.

Riemann's theorem on rearrangements

Canonical name	RiemannsTheoremOnRearrangements
Date of creation	2013-03-22 17:31:52
Last modified on	2013-03-22 17:31:52
Owner	Gorkem (3644)
Last modified by	Gorkem (3644)
Numerical id	23
Author	Gorkem (3644)
Entry type	Theorem
Classification	msc 40A05
Synonym	Riemann series theorem
Related topic	UnconditionallyConvergent
Related topic	FiniteChangesInConvergentSeries
Related topic	FiniteChangesInConvergentSeries2

If the map $n \mapsto n'$ is a bijection on \mathbb{N} , we say that the sequence $(a_{n'})$ is a rearrangement of (a_n) .

The following theorem, which is due to Riemann, shows that the convergence of a conditionally convergent series depends so much on the order of its terms; in particular, a conditionally convergent series can be made to converge to any real number by changing the order of its terms.

Theorem (Riemann series theorem). Let (a_n) be a sequence in \mathbb{R} such that $\sum_{n=1}^{\infty} a_n$ converges but $\sum_{n=1}^{\infty} |a_n| = \infty$, i.e., $\sum a_n$ is conditionally convergent. Let $-\infty \leq \alpha < \beta \leq \infty$ be arbitrary. Then there exists a rearrangement $(a_{n'})$ such that

$$\liminf_N \sum_{n'=1}^N a_{n'} = \alpha \quad \text{and} \quad \limsup_N \sum_{n'=1}^N a_{n'} = \beta.$$

Proof. Let $a_n^+ = \max\{0, a_n\}$ and $a_n^- = \min\{0, -a_n\}$. Then we have $a_n = a_n^+ - a_n^-$ and $|a_n| = a_n^+ + a_n^-$. Since $\sum a_n < \infty$, both $\sum a_n^+$ and $\sum a_n^-$ diverge or converge simultaneously. But since $\sum |a_n| = \infty$, we see that at least one of $\sum a_n^+$ and $\sum a_n^-$ must diverge. It follows that $\sum a_n^+ = \infty$ and $\sum a_n^- = \infty$.

Also by the n th term test, $\lim_n a_n^+ = \lim_n a_n^- = 0$.

Now we pass to subsequence of (a_n^+) by removing all terms with $a_n^+ = 0$ and $a_n \neq 0$. For a_n^- , we remove all terms with $a_n^- = 0$. Let us denote the subsequences still as (a_n^+) and (a_n^-) . Since only zeros have been removed, $\sum a_n^+$ and $\sum a_n^-$ are still divergent.

Now we will define integers m_j and k_j for $j \in \mathbb{N}$, and consider the series

$$\begin{aligned} & a_1^+ + a_2^+ + \cdots + a_{m_1}^+ \\ & - a_1^- - a_2^- - \cdots - a_{k_2}^- \\ & + a_{m_1+1}^+ + a_2^+ + \cdots + a_{m_2}^+ \\ & - a_{k_1+1}^- - a_2^- - \cdots - a_{k_2}^- + \\ & \vdots \end{aligned}$$

This series is clearly a rearrangement of $\sum a_n$, by our choice of the subsequences a_n^+ and a_n^- .

We pick up two sequences α_j and β_j such that $\alpha_j \rightarrow \alpha$, $\beta_j \rightarrow \beta$, $\alpha_n \leq \beta_n$ and $\beta_1 > 0$. We choose m_1 such that $\sum_{n=1}^{m_1} \alpha_n \geq \beta_1$ but $\sum_{n=1}^{m_1-1} \alpha_n < \beta_1$. We choose

k_1 such that $\sum_{n=1}^{m_1} - \sum_{n=1}^{k_1} \leq \alpha_1$ but $\sum_{n=1}^{m_1} - \sum_{n=1}^{k_1-1} > \alpha_1$. We continue this way, inductively.

Since $\lim_n a_n^+ = \lim_n a_n^- = 0$, the subsequences of the sequence of partial sums that end with $a_{m_j}^+$ and $a_{k_j}^-$ converge to β and α , and it can be seen that no subsequence can be found with a limit larger than β or lower than α . \square

References

- [1] Rudin, W., *Principles of Mathematical Analysis*, McGraw Hill, 1976.