



planetmath.org

Math for the people, by the people.

Abel's lemma

Canonical name	AbelsLemma
Date of creation	2013-03-22 13:19:49
Last modified on	2013-03-22 13:19:49
Owner	mathcam (2727)
Last modified by	mathcam (2727)
Numerical id	14
Author	mathcam (2727)
Entry type	Theorem
Classification	msc 40A05
Synonym	summation by parts
Synonym	Abel's partial summation
Synonym	Abel's identity
Synonym	Abel's transformation
Related topic	PartialSummation

Theorem 1 Let $\{a_i\}_{i=0}^N$ and $\{b_i\}_{i=0}^N$ be sequences of real (or complex) numbers with $N \geq 0$. For $n = 0, \dots, N$, let A_n be the partial sum $A_n = \sum_{i=0}^n a_i$. Then

$$\sum_{i=0}^N a_i b_i = \sum_{i=0}^{N-1} A_i (b_i - b_{i+1}) + A_N b_N.$$

In the trivial case, when $N = 0$, then sum on the right hand side should be interpreted as identically zero. In other words, if the upper limit is below the lower limit, there is no summation.

An inductive proof can be found <http://planetmath.org/ProofOfAbelsLemmaByInduction> The result can be found in [?] (Exercise 3.3.5).

If the sequences are indexed from M to N , we have the following variant:

Corollary Let $\{a_i\}_{i=M}^N$ and $\{b_i\}_{i=M}^N$ be sequences of real (or complex) numbers with $0 \leq M \leq N$. For $n = M, \dots, N$, let A_n be the partial sum $A_n = \sum_{i=M}^n a_i$. Then

$$\sum_{i=M}^N a_i b_i = \sum_{i=M}^{N-1} A_i (b_i - b_{i+1}) + A_N b_N.$$

Proof. By defining $a_0 = \dots = a_{M-1} = b_0 = \dots = b_{M-1} = 0$, we can apply Theorem 1 to the sequences $\{a_i\}_{i=0}^N$ and $\{b_i\}_{i=0}^N$. \square

References

- [1] R.B. Guenther, L.W. Lee, *Partial Differential Equations of Mathematical Physics and Integral Equations*, Dover Publications, 1988.