



planetmath.org

Math for the people, by the people.

power series

Canonical name	PowerSeries
Date of creation	2013-03-22 12:32:55
Last modified on	2013-03-22 12:32:55
Owner	azdbacks4234 (14155)
Last modified by	azdbacks4234 (14155)
Numerical id	23
Author	azdbacks4234 (14155)
Entry type	Definition
Classification	msc 40A30
Classification	msc 30B10
Related topic	TaylorSeries
Related topic	FormalPowerSeries
Related topic	TermwiseDifferentiation
Related topic	AbelsLimitTheorem
Defines	constant term

A *power series* is a series of the form

$$\sum_{k=0}^{\infty} a_k (x - x_0)^k,$$

with $a_k, x_0 \in \mathbb{R}$ or $\in \mathbb{C}$. The a_k are called the coefficients and x_0 the center of the power series. a_0 is called the *constant term*.

Where it converges the power series defines a function, which can thus be represented by a power series. This is what power series are usually used for. Every power series is convergent at least at $x = x_0$ where it converges to a_0 . In addition it is absolutely and uniformly convergent in the region $\{x \mid |x - x_0| < r\}$, with

$$r = \liminf_{k \rightarrow \infty} \frac{1}{\sqrt[k]{|a_k|}}$$

It is divergent for every x with $|x - x_0| > r$. For $|x - x_0| = r$ no general predictions can be made. If $r = \infty$, the power series converges absolutely and uniformly for every real or complex x . The real number r is called the **radius of convergence** of the power series.

Examples of power series are:

- Taylor series, for example:

$$e^x = \sum_{k=0}^{\infty} \frac{x^k}{k!}.$$

- The geometric series:

$$\frac{1}{1-x} = \sum_{k=0}^{\infty} x^k,$$

with $|x| < 1$.

Power series have some important :

- If a power series converges for a $z_0 \in \mathbb{C}$ then it also converges for all $z \in \mathbb{C}$ with $|z - x_0| < |z_0 - x_0|$.
- Also, if a power series diverges for some $z_0 \in \mathbb{C}$ then it diverges for all $z \in \mathbb{C}$ with $|z - x_0| > |z_0 - x_0|$.

- For $|x - x_0| < r$ Power series can be added by adding coefficients and multiplied in the obvious way:

$$\sum_{k=0}^{\infty} a_k (x-x_0)^k \cdot \sum_{l=0}^{\infty} b_l (x-x_0)^l = a_0 b_0 + (a_0 b_1 + a_1 b_0)(x-x_0) + (a_0 b_2 + a_1 b_1 + a_2 b_0)(x-x_0)^2 \dots$$

- (Uniqueness) If two power series are equal and their domains are the same, then their coefficients must be equal.
- Power series can be termwise differentiated and integrated. These operations keep the radius of convergence.